

# Northumbria Research Link

Citation: Oseni, Zainab, Seedat, Farah and Kandala, Ngianga-Bakwin (2018) HIV Epidemic Heterogeneity in Zimbabwe: Evidence from successive Demographic Health Surveys. *Journal of Biosocial Science*, 50 (6). pp. 840-852. ISSN 0021-9320

Published by: Cambridge University Press

URL: <http://dx.doi.org/10.1017/S0021932017000657>  
<<http://dx.doi.org/10.1017/S0021932017000657>>

This version was downloaded from Northumbria Research Link:  
<http://nrl.northumbria.ac.uk/id/eprint/32502/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



**Northumbria  
University**  
NEWCASTLE



**UniversityLibrary**

*HIV epidemic heterogeneity in Zimbabwe*

## HIV EPIDEMIC HETEROGENEITY IN ZIMBABWE: EVIDENCE FROM SUCCESSIVE DEMOGRAPHIC HEALTH SURVEYS

ZAINAB OSENI\*<sup>1</sup>, FARAH SEEDAT\* AND NGIANGA-BAKWIN KANDALA\*<sup>†‡§</sup>

\**Division of Health Sciences, Warwick Medical School, University of Warwick, Coventry, UK*, †*Department of Mathematics, Physics and Electrical Engineering, Northumbria University, Newcastle upon Tyne, UK*, ‡*Faculty of Health and Sport Sciences, University of Agder, Kristiansand, Norway* and §*Division of Epidemiology and Biostatistics, School of Public Health, University of the Witwatersrand, Johannesburg, South Africa*

<sup>1</sup>Corresponding author. Email: [ngianga-bakwin.kandala@northumbria.ac.uk](mailto:ngianga-bakwin.kandala@northumbria.ac.uk)

**Summary.** Zimbabwe has one of the worst HIV epidemics in the world. This study investigated data from two successive Zimbabwe Demographic and Health Surveys (ZDHS) conducted in 2005–06 and 2010–11. A random representative sample of 30,000 men aged 15–59 and women aged 15–49 was selected from the two surveys. The HIV prevalence was mapped with a flexible, coherent regression framework using a geo-additive semi-parametric mixed model. HIV indicator prevalence maps were constructed at the regional level, and at the administrative level relevant for policy design, planning and decision-making. Substantial regional variation was found, not only in the burden of HIV, but also in its risk factors. The results suggest that responses/policies should vary at the regional level to ensure that the often diverse needs of populations across a country are met and incorporated into planning the HIV response. The use of geographically referenced data in two successive ZDHS provides crucial new insights into the spatial characteristics of the HIV epidemic in Zimbabwe. In particular, it highlights the HIV heterogeneity across Zimbabwe, with substantial regional variation, not only in the burden of HIV, but also in its risk factors.

### Introduction

Zimbabwe, a southern African country with a population of 15,603,000 people, has one of the worst HIV epidemics in the world with an estimated prevalence in 2014 of 16.7% amongst 15- to 49-year-olds (UNAIDS, 2015a; WHO, 2017); this was down from a peak prevalence of 27.7% in 1997 (UNAIDS, 2015b). However, a reduction in prevalence reported nationally could be obscuring the spatial prevalence and the temporal dynamics of the HIV epidemic at a sub-national level. Zimbabwe is made up of 10 regions (Figure 1). Gonese *et al.* (2015) found striking variations in the change of prevalence amongst males and females across different Zimbabwean provinces. Over a 6-year period in Zambia, regions changed from being among the highest HIV prevalence areas to becoming one of the lowest, and vice versa (Kandala *et al.*, 2011). Another study found greater significant declines in HIV prevalence among young urban women compared with young rural women in Zambia (Sandoy *et al.*, 2006). Cuadros and Abu-Raddad (2014) found that areas with low HIV prevalence in several sub-Saharan countries had experienced further sharp declines, thus driving a national reduction in prevalence. However, prevalence has continued to rise or remain steady in some areas of sub-Saharan Africa with high HIV prevalence. Therefore, consideration of the spatio-temporal dynamics of the HIV epidemic at sub-national level is important as this could guide effective and targeted service delivery as well as intervention planning for the future.

The current study aimed to investigate the sub-regional changes in HIV prevalence in Zimbabwe over the 5- to 6-year period between 2005–06 and 2010–11. Thus, the study contributes to the Zimbabwe National HIV and AIDS Strategic Plan II (ZNASPII), which aims to promote the equitable distribution and delivery of services countrywide (National AIDS Council, 2011).

### Methods

#### *Data collection*

Data were from two successive Zimbabwe Demographic and Health Surveys (ZDHS) from 2005–06 and 2010–11. The DHS are funded by the United States Agency for International Development (USAID). They are a well-established source of population-level data with information about knowledge and behaviour regarding HIV/AIDS and other sexually transmitted infections. The objectives, organization, sample design and questionnaires used in the two Zimbabwe DHS are described elsewhere (Central Statistics Office, 2006). In brief, a random sample of men aged 15–59 and women aged 15–49 was selected in both surveys. Three different questionnaires were used to obtain socio-demographic data from participating households. Capillary blood samples were collected on filter paper card for laboratory testing of HIV in each household. The protocol used for collection and analysis of blood specimens was based on an anonymous linked protocol that allows for HIV test results to be merged with an individual's sociodemographic data without identifying the individual concerned. In this study, HIV status was linked to a limited number of socio-demographic variables including respondent's sex, age, place of residence, education level, wealth index and region of residence. (Table 1). The main variable investigated to have an impact on HIV status was the respondent's geographic location, i.e. region of residence at the time of the survey (Fig. 1), in addition to various control variables such as socio-demographic variables known to be associated with HIV status. Age of the respondent at the time of survey was included as an indicator of the birth cohort. Other socio-demographic covariates were sex (female vs male), wealth index (poorest vs poorer, middle, richer and richest), education of the respondent (no education vs primary, secondary and higher education) and place (locality) of residence (rural vs urban).

### *Statistical analysis*

The study applied a unified approach by exploring the spatial patterns in the prevalence of HIV infection at the regional level and possible non-linear effects within a simultaneous, coherent regression framework using a geo-additive semi-parametric mixed model. The model was also able to quantify the spatial autocorrelation of HIV infection in Zimbabwe arising from the population's mobility. The model employed a fully Bayesian approach using Markov Chain Monte Carlo (MCMC) techniques for inference and model checking (Fahrmeir & Lang, 2001; Kandala *et al.*, 2008). The response variable was defined as  $y_i=1$  if HIV sero-positive and  $y_i=0$  otherwise. The standard measure of effect was the Posterior Odds Ratio (POR). Although this estimation process is used less frequently in the literature, the estimated PORs produced can be interpreted in the same way as ordinary logistic models. The model and statistical techniques utilized have been described in more detail elsewhere (Kandala *et al.*, 2011).

The model was implemented using the BayesX (version 2.0.1) software package (Brezger *et al.*, 2005), which permits Bayesian inference, based on MCMC simulation techniques (University of Munich, Munich, Germany). The statistical significance of apparent associations between potential risk factors and the prevalence of HIV was explored using chi-squared and Mann–Whitney *U*-tests, as appropriate. Multivariate analysis was used to evaluate the significance of the Posterior Odds Ratio determined for the fixed non-linear effects and spatial effects.

## **Results**

The distribution of respondents by socio-demographic variables and HIV status in the two successive ZDHS is shown in Tables 1 and 2. The overall HIV prevalence for 2005–06 was 17.9%, and it was higher among females (20.7%) than in males (14.1%). Urban areas had a higher prevalence (18.8%) than rural areas (17.5%), but the difference was not significant in the bivariate analysis. The 15–25 age group had the lowest prevalence (8.5%), and the prevalence rose markedly to 27.9% and 27.7% amongst the 26–35 and 36–49 age groups respectively, before declining to 19.3% in those aged 50 years and over. As education level increased, HIV rates decreased from 21.6% in the 'no education' group to 17.3% in the 'incomplete secondary education' group. Prevalence was 6.7% in people with complete secondary education. Prevalence rose again to 15.3% in people with education higher than secondary school. Wealth index was also a significant predictor of HIV rates, and as wealth index increased so did HIV prevalence, with a rise from 16.9% in the poorest group to 21.7% in the richer group, after which there was a sharp decrease to 15.6% in the richest group.

Matabeleland South (21.1%), Harare (19.3%), Matabeleland North (19.3%), Mashonaland West (19.1%) and Mashonaland East (18%) had the highest prevalences of HIV in 2005, exceeded the national prevalence level. The lowest prevalence was in Mashonaland Central (16.2%) and Midlands (16.3%).

The 2010–11 data show a decline in overall HIV prevalence to 16.4%. Similar to the 2005 data, HIV prevalence was higher in females (18.6%) compared with males (13.4%). However, there was a reduced gap in prevalence between males and females due both to a reduction in the proportion of infected females and a decrease in the proportion of infected males. The lowest prevalence was also noted in the 15–25 age groups (6.6%). This rose to 22.8% in the 26–35 age groups before reaching a peak at 27.5% amongst the 36–49 age groups. Prevalence then declined to 20.4% in the 50 year and over group. In 2010–11, those with no secondary education still had the highest rates of HIV, but individuals with no or incomplete primary education (17.0% and 18.4% respectively) had lower

rates of HIV compared with individuals with complete primary education, who had the highest HIV rate amongst the education groups (19.4%). By 2010–11 the gap between urban and rural areas had widened and reached significance: urban areas (17.9%) had a higher HIV prevalence than rural areas (15.6%). The relationship between wealth and HIV rates had also changed, as the HIV prevalence in the richer group had decreased from 21.7% to 17.4%, whereas the HIV rate in the poorest group had increased from 16.9% to 17.5%, and was now the group with highest HIV risk.

Matebeleland North and Matebeleland South remained areas with the highest HIV prevalence with this staying more or less constant at 21.2% and 19.8% respectively. All other regions except Bulawayo showed a decrease in prevalence. In Bulawayo, there was an increase in prevalence to 19.5%, changing it from one of the lower prevalence areas to one of the highest. Although the Midlands region experienced a decrease in prevalence, it went from being one of the lowest to becoming a relatively high-prevalence area. Harare, which was among the highest prevalence areas in 2005, in 2010–11 had become the lowest prevalence area (14.2%). The HIV prevalences in Matebeleland South, Matebeleland North and Bulawayo were higher than the national average.

Table 3 shows the results of the multivariate analysis using Bayesian geo-additive regression analyses, i.e. adjusting for many confounding factors (including geographic location), indicating that trends in HIV prevalence had changed over time, which is consistent with the unadjusted results. The analysis showed that in 2005–06 HIV rates were highest among females, individuals who were 36–49 years old, those with complete primary education and those in the richer wealth index category. In 2010–11, females, individuals who were 36–49 years old, those with complete primary education, those living in urban areas and in the middle wealth category had the highest rates of HIV. After adjusting for other factors, Matebeleland South and Matebeleland North still had consistently higher HIV prevalence rates across the provinces of Zimbabwe. However, rates across many provinces changed in different directions. The most drastic changes were seen in Harare and Mashonaland West, where HIV rates decreased extensively. Harare became the province with the lowest HIV burden in 2011, while Mashonaland West dropped from the third highest rate to number eight. Conversely, Bulawayo had a higher burden of HIV in 2011 compared with 2005, becoming the province with the third highest HIV rate. Rates in Masvingo and Manicaland remained relatively consistent at moderate HIV levels.

Two important observations emerged from the province-specific net spatial patterns of HIV, which included the total residual spatial patterns of the province (i.e. the sum of the structured and unstructured spatial patterns) (Figure 2 and Figure 3). Statistically significant higher HIV rates of 10% and 13% (POR of 1.10 and 1.13) were found in the western provinces (Matebaland North and South, respectively) in 2005 (Figure 2), and in 2011 (Figure 3) of 26% and 53% (POR 1.26 and 1.53) (Figures 2 and 3). Secondly, trends in the remaining provinces changed over time, from a significant negative association to no association (Figures 2 and 3). Most of the provinces had lower HIV rates in 2005, with the exception of Mashonaland West (POR of 1.04; Figure 2). By 2011, this had changed, and there was no spatial association for any of the remaining provinces, with the exception of Harare, which still had a negative association with HIV rates (POR of 0.69; Figure 3). Over and above the impact of the fixed effects, there appeared to be a higher risk of HIV in the eastern districts (Matebelaland North and South) that was consistent over time.

The estimated non-linear effects of age at HIV diagnosis were plotted as PORs of the risk of HIV against age (Figure 4). Figure 4 shows the PORs together with the 95% point-wise credible intervals. There was a non-linear relationship between the risk of HIV and age during both periods. As expected, in 2005 the likelihood of infection increased with age reaching a peak at around 35 years and then decreasing slightly until 50 years before rising again thereafter (Figure 4, left-hand panel). In 2011, the likelihood of infection was similar and increased with age until 50 years (male only), then it levelled off (for males only) and no longer increased compared with [unlike in?] 2005 (Figure. 4, right-hand panel). It should be pointed out that the age range for females was 15–49 and that for males was 15–59, so the above comparison for age 50 and above only refers to male respondents.

## Discussion

This study investigated inequalities in HIV prevalence at the provincial level in Zimbabwe using Bayesian methods and a nationally representative household sample. The analysis took into account the effects of both individual- and regional-level measures of social inequality in order to gain insight into the influences of socioeconomic, demographic and geographical factors on HIV prevalence in the country.

The results show that HIV rates in Zimbabwe reduced over time to 16.4% in 2010–11, which is higher than the figure of around 15% reported in other studies (Zimbabwe National Statistics Agency, 2012; Glory, 2014). This could be because the estimate was unweighted and included individuals aged over 50, while previous studies only included individuals aged 15–49 years. The marginal difference in reported values is statistically insignificant.

The most striking spatial change found in this study was that Harare in 2005 had the third highest HIV rate of all the Zimbabwean provinces, but by 2010–11 it had the lowest. This could be attributed to the success of interventions in this urban area. Conversely, Bulawayo had the lowest HIV rate in 2005 but by 2010–11 had the

third highest rate. Therefore, the spatial findings from the 2010–11 data generally indicate that HIV rates progressively increased from the south-western regions to the north-eastern regions.

Over time, the highest HIV rates were found in females, the 26–49-year cohort and in individuals with no secondary education. Females were usually at higher risk of HIV due to early age of intercourse, economic factors (such as poverty and exposure to sex trafficking) as well as cultural factors where they may have less power to demand protective measures or refuse sex (Duffy, 2005; Karim *et al.*, 2010). Another possible explanation for the higher risk of HIV observed among females may be the large mucosal surface of the vagina, which is more prone to injury and infection than the penis.

It is vital to tailor HIV prevention efforts to women to reduce the burden of the epidemic. Individuals aged 26–49 had higher rates of HIV compared with other groups, which may be because this is the group that has the resources, opportunity and access to a social network to engage in risky sexual behaviour.

The relationship between wealth index category and HIV rates changed between 2005 and 2010–11; whilst the richer group had higher HIV rates in 2005, in 2010–11 the poorest group had the highest risk. This could be due to the loss of household income and assets due to HIV so that those in the richer wealth category move to the poorer category (Parkhurst, 2010). Previous findings have shown that the decline in HIV incidence has largely been in the top of the wealth index category (Lopman *et al.*, 2007). Those in the poorest wealth index category are also less likely to have had a secondary education and possibly less access to health education and promotion materials. A systematic review found that HIV prevalence falls more consistently among highly educated groups than the less educated, even when overall population prevalence is falling (Hargreaves *et al.*, 2008). Furthermore, people in the lower wealth index group might have fewer available resources to put health promotion recommendations into practice. For instance, transmission of the HIV virus through breast-feeding can be prevented by substituting breast milk with formula milk. However, the poor may not have access to clean water and may be unable to afford formula milk.

At the individual level, people in the richer wealth category were possibly at higher risk of HIV due to opportunity and the social network available for engagement in risky sexual behaviours, whilst people in the poorest group might be at increased risk due to early sexual debut and transactional sex (Fenton, 2004; Shelton *et al.*, 2005; Parkhurst, 2010). Thus, HIV infection risk in the higher wealth category is based on risky behaviour related to wealth, and risk in the poorer group is based on risky behaviour related to poverty (Parkhurst, 2010). Therefore, it is important for policymakers to move beyond the assumption that HIV infection is strictly related to poverty or wealth and instead focus on addressing HIV risk factors and contributing factors unique to each group (Parkhurst, 2010).

Consistent with the situation in other African countries (Dyson, 2003), urban dwellers in Zimbabwe are at higher risk of HIV compared with rural dwellers, and the gap between the two has increased over time. Some studies have shown this to be partly the result of successful interventions in rural compared with urban settings (Orne-Gliemann *et al.*, 2006). Industrialized cities with developed infrastructure encourage migration from rural to urban areas causing higher density of the population and increasing contact and transmission with HIV (Voeten *et al.*, 2010). Migration also increases the level of commercial sex work in urban settings (Bloom & Canning, 2003; Kalipeni *et al.*, 2007).

HIV rates in Zimbabwe over time have remained spatially distributed, with consistently highest rates in Matebeleland South and Matebeleland North. The reasons for the higher rate of HIV infection in specific provinces have not been extensively studied in Zimbabwe. One possibility for the high HIV trends across both Matebeleland provinces is the higher prevalence of poverty in these regions (Zimbabwe National Statistics Agency, 2013). Equally, Harare had the lowest rate of HIV prevalence in 2010–11 and also had lower levels of poverty than other provinces. This possible association between poverty and HIV rates also fits with the findings in this study where in 2010–11 individuals in the poorest wealth categories had the highest rates of HIV. Conversely, Bulawayo had the second lowest rate of poverty yet had one of the highest HIV prevalence rates compared with other regions, and Mashonaland Central had one of the lowest HIV rates in 2005 but is a region with high poverty levels (Sandoy *et al.*, 2006). Another hypothesis is that Matebeland South and North have higher HIV rates as they share borders with the districts of Botswana that have the highest HIV rates (Kandala *et al.*, 2012). Therefore, migration between the areas may increase HIV rates in the Matebeleland provinces.

The major strength of this study was that it used a nationally representative household survey to investigate and explain inequalities in HIV rates accounting for individual, household and regional factors. Furthermore, Bayesian methodology was applied, which quantifies observable and non-observable risk factors of HIV whilst taking into account the auto-correlation in the data that exists between neighbouring geographic locations as people with HIV interact. A limitation of the study is that the data were cross-sectional, which means a causal association between HIV prevalence and its risk factors could not be inferred. Likewise, there was a time lag between exposure to HIV and its detection. Therefore, a risk factor during the time period it was analysed may not necessarily inform the exposure to HIV infection during the earlier years. Finally, other factors that were not included in the model analysis, such as condom use and number of sexual partners, may also contribute to the inequality in HIV prevalence.

This study was an extensive investigation into the spatial variation in HIV infection rates in Zimbabwe in 2005–06 and 2010–11. The study showed a significant geographic pattern of HIV infection, with the highest prevalence being around the south-western provinces of Zimbabwe, especially Matabeleland South and North. There is a need to investigate the reasons why these regions are at greater risk of HIV infection so that they can be targeted with HIV intervention programmes. The HIV distributions provided by this study can be used by policymakers to plan and implement HIV/AIDS interventions across Zimbabwe for the efficient allocation of resources to those at greatest risk.

### Acknowledgments

The authors thank the Demographic and Health Survey Programme (Macro International) for providing the dataset and Caroline Gallimore, the Associate Editor for providing useful comments and suggestions to improve the clarity of the paper.

*Ethical Approval.* The study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki as reflected in a *priori* approval by the institution's human research committee. Ethical approval was granted by the Ethics Committee of the National Statistical Office of Zimbabwe.

*Conflicts of Interest.* The authors have no conflicts of interest to declare.

*Funding.* This study was supported through the DELTAS Africa Initiative SSACAB (Grant No. DEL-15-005). The DELTAS Africa Initiative is an independent funding scheme of the African Academy of Sciences (AAS) Alliance for Accelerating Excellence in Science in Africa (AESA) and supported by the New Partnership for Africa's Development Planning and Coordinating Agency (NEPAD Agency) with funding from the Wellcome Trust (Grant No. 107754/Z/15/Z) and the UK government. The views expressed in this publication are those of the author(s) and not necessarily those of the AAS, NEPAD Agency, Wellcome Trust or the UK government.

### References

- Bloom, D. & Canning, D.** (2003) The health and poverty of nations: from theory to practice. *Journal of Human Development* 4(1), 47–71.
- Brezger, A., Kneib, T. & Lang, S.** (2005) BayesX-Software for Bayesian inference based on Markov chain Monte Carlo simulation techniques. *Journal of Statistical Software* 14(11).
- Central Statistics Office.** (2006) Zimbabwe Demographic and Health Survey 2005–2006. URL: <http://www.dhsprogram.com/pubs/pdf/FR186/FR186.pdf> (accessed 14th November 2017).
- Cuadros, D. F. & Abu-Raddad, L. J.** (2014) Spatial variability in HIV prevalence declines in several countries in sub-Saharan Africa. *Health & Place* 28, 45–49.
- Duffy, L.** (2005) Culture and context of HIV prevention in rural Zimbabwe: the influence of gender inequality. *Journal of Transcultural Nursing* 16(1), 23–31.
- Dyson, T.** (2003) HIV/AIDS and urbanization. *Population and Development Review* 29, 427–442.
- Fahrmeir, L., & Lang, S.** (2001). Bayesian inference for generalized additive mixed models based on Markov random field priors. *Journal of the Royal Statistical Society: Series C (Applied Statistics)* 50(2), 201–220.
- Fenton, L.** (2004) Preventing HIV/AIDS through poverty reduction: the only sustainable solution? *The Lancet* 364, 1186–1187.
- Glory, A.** (2014) Prevalence, spatial patterns and factors associated with HIV infection in Zimbabwe, 2011. Master's thesis, University of Witwatersrand, Johannesburg.
- Gonese, E., Mapako, T., Dzagare, J., Rusakaniko, S., Kilmarx, P. H., Postma, M. J. & Mugurungi, O.** (2015) Within-gender changes in HIV prevalence among adults between 2005/6 and 2010/11 in Zimbabwe. *PLoS One* 10, e0129611.
- Hargreaves, J. R., Bonell, C. P., Boler, T., Boccia, D., Birdthistle, I., Fletcher, A. & Glynn, J. R.** (2008) Systematic review exploring time trends in the association between educational attainment and risk of HIV infection in sub-Saharan Africa. *AIDS* 22(3), 403–414.
- Kalipeni, E., Oppong, J. & Zerai, A.** (2007) HIV/AIDS, gender, agency and empowerment issues in Africa. *Social Science & Medicine* 64, 1015–1018.
- Kandala, N. B., Ji, C., Cappuccio, P. F. & Stones, R. W.** (2008) The epidemiology of HIV infection in Zambia. *AIDS Care* 20(7), 812–819.
- Kandala, N. B., Brodish, P., Buckner, B., Foster, S. & Madise, N.** (2011) Millennium Development Goal 6 and HIV infection in Zambia: what can we learn from successive household surveys? *AIDS* 25(1), 95.
- Kandala, N. B., Campbell, E. K., Rakgoasi, S. D., Madi-Segwagwe, B. C. & Fako, T. T.** (2012) The geography of HIV/AIDS prevalence rates in Botswana. *HIV/AIDS* 4, 95–102.
- Karim, Q. A., Sibeko, S. & Baxter, C.** (2010) Preventing HIV infection in women: a global health imperative. *Clinical Infectious Diseases* 50, S122–S129.
- Lopman, B., Lewis, J., Nyamukapa, C., Mushati, P., Chandiwana, S. & Gregson, S.** (2007) HIV incidence and poverty in Manicaland, Zimbabwe: is HIV becoming a disease of the poor? *AIDS* 21, 57.
- Ministry of Health and Child Welfare** (2009) *Zimbabwe National HIV and AIDS Estimates 2009*
- National AIDS Council** (2011) *Zimbabwe National HIV and AIDS Strategic Plan [ZNASP II] 2011–2015*. [http://www.nationalplanningcycles.org/sites/default/files/country\\_docs/Zimbabwe/znaspi\\_ii\\_aids\\_2011-2015.pdf](http://www.nationalplanningcycles.org/sites/default/files/country_docs/Zimbabwe/znaspi_ii_aids_2011-2015.pdf) (accessed 30th November, 2017)

- Orne-Gliemann, J., Mukotekwa, T., Perez, F., Miller, A., Sakarovitch, C., Glenshaw, M. & Dabis, F. (2006) Improved knowledge and practices among end-users of mother-to-child transmission of HIV prevention services in rural Zimbabwe. *Tropical Medicine & International Health* **11**, 341–349.
- Parkhurst, J. O. (2010) Understanding the correlations between wealth, poverty and human immunodeficiency virus infection in African countries. *Bulletin of the World Health Organization* **88**, 519–526.
- Sandøy, I. F., Kvåle, G., Michelo, C. & Fylkesnes, K. (2006) Antenatal clinic-based HIV prevalence in Zambia: declining trends but sharp local contrasts in young women. *Tropical Medicine & International Health* **11**, 917–928.
- Shelton, J. D., Cassell, M. M. & Adetunji, J. (2005) Is poverty or wealth at the root of HIV? *The Lancet* **366**, 1057–1058.
- UNAIDS (2015a) *How AIDS. Changed Everything. MDG 6: 15 Years, 15 Lessons of Hope from the AIDS Response*. Joint United Nations Programme on HIV/AIDS, Geneva.
- UNAIDS (2015b) *Global Aids Response Progress Report 2015: Zimbabwe Country Report*. URL: [http://www.unaids.org/sites/default/files/country\\_documents/ZWE\\_narrative\\_report\\_2015.pdf](http://www.unaids.org/sites/default/files/country_documents/ZWE_narrative_report_2015.pdf) (accessed 30th November 2017)
- Voeten, H. A., Vissers, D. C., Gregson, S., Zaba, B., White, R. G., de Vlas, S. J. & Habbema, J. D. F. (2010) Strong association between in-migration and HIV prevalence in urban sub-Saharan Africa. *Sexually Transmitted Diseases* **37**(4), 240.
- WHO Zimbabwe (2017) *Country Information; Data and Statistics*. URL: <http://www.afro.who.int/countries/zimbabwe> (accessed 14th September 2017).
- Zimbabwe National Statistics Agency (2012) *Zimbabwe Demographic and Health Survey 2010–11*. URL: <https://dhsprogram.com/pubs/pdf/FR254/FR254.pdf> (accessed 14th September 2017)
- Zimbabwe National Statistics Agency (2013) *Poverty and Poverty Datum Line Analysis in Zimbabwe 2011/12*. URL: <http://www.zimstat.co.zw/sites/default/files/img/publications/Finance/Poverty2011.pdf> (accessed 14th September 2017)

**Fig. 1.** Map of Zimbabwe showing its ten regions.

**Fig. 2.** Left: total residual spatial effects of the risk of HIV/AIDS at region level in Zimbabwe in 2005–06. Shown are the posterior odds ratios. Right: corresponding posterior probabilities at 90% nominal level.

**Fig. 3.** Left: total residual spatial effects of the risk of HIV/AIDS at region level in Zimbabwe in 2010–11. Shown are the posterior odds ratios. Right: corresponding posterior probabilities at 90% nominal level.

**Fig. 4.** Estimated non-linear (logit) effects of respondent's age on the risk of HIV/AIDS at the time of HIV/AIDS diagnosis in 2005–06 (left) and 2010–11 (right). Shown are the posterior logit within the 95% and 80% credible intervals (ZDHS, 2005–06 and 2010–11). Note that for females age ranges from 15 to 49 and for males from 15 to 59.

**Table 1.** Baseline characteristics of the study respondents, ZDHS 2005–06 and 2010–11

Variable	ZDHS 2005–06 (N=13,069)	ZDHS 2010–11 (N=16,651)
Mean age (SD) <sup>a</sup>	27.7 (9.9)	28.4 (9.9)
Sex		
Male	42.6	44.9
Female	57.4	55.1
HIV status (%)		
Yes	17.9	16.4
No	82.1	83.6
Place of residence (%)		
Urban	31.7	35.9
Rural	68.3	64.1
Education (%)		
None	3.1	1.9
Incomplete primary	27.4	10.9
Complete primary	5.1	16.3
Incomplete secondary	59.3	63.3
Complete secondary	1.6	2.2
Higher	3.4	5.3
Wealth Index category (%)		
Poorest	18.9	18.0
Poorer	19.6	17.3
Middle	19.3	18.0
Richer	22.2	22.5
Richest	20.0	24.1
Region of residence (%)		
Manicaland	11.5	11.0
Mahonaland Central	8.1	10.4
Mashonaland East	8.6	9.6
Mashonaland West	9.3	11.1
Matebeleland North	7.6	8.2

---

Variable	ZDHS 2005–06 (N=13,069)	ZDHS 2010–11 (N=16,651)
Matebeleland South	6.5	9.1
Midlands	15.0	11.0
Masvingo	11.3	8.2
Harare	13.5	12.8
Bulawayo	8.6	8.6

---

Data expressed as mean (standard deviations) or as percentages.

<sup>a</sup>Age ranges from 15 to 59 years.

**Table 2.** Baseline characteristics of the study respondents by HIV status, Zimbabwe DHS, 2005–06 and 2010–11

Variable	ZDHS 2005–06			ZDHS 2010–11		
	HIV negative (N=10,714)	HIV positive (N=2335)	<i>p</i> -value <sup>a</sup>	HIV negative (N=8455)	HIV positive (N=5277)	<i>p</i> -value <sup>a</sup>
Age group (%)			<0.001			<0.001
15–25 years	91.5	8.5		93.4	6.6	
26–35 years	72.1	27.9		77.2	22.8	
36–49 years	72.3	27.7		72.5	27.5	
>49 years	80.7	19.3		79.6	20.4	
Place of residence (%)			0.08			0.001
Urban	81.2	18.7		82.1	17.9	
Rural	82.5	17.5		84.3	15.6	
Sex			<0.001			<0.001
Male	85.9	14.1		86.6	13.4	
Female	79.3	20.7		81.4	18.6	
Education (%)			<0.001			<0.001
None	78.3	21.6		83.0	17.0	
Incomplete primary	80.6	19.4		81.6	18.4	
Complete primary	80.4	19.6		80.6	19.4	
Incomplete secondary	82.7	17.3		84.2	15.8	
Complete secondary	93.3	6.7		91.5	8.5	
Higher	84.6	15.3		88.1	11.9	
Wealth index category (%)			<0.001			0.001
Poorest	83.1	16.9		82.5	17.5	
Poorer	82.2	17.8		84.1	15.9	
Middle	83.0	17.0		82.8	17.2	
Richer	78.3	21.7		82.6	17.4	
Richest	84.4	15.6		86.0	14.0	
Region of residence (%)			0.035			<0.001
Manicaland	82.6	17.4		85.1	14.9	
Mahonaland Central	83.8	16.2		85.2	14.8	

Variable	ZDHS 2005–06			ZDHS 2010–11		
	HIV negative	HIV positive	<i>p</i> -value <sup>a</sup>	HIV negative	HIV positive	<i>p</i> -value <sup>a</sup>
	( <i>N</i> =10,714)	( <i>N</i> =2335)		( <i>N</i> =8455)	( <i>N</i> =5277)	
Mashonaland East	82.0	18.0		84.6	15.4	
Mashonaland West	80.8	19.2		84.2	15.8	
Matebeleland North	80.7	19.3		80.2	19.8	
Matebeleland South	78.9	21.1		78.8	21.2	
Midlands	83.6	16.3		84.1	15.9	
Masvingo	83.3	16.7		85.0	15.0	
Harare	80.7	19.3		85.8	14.2	
Bulawayo	82.8	17.2		84.5	19.5	

Data expressed as mean (standard deviation) or as percentages.

<sup>a</sup>*p*-value comparison across HIV-positive and -negative cases using the chi-squared test for categorical variables.

**Table 3.** Unadjusted and fully adjusted odds ratios HIV status cross selected covariates, 2005–06 and 2010–11 ZDHS

Variable	ZDHS 2005–06		ZDHS 2010–11	
	Marginal fully-adjusted OR (95%CI) <sup>a</sup>	Bayesian fully adjusted OR (95% CI) <sup>b</sup>	Marginal fully-adjusted OR (95%CI) <sup>a</sup>	Bayesian fully adjusted OR (95% CI) <sup>b</sup>
<b>Age group (%)</b>				
15–25 years (Ref.)	1.00		1.00	
26–35 years	4.19 (3.73, 4.70)	Figure 4 left	4.32 (3.82, 4.90)	Figure 4 right
36–49 years	4.37 (3.84, 4.97)		5.72 (5.02, 6.52)	
>49 years	3.54 (2.52, 4.97)		5.13 (3.77,6.98)	
<b>Place of residence (%)</b>				
Urban	1.29 (1.05, 1.60)	1.23 (1.02, 1.48)	1.97 (1.66, 2.34)	1.87 (1.60, 2.19)
Rural (Ref.)	1.00	1.00	1.00	1.00
<b>Sex</b>				
Female	1.58 (1.43, 1.75)	1.51 (1.36, 1.68)	1.46 (1.32, 1.62)	1.44 (1.31, 1.60)
Male (Ref.)	1.00	1.00	1.00	1.00
<b>Education (%)</b>				
No education	1.05 (0.72, 1.53)	1.28 (0.94, 1.91)	0.98 (0.64, 1.50)	1.06 (0.71, 1.52)
Incomplete primary	1.41 (1.05, 1.89)	1.67 (1.26, 2.32)	1.58 (1.19, 2.11)	1.69 (1.12, 2.17)
Complete primary	1.63 (1.16, 2.30)	1.93 (1.30, 2.63)	1.72 (1.31, 2.26)	1.79 (1.38, 2.27)
Incomplete secondary	1.53 (1.16, 2.02)	1.70 (1.31, 2.31)	1.68 (1.31, 2.16)	1.74 (1.44, 2.24)
Complete secondary	0.76 (0.41, 1.41)	0.74 (0.44, 1.26)	1.11 (0.67, 1.84)	1.01 (0.52, 1.57)
Higher education (Ref.)	1.00	1.00	1.00	1.00
<b>Wealth index category (%)</b>				
Poorest	1.25 (0.98, 1.59)	1.18 (0.93, 1.47)	1.66 (1.34, 2.05)	1.60 (1.34, 1.96)
Poorer	1.39 (1.10, 1.75)	1.33 (1.07, 1.63)	1.63 (1.33, 2.00)	1.61 (1.33, 2.02)
Middle	1.38 (1.10, 1.73)	1.35 (1.11, 1.76)	1.75 (1.45, 2.12)	1.73 (1.43, 2.09)
Richer	1.60 (1.37, 1.88)	1.54 (1.32, 1.87)	1.36 (1.17, 1.59)	1.33 (1.17, 1.54)
Richest	1.00	1.00	1.00	1.00
<b>Region of residence (%)</b>				
Manicaland	1.03 (0.82, 1.28)	Figure 2	0.93 (0.76, 1.15)	Figure 3
Mahonaland Central (Ref.)	1.00	0.95 (0.79, 1.07)	1.00	0.94 (0.82, 1.16)

Variable	ZDHS 2005–06		ZDHS 2010–11	
	Marginal fully-adjusted OR (95% CI) <sup>a</sup>	Bayesian fully-adjusted OR (95% CI) <sup>b</sup>	Marginal fully-adjusted OR (95% CI) <sup>a</sup>	Bayesian fully-adjusted OR (95% CI) <sup>b</sup>
Mashonaland East	1.05 (0.83, 1.33)	0.97 (0.88, 1.06)	1.02 (0.82, 1.26)	0.95 (0.82, 1.13)
Mashonaland West	1.20 (0.96, 1.51)	1.04 (0.92, 1.19)	0.95 (0.78, 1.17)	0.93 (0.79, 1.05)
Matebeleland North	1.28 (1.01, 1.63)	1.10 (1.00, 1.27)	1.32 (1.07, 1.65)	1.26 (1.07, 1.51)
Matebeleland South	1.36 (1.06, 1.73)	1.13 (1.02, 1.30)	1.64 (1.33, 2.01)	1.53 (1.26, 1.84)
Midlands	1.00 (0.81, 1.24)	0.94 (0.86, 1.01)	1.00 (0.82, 1.23)	0.99 (0.87, 1.12)
Masvingo	1.04 (0.84, 1.30)	0.97 (0.86, 1.08)	0.96 (0.77, 1.20)	0.92 (0.76, 1.12)
Harare	1.13 (0.87, 1.46)	1.02 (0.89, 1.19)	0.67 (0.53, 0.85)	0.69 (0.58, 0.81)
Bulawayo	1.00 (0.75, 1.32)	0.96 (0.81, 1.12)	1.11 (0.85, 1.44)	1.12 (0.91, 1.36)

Rank of marginal OR in 2005–06: Matebeleland South (1), Matebeleland North (2), Mashonaland West (3), Harare (4), Mashonaland East (5), Masvingo (6), Manicaland (7), Midlands (8), (9), Mahonaland Central (10).

Rank of marginal OR in 2010–11: Matebeleland South (1), Matebeleland North (2), Bulawayo (3), Mashonaland East (4), Midlands (5), Mahonaland Central (6), Masvingo (7), Mashonaland West (8), Manicaland (9), Harare (10).

<sup>a</sup>Marginal fully adjusted odds ratio (OR) from standard logistic regression models and ranked from the highest to the lowest OR. Mahonaland Central is taken as reference because of its low HIV prevalence in both 2005 and 2011.

<sup>b</sup>Spatially adjusted posterior odds ratio (POR) obtained from the Bayesian geo-additive regression mixed model after controlling for non-linear effect of age, categorical variables and the region of residence (spatial effects: see also Figs 2 and 3).

Ref., reference category.