

# Northumbria Research Link

Citation: Klonizakis, Markos, Gumber, Anil, McIntosh, Emma, King, Brenda, Middleton, Geoff, Michaels, Jonathan and Tew, Garry (2018) Exercise fidelity and progression in a supervised exercise programme for adults with venous leg ulcers. *International Wound Journal*, 15 (5). pp. 822-828. ISSN 1742-4801

Published by: Wiley

URL: <http://doi.org/10.1111/iwj.12933> <<http://doi.org/10.1111/iwj.12933>>

This version was downloaded from Northumbria Research Link:  
<http://nrl.northumbria.ac.uk/id/eprint/33979/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

## **Exercise Fidelity and Progression in a Supervised Exercise Programme for Adults with Venous Leg Ulcers.**

Markos Klonizakis DPhil<sup>1</sup>, Anil Gumber PhD<sup>2</sup>, Emma McIntosh Msc<sup>1</sup>, Brenda King Msc<sup>3</sup>, Geoff Middleton Msc<sup>4</sup>, Jonathan A. Michaels MChir<sup>5</sup>, Garry A. Tew PhD<sup>6</sup>.

<sup>1</sup>Centre for Sport and Exercise Science, Sheffield Hallam University, Sheffield, United Kingdom, S10 2BP

<sup>2</sup>Department of Allied Health Professions, Faculty of Health and Wellbeing, Sheffield Hallam University, Sheffield, United Kingdom, S10 2BP

<sup>3</sup>Manor Clinic, Sheffield Teaching Hospitals NHS Foundation Trust, Sheffield, United Kingdom, S12 2ST

<sup>4</sup>School of Sport and Exercise Science, University of Lincoln, Brayford Pool, Lincoln, Lincolnshire LN6 7TS

<sup>5</sup>School of Health and Related Research, University of Sheffield, Sheffield, United Kingdom, S1 4DA.

<sup>6</sup>Department of Sport, Exercise and Rehabilitation, Northumbria University, Newcastle-upon-Tyne, United Kingdom, NE1 8ST

\*Correspondence and requests for reprints to:

Dr Markos Klonizakis, Centre for Sport and Exercise Science, Sheffield Hallam University, Sheffield, S10 2BP. Email: [M.Klonizakis@shu.ac.uk](mailto:M.Klonizakis@shu.ac.uk) Tel: +44(0)1142252465

Keywords: venous ulcers, aerobic exercise, intervention fidelity, exercise progression, safety

## **Abstract**

Purpose of investigation: Despite exercise being included in the recommended advice for patients with venous leg ulcers, there is a fear shared by clinicians and patients that exercise may be either inappropriate or harmful and actually delay rather than promote healing.

Therefore, before implementing a larger scale study, exploring the effects of a supervised exercise programme in patients with venous ulcers being treated with compression therapy, it is important to assess exercise safety as well as fidelity and progression in a feasibility study.

Methods: Eighteen participants randomised in the exercise group were asked to undertake 36 (3 times/week for 12 weeks), 60-minute exercise sessions, each comprising moderate-intensity aerobic, resistance and flexibility exercise components. For the purposes of this paper we analysed the data collected during the exercise sessions.

Results: The overall session attendance rate was 79%, with 13/18 participants completing all sessions. No in-session adverse events were reported. 100% aerobic components and 91% of resistance components were completed within the desired moderate-intensity target. Similarly, 81% of aerobic components and 93% of flexibility components were completed within the prescribed duration targets.

Conclusions: Our data showed that patients with venous ulcers could safely follow a supervised exercise programme incorporating moderate-intensity aerobic, resistance and flexibility components.

## Introduction

Venous leg ulceration is a chronic and devastating condition that affects approximately 1% of the adult population in the Western world (1). It costs up to 198 million sterling pounds in national healthcare expenditure in the U.K. alone (2), affecting significantly, in a negative manner patients' quality of life (3). Moreover, venous leg ulcers tend to recur quite frequently, with recurrence rates reaching 70% within a year of healing (4).

With such costs involved and the considerable devastation in patients' lives, it is no surprise that adjunct and alternative therapies to compression therapy (which is considered as the golden standard) (5) have been pursued (e.g. ultrasound (6), larval therapy (7), biomaterials (8)), with exercise and physical activity promotion being considered as well (e.g., walking (9), increased physical activity (10), resistance exercise (11)).

The concept of using exercise as an adjunct therapy to compression isn't new and indeed exercise is included as a recommendation in the NICE Clinical Knowledge Summary for venous leg ulcers' management (e.g., "regular walking", "exercising to improve calf muscle pump function") (12). Nevertheless, there is a fear shared by both clinicians and the patients that exercise may be either inappropriate or harmful and actually delay rather than promote healing (13,14). This notion together with the mixed results of previous studies (13,15-16), has limited the exploration of regimes that could potentially benefit patients and improve clinical outcomes. Overall, there is little published data on the ability of this patient group to undertake different types of exercise training and on rates of exercise progression. The data has the potential to inform practitioners and researchers involved in prescribing and supervising exercise with venous ulcer patients. .

FISCU (17) is a recently-completed, two-center study exploring the feasibility of using exercise as an adjunct therapy to compression in patients with venous leg ulcers. This trial represents an attempt to implement a supervised exercise programme with this patient population, in a manner similar to what has been promoted successfully in other clinical populations in the UK (e.g. peripheral arterial disease (18), chronic obstructive pulmonary disease (19), cardiac diseases (20)). Central to the internal validity of all intervention trials is intervention fidelity, which refers to the extent an experimental manipulation has been implemented in a comparable manner to all participants, as intended (21). Furthermore, it is important to present all in-session exercise safety data to better inform clinicians, policy makers and patients with venous ulcers. As such, having published the main study findings, which supported the feasibility of conducting a future full-scale trial (17), our aim here was to present a detailed appraisal of exercise data collected during the FISCU trial, focussing on treatment fidelity and exercise progression.

## **Methods**

FISCU was a two-arm, parallel-group, randomised feasibility trial that received ethical clearance from the NHS National Research Ethics Committee for Yorkshire and the Humber (14/YH/0091), and was prospectively registered (ISRCTN09433624). Thirty-eight adults who were receiving lower-limb compression for a new venous leg ulcer of greater than 1 cm diameter were recruited from tissue viability clinics and newspaper advertisement in Sheffield, United Kingdom. Following provision of consent and baseline assessment, participants were randomly assigned to receive usual care (n=20) or usual care plus a 12-week supervised exercise programme (n=18). A full description of the protocol is available

elsewhere (22); however, for the purpose of this article the exercise training protocol is described below.

### **Exercise protocol**

Following study enrolment and randomisation, exercise group participants were referred for a 12-week exercise intervention, undertaken 3 times per week (typically being delivered on Mondays, Wednesdays and Fridays to allow sufficient recovery between sessions). A maximum of an additional 2 weeks was allowed for the participants to complete the 36 sessions in case sessions were missed because of illness, family/work commitments or holiday. The sessions were supervised by an exercise physiologist and were typically undertaken in a group form (no more than 4 patients per session, to ensure proper supervision and adequate progression monitoring). Each exercise session lasted approximately 60 minutes and comprised a combination of aerobic, resistance and flexibility exercises. Each session began and ended with 5 minutes of low-intensity treadmill walking or cycling for a warm-up and cool-down, respectively. The aerobic component was aimed to last approximately 30 minutes, with the exercise mode being treadmill walking, cycling, or a combination of both, with the mode being determined by the physical function and preference of participants.

Resistance and flexibility exercises were performed for approximately 20 minutes in order to improve calf muscle pump function, leg (predominantly calf) muscle strength, and joint (predominantly ankle) mobility. Resistance exercises mainly involved dynamic body-weight exercises with or without the use of dumbbells and stability balls (e.g., calf raises and partial

squats). Exercise was aimed to be performed for two or three sets of 10 to 15 repetitions to the point of moderate muscle fatigue (23). For flexibility, static stretches were performed for all of the major muscle groups of the legs, for a total of 60 seconds per muscle group (comprising 3 × 20-second stretches), held at the point of mild discomfort (23).

### **Exercise intensity: prescription and measurement**

The intensity of aerobic and resistance exercises was guided using Borg's 6-20 ratings of perceived exertion (RPE) scale (24), aiming for an exertion level of 12 to 14 ("moderate" to "somewhat hard") on the 6-20 scale, which equates to the ventilatory threshold (24). Each patient was familiarized with the scale and the recommended researcher instructions for scale administration were used (25). Perceived exertion, heart rate (via telemetry; Polar RS400, Kempele, Finland), and aerobic and resistance exercise indices (e.g., treadmill speed and gradient) were recorded at regular intervals during the whole session to allow accurate quantification of the exercise stimulus and to facilitate progression of the programme over time.

### **Exercise safety**

Compression garments (stockings/bandages) were monitored throughout each exercise session. The exercise supervisor was instructed to terminate the session if these were affected by exercise, with participants being referred to the tissue-viability nursing team for re-application, and additional visits were to be noted for the health-economics analyses. Our safety monitoring procedure indicated that all serious adverse events, as well as all non-

serious adverse events that are deemed to be related to participation in the research (e.g., exercise strains or injuries, excessive wound discharge, in-session exercise bandage slipping) were to be recorded during the period between provision of informed consent through to 12 months after randomisation. Participants were asked to contact the study team to inform them about adverse events if and when they occur. Study investigators also questioned participants about the occurrence of adverse events during each participant study visit.

### **Statistical analysis**

Descriptive statistics were used to calculate the session attendance data, completion rates as per protocol for aerobic (duration, intensity, combination of duration and intensity), resistance (intensity, number of exercises, sets, repetitions) and flexibility exercises (number of exercises, duration, intensity), and present baseline demographics. The Shapiro-Wilk test was used to assess data normality and Mauchly's Test of Sphericity was used to indicate data sphericity (the assumption of sphericity was not violated in any case). Exercise progression was assessed by comparing Session 1 (baseline), with Sessions 18 (midpoint) and 36 (intervention completion) using Analysis of Variance (ANOVA) for Repeated Measures (SPSS v.23, Armonk, NY: IBM Corp). Post-hoc analysis was undertaken using Bonferroni corrected t-tests. To calculate the effect sizes we used eta-square for ANOVA assessments and Cohen's d for post-hoc analysis, using the magnitudes determined by Cohen (26): For  $\eta^2$  0.01 is considered a small effect, 0.06 is considered a medium effect and 0.14 is considered a large effect. Similarly for Cohen's d: 0.2 is considered a small effect, 0.5 is considered a medium effect and 0.8 is considered a large effect. Data are described as means (SD), unless otherwise stated. Significance set at  $p < 0.05$  and for post hoc analysis at  $p < 0.0167$ .

## Results

### *Participants*

Characteristics of the 18 exercise-group participants are shown in Table 1. Ten of these participants were female and the mean  $\pm$  SD age, stature and body mass were  $66.9 \pm 13.9$  years,  $171.1 \pm 11.9$  cm and  $102.1 \pm 29.4$  kg, respectively. Median ulcer size was  $4.9 \text{ cm}^2$ .

### *Attendance*

The overall exercise attendance rate was 79% (512/648), with 13 of the 18 participants (72%) attending all exercise sessions. Amongst those who completed the study, 411/468 sessions were completed within the 12-week period, with the rest (57/468) being completed within the additional 2-week period. Of the five participants who did not complete all sessions, one withdrew fully from the trial before the 3-month follow-up assessment due to non-ulcer-related health problems, and four withdrew from treatment (i.e. stopped attending before the end of the 12-week intervention period) but remained in the study (one due to ulcer-related problems, three due to non-ulcer-related health problems). These five participants had completed 2, 4, 6, 15 and 17 exercise sessions, respectively, before withdrawing. Reasons for not attendance included lack of transportation (n=34), non-ulcer related health reasons (n=74) and ulcer-related health reasons (n=32), with more than one reasons given on some occasions.

### *Exercise Safety*

No serious, in-session adverse events were experienced and the bandaging was also not disrupted during any exercise session. Two incidents of excessive fluid discharge were detected the day after exercise sessions, possibly or probably related to exercise. Following consultations with healthcare personnel, these were dealt by postponing the exercise session following the incident reporting (incident 1) and temporarily removing the resistance element from the training programme (incident 2).

### *Exercise choices*

The majority of the participants (72%) chose treadmill as their main aerobic mode of training at baseline, with the rest preferring cycling due to frailty and lack of confidence with exercising on the treadmill. One participant changed briefly from treadmill to exercise cycle, before reverting to treadmill again. Only one of the participants was training via exercise cycle at the end of the 12-week intervention, with the rest of the participants that completed the intervention using the treadmill instead.

In regards to the resistance element of the intervention, four participants started the programme stating that they were unable to do squats, step-ups or calf raises. This number was reduced to two at the end of their participation (they were however, able to complete the rest of the regime). Finally, one of the participants could not do squats on Session 18, due to a pre-existing pain unrelated to exercise, completing however, the rest of the session without issues. The participant completed his programme as well.

### *Exercise Intensity*

All of the aerobic and 91% of the resistance training components, across all participants, were performed at the desired moderate intensity, as determined using RPE responses in the 12-14 range (Tables 2 and 4).

### *Exercise Progression*

Table 3 presents data on changes in the duration of the aerobic component and the number of repetitions completed for four lower-limb resistance exercises. The number of minutes spent on aerobic exercise increased through the 12-week period (Baseline: 19 min (8), Mid-point: 26 min (5), End-point: 29 min (3)).

Performance of the participants in the resistance exercise indices was also improved (Table 3): For example, calf raises increased from 19 (13) at baseline, to 36 (13) at mid-point, reaching 42 (14) at the end of the intervention.

### *Exercise Fidelity*

For the aerobic exercise element all completed sessions were completed according to the prescribed intensity. For resistance this was the case in 466/512 (91%) completed sessions. Duration of the exercise elements was close to the prescribed duration as well (413/512 = 81% for aerobic, 474/512 = 93% for resistance). The majority of those not completing the prescribed duration were at the beginning of their programme, and was due to lack of

physical fitness (n=4), discomfort (n=2) and unfamiliarity with the training equipment/exercises (n=4) – with more than one reason being given by some participants.

Similarly, the main reason for resistance components not being completed according to protocol was lack of physical fitness. This, however, became less of an issue as the programme progressed, reaching almost 100% completion in the last sessions.

Finally, flexibility exercises were completed as per protocol in regards to duration and number of exercises.

## **Discussion**

Using a supervised exercise regime as an adjunct therapy to reduce venous leg ulcer healing time, represents a plausible, yet largely unassessed therapeutic strategy (16). The lack of appropriately designed studies, which would substantiate its use and the fear of healthcare professionals and the patients themselves about the safety and applicability of exercise are two main reasons, why the advice of a more "active lifestyle" is not being taken up more widely within this patient population (13,14).

We have recently presented data supporting the feasibility of a full-scale trial of adjunctive exercise therapy for venous leg ulceration (17). The aim of the current paper was to undertake a detailed evaluation of the exercise session data. When adhering to pre-determined safety criteria, our results show primarily a very high fidelity of our proposed programme. It is evident from our data that not only is it possible to exercise this primarily-older and largely-frail, patient population at moderate intensities, but it is also possible to

see a positive exercise progression over the duration of a medium-term training programme. This is the first study to report in-session data on this patient group and this acts as a comparator for researchers and practitioners embarking on similar trials with exercise as a therapy with this patient group.

#### *Attendance, compliance, and safety*

Our overall session attendance (79%) for the 18 participants across the 12-week exercise intervention compares well with an attendance range of 58–77% for other exercising, clinical populations (27,28). Our attendance results can be interpreted even more favourably to those achieved in other exercise studies, considering the fact that ours was a time-demanding (e.g. 3 times per week), 3-month intervention, focusing on a group which is older, sedentary and without an exercising culture; the large majority of our participants have not previously followed an exercise programme. Consequently, it can be postulated that our participants were keen to embrace such an intervention and participated wholeheartedly. Our results also show that most missed sessions can be accounted to reasons unrelated to the exercise programme (e.g. illnesses and family commitments) rather than the exercise programme itself. This knowledge, combined with the very good safety record (e.g., no participants had their compression garments affected during the exercise sessions), is a sign of trust of moving the intervention into the next stage, that of the definitive trial. Nevertheless, much more data is required to evaluate the safety of the intervention properly in this patient group.

---

When evaluating the fidelity of exercise training interventions researchers should ideally consider both session attendance and meeting the prescribed exercise intensity, as this interaction constitutes the dose of the intervention and influences the physiological response to exercise training (21). Although in our case this might have been considered as a relatively easy task (as our aim was to have participants exercising at "moderate" intensity, e.g. 12-14 in the 6-20 RPE Borg scale), which is considerably lower to that sought by high intensity training (e.g., 85–95% of peak heart rate) (29) exercise regimes, results should not be overlooked: our participants' unfamiliarity with exercise interventions and in some cases frailty, meant that even the intended moderate intensity could potentially be difficult to achieve in practice. For the aerobic exercise element of our intervention this was achieved and maintained throughout the duration of the intervention, matching the performance of other regimes, conducted in clinical settings, in older clinical populations (e.g. Alzheimer's Disease) (30). Results differ in regards to resistance and flexibility, as certain participants found difficulty to complete all resistance exercises to the required level (Table 3) or intensity (Table 4). This was mainly due to frailty and lack of physical fitness (number of sets/repetitions for resistance) or patients finding the exercises easier than expected (intensity for flexibility). This can only be considered as part of our learning process to introduce more challenging exercises (for flexibility) and a varying introductory pace (for resistance), in the future study stages.

### *Exercise progression*

---

The main aim of this article was to present our findings on attendance, compliance and safety. Nevertheless, our detailed collection and analysis of exercise training data permits the objective appraisal of our regime in regards to exercise progression as well: To facilitate a positive adaptation to training, the prescription of exercise needs to advance over time (27). Many programmes have failed to achieve this, presenting a need to re-define targets, following an in-programme assessment (31) (which can be costly and resource-intensive). In the study presented in this article, we used relative measures of exercise intensity to assess adherence to the prescribed intensity. The fact that our aim was achieved was reflected in all of our exercise indices, which show a statistically-significant increase in most measures, as well as a moderate-to-large effect sizes: this demonstrates a clear exercise progression. Although it is difficult to compare our findings with that of other trials in clinical or older populations (as in-session data is not usually reported), our data is equally- or more favourably- comparable to similar interventions in other clinical populations where physical functioning indices appear to be reduced (e.g. chronic kidney disease) (32) or improved (e.g. older people living in retirement communities (33), multiple sclerosis (34)) when compared with normative values. It remains to see whether this exercise progression will be achievable in the definitive trial as well, nevertheless, the indicators are encouraging, suggesting that participants with venous ulcers can benefit in multiple ways (e.g. improved cardiorespiratory endurance (35) and better physical function (27), which are related to high exercise session attendance) by taking part in such an intervention combining medium-intensity aerobic, flexibility and resistance exercise, as previous studies in clinical populations have shown (36,37).

### *Limitations*

With this study exploring the feasibility of the intervention, the number of participants was relatively small to what the definitive trial is expected to include. With that in mind findings should be treated as indicative. Additionally, an in-depth assessment of fidelity in a definitive, multi-centre exercise intervention will examine the consistency of the exercise dose across the different sites, something that was not possible on this occasion. Finally, we cannot rule out the possibility of underreported RPE scores due to the influence of observer sex as it has been suggested that male participants report lower RPE values when a female observer, as opposed to male, is in the room (38). Nevertheless, our sessions were delivered by both male and female physiologists and our findings appear to be consistent throughout the intervention, hence the likelihood of that is small.

### *Conclusions*

This is the first study to provide a detailed quantification of the exercise sessions performed across an exercise intervention combining aerobic, resistance and flexibility exercises for patients with venous ulcers. The data will act as a comparator for researchers embarking on similar trials and advocating exercise to this patient group in their practice. Our findings showed that our participants trained at the intended exercise intensity, improving their performance amongst all exercise domains in which they trained (e.g., number of minutes in aerobic exercise, number of squats and calf raises etc), without having their safety compromised. We conclude that it is possible to exercise this patient population at moderate exercise intensities. This is purposeful for further studies which consider

deploying similar supervised exercise regimes as an adjunct therapy to compression, in an attempt to reduce healing times in patients with venous ulcers.

---

**Key points:**

- Supervised exercise has been used successfully in many clinical populations. Nevertheless – and despite exercise being included in the clinical recommendations for people with venous ulcers (VLUs), it has never been tested in this clinical population.
- We aimed to evaluate the fidelity and exercise progression of a supervised exercise programme in patients with venous ulcers being treated with compression therapy. Eighteen people with VLUs were asked to attend 36 sessions of supervised exercise consisting of aerobic, resistance and flexibility exercises, within a space of 12 weeks.
- The overall session attendance rate was 79%, with 13/18 participants completing all 36 sessions, while no in-session adverse events were reported. The vast majority of aerobic and resistance exercise components were completed within the desired moderate-intensity targets and within the prescribed duration targets.

**Acknowledgements**

The FISCU study was funded by the National Institute for Health Research (NIHR) Research for Patient Benefit Programme (grant PB-PG-0213-30029). This funding source had no role in the design of this study and will not have any role during its execution, analyses and

interpretation of the data. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

### Conflicts of Interest

None to declare.

### References

---

1. Donnelly R, London NJ. The Ulcerated lower limb. In: Donnelly R, London N. NJ eds., *ABC of Arterial and Venous Disease*. 2<sup>nd</sup> ed. Chichester, United Kingdom; Wiley-Blackwell; 2009:77-83.
2. Posnett J, Franks PJ. The burden of chronic wounds in the UK. *Nurs Times*. 2008;**104**:44-45.
3. Carradice D, Mazari FA, Samuel N, Allgar V, Hatfield J, Chetter IC. Modelling the effect of venous disease on quality of life. *Br J Surg*. 2011;**98**:1089-1098.
4. Barwell JR, Davies CE, Deacon J, Harvey K, Minor J, Sassano A, Taylor M, Usher J, Wakely C, Earnshaw JJ, Heather BP. Comparison of surgery and compression with compression alone in chronic venous ulceration (ESCHAR study): randomised controlled trial. *Lancet*. 2004;**363**:1854-1859.
5. O'Meara S, Cullum N, Nelson EA, Dumville JC. Compression for venous leg ulcers. *Cochrane Database Syst Rev*. 2012;**11**:CD000265.

6. Watson JM, Kang'ombe AR, Soares MO, et al. Use of weekly, low dose, high frequency ultrasound for hard to heal venous leg ulcers: the VenUS III randomised controlled trial. *BMJ*. 2011;**342**:d1092-1101.
7. Dumville JC, Worthy G, Bland JM, Cullum N, Dowson C, Iglesias C, Mitchell JL, Nelson EA, Soares MO, Torgerson DJ. Larval therapy for leg ulcers (VenUS II): randomised controlled trial. *BMJ*. 2009;**338**:b773-780.
8. Mostow EN, Haraway GD, Dalsing M, Hodde JP, King D, OASIS Venous Ulcer Study Group. Effectiveness of an extracellular matrix graft (OASIS Wound Matrix) in the treatment of chronic leg ulcers: a randomized clinical trial. *Journal of vascular surgery. J Vasc Surg*. 2005;**41**:837-843.
9. Meagher H, Ryan D, Clarke-Moloney M, O'Laighin G, Grace PA. An experimental study of prescribed walking in the management of venous leg ulcers. *J Wound Care*. 2012;**21**:421-430.
10. Heinen MM, Evers AW, Van Uden CJ, Van der Vleuten CJ, van de Kerkhof PC, Van Achterberg T. Sedentary patients with venous or mixed leg ulcers: determinants of physical activity. *J Adv Nurs*. 2007;**60**:50-57.
11. O'Brien J, Finlayson K, Kerr G, Edwards H. Evaluating the effectiveness of a self-management exercise intervention on wound healing, functional ability and health-related quality of life outcomes in adults with venous leg ulcers: a randomised controlled trial. *Int Wound J*. 2017;**14**:130-137.
12. Clinical Knowledge Summaries: Leg ulcer – venous, National Institute for Health and Clinical Excellence <https://cks.nice.org.uk/leg-ulcer-venous#!topicsummary>. Accessed Jan 25, 2018.

13. O'Brien J, Finlayson K, Kerr G, Edwards H. The perspectives of adults with venous leg ulcers on exercise: an exploratory study. *J Wound Care*. 2014;**23**:496-509.
14. McCulloch J, Mahoney E, McCallon S. Enhancing the role of physical therapy in venous leg ulcer management. *JAMA Dermatol*. 2015;**151**:327-327. doi: 10.1001/jamadermatol.2014.4042
15. Jull A, Parag V, Walker N, Maddison R, Kerse N, Johns T. The prepare pilot RCT of home-based progressive resistance exercises for venous leg ulcers. *J Wound Care*. 2009;**18**:497-503.
16. Yim E, Kirsner RS, Gailey RS, Mandel DW, Chen SC, Tomic-Canic M. Effect of physical therapy on wound healing and quality of life in patients with venous leg ulcers: a systematic review. *JAMA Dermatol*. 2015;**151**:320-327.
17. Klonizakis M, Tew GA, Gumber A, Crank H, King B, Middleton G, Michaels JA. Supervised exercise training as an adjunct therapy for venous leg ulcers: a randomised controlled feasibility trial. *Br J Dermatol*. 2017;**27**: In Press. doi: 10.1111/bjd.16089
18. Layden J, Michaels J, Bermingham S, Higgins B, Guideline Development Group. Diagnosis and management of lower limb peripheral arterial disease: summary of NICE guidance. *BMJ*. 2012;**345**:e4947. doi: 10.1136/bmj.e4947
19. Liddell F, Webber J. Pulmonary rehabilitation for chronic obstructive pulmonary disease: a pilot study evaluating a once-weekly versus twice-weekly supervised programme. *Physiotherapy*. 2010;**96**:68-74.
20. Sandercock GR, Cardoso F, Almodhy M, Pepera G. Cardiorespiratory fitness changes in patients receiving comprehensive outpatient cardiac rehabilitation in the UK: a multicentre study. *Heart*. 2013;**99**:785-90.

21. Taylor KL, Weston M, Batterham A. Evaluating intervention fidelity: an example from a high-intensity interval training study. *PLoS One*. 2015;**10**:e0125166. doi: 10.1371/journal.pone.0125166
22. Tew GA, Michaels J, Crank H, Middleton G, Gumber A, Klonizakis M. Supervised exercise training as an adjunctive therapy for venous leg ulcers: study protocol for a randomised controlled trial. *Trials*. 2015;**6**:443-453.
23. Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, Nieman DC, Swain DP. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc*. 2011;**43**:1334-1359.
24. Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc*. 1982;**14**:377-381.
25. Purvis JW, Cureton KJ. Ratings of perceived exertion at the anaerobic threshold. *Ergonomics*. 1981;**24**:295-300.
26. Cohen J. *Statistical power analysis for the behavioral sciences*. 2<sup>nd</sup> ed. Hillsdale, NJ: Lawrence Earlbaum Associates; 1988.
27. Picorelli AM, Pereira LS, Pereira DS, Felício D, Sherrington C. Adherence to exercise programs for older people is influenced by program characteristics and personal factors: a systematic review. *J Physiother*. 2014;**60**:151-6.
28. Weston M, Batterham AM, Tew GA, Kothmann E, Kerr K, Nawaz S, Yates D, Danjoux G. Patients Awaiting Surgical Repair for Large Abdominal Aortic Aneurysms Can Exercise at Moderate to Hard Intensities with a Low Risk of Adverse Events. *Front Physiol*. 2017; **9**:684.

29. Weston KS, Wisløff U, Coombes JS. High-intensity interval training in patients with lifestyle-induced cardiometabolic disease: a systematic review and meta-analysis. *Br J Sports Med.* 2014;**48**:1227-1234.
30. Watson E, Yu F. Monitoring exercise delivery to increase participation adherence in older adults with Alzheimer's disease. *J Gerontol Nurs.* 2013;**39**:11-4.
31. West MA, Loughney L, Lythgoe D, Barben CP, Sriadam R, Kemp GJ, Grocott MP, Jack S. Effect of rehabilitation on objectively measured physical fitness after neoadjuvant treatment in preoperative rectal cancer patients: a blinded interventional pilot study. *Br J Anaesth.* 2015;**114**:244-251.
32. Padilla J, Krasnoff J, Da Silva M, Hsu CY, Frassetto L, Johansen KL, Painter P. Physical functioning in patients with chronic kidney disease. *J Nephrol.* 2008;**21**:550-559.
33. Baker MK, Kennedy DJ, Bohle PL, Campbell DS, Knapman L, Grady J, Wiltshire J, McNamara M, Evans WJ, Atlantis E, Fiatarone Singh MA. Efficacy and feasibility of a novel tri-modal robust exercise prescription in a retirement community: a randomized, controlled trial. *J Am Geriatr Soc.* 2007;**55**:1-10.
34. Carter AM, Daley AJ, Kesterton SW, Woodroffe NM, Saxton JM, Sharrack B. Pragmatic exercise intervention in people with mild to moderate multiple sclerosis: a randomised controlled feasibility study. *Contemp Clin Trials.* 2013;**35**:40-47.
35. Rhodes RE, Martin AD, Taunton JE, Rhodes EC, Donnelly M, Elliot J. Factors associated with exercise adherence among older adults. An individual perspective. *Sports Med.* 1999; **28**:397-411.
36. Pothirat C, Chaiwong W, Phetsuk N. Efficacy of a simple and inexpensive exercise training program for advanced chronic obstructive pulmonary disease patients in community hospitals. *J Thorac Dis.* 2015;**7**:637-643.

37. Taaffe DR, Newton RU, Spry N, Joseph D, Chambers SK, Gardiner RA, Wall BA, Cormie P, Bolam KA, Galvão DA. Effects of Different Exercise Modalities on Fatigue in Prostate Cancer Patients Undergoing Androgen Deprivation Therapy: A Year-long Randomised Controlled Trial. *Eur Urol.* 2017;**72**:293-299.
38. Winchester R, Turner LA, Thomas K, Ansley L, Thompson KG, Micklewright D, Gibson AS. Observer effects on the rating of perceived exertion and affect during exercise in recreationally active males. *Percept Mot Skills.* 2012;**115**:213-227.

## Tables

<b>Variable</b>	<b>Exercise group (n=18)</b>
<b>Age, years</b>	66.9 (13.9)
<b>Gender, number male/female</b>	8/10
<b>Stature, cm</b>	171.1 (11.9)
<b>Body mass, kg</b>	102.1 (29.4)
<b>Ulcer size, cm<sup>2</sup>, median (range)</b>	4.9 (1.9 to 136.4)
<b>Duration of ulcer, months, median (range)</b>	5 (1 to 72)
<b>Ankle-brachial index</b>	1.05 (0.14)
<b>Ankle circumference, cm</b>	27.1 (5.5)
<b>Calf circumference, cm</b>	37.3 (7.6)
<b>Comorbidities, n (%)</b>	
<b>Hypertension</b>	7 (39)
<b>History of other CVD</b>	1 (6)
<b>Non-insulin-dependent diabetes</b>	4 (22)
<b>History of cancer</b>	2 (11)
<b>Hypercholesterolemia</b>	1 (6)

Table 1: Exercise group participant characteristics (Values are mean (SD) unless otherwise stated; CVD, cardiovascular disease).

<b>Variable</b>	<b>Estimated Range</b> <i>(Moderate Intensity 60% - 80%)</i>	<b>Base-Line</b> <b>(n= 18)</b>	<b>Mid-point</b> <b>(n= 13)</b>	<b>Intervention End</b> <b>(n=13)</b>	<b>P value;</b> <b><math>\eta^2</math></b>	<b>Post Hoc (Baseline-Midpoint);</b> <b>Cohen's d</b>	<b>Post Hoc (Baseline-Intervention End);</b> <b>Cohen's d</b>	<b>Post Hoc (Midpoint-Intervention End);</b> <b>Cohen's d</b>
<b>Aerobic Training HR</b>	91-121 (10-13)	103 (14)	107 (12)	112 (18)	0.7; 0.01	0.38; 0.32	0.14; 0.35	0.45; 0.30
<b>Aerobic Training RPE</b>	12-14	12(0)	12(0)	12(0)	0.3; 0.05	0.71; 0.65	0.14; 0.26	0.13; 0.37
<b>Resistance Training RPE</b>	12-14	12(1)	12(0)	12(0)	0.3; 0.05	0.5; 0.25	0.14; 0.62	0.92; 0.38

Table 2: Changes in Exercise Intensity Indices between Exercise Sessions ( $p < 0.05$  for Repeated Measures ANOVA and  $p < 0.0167$  for post-hoc analysis).

	<b>Base- Line (n=18)</b>	<b>Mid- point (n=13)</b>	<b>Intervention End (n=13)</b>	<b>P value; <math>\eta^2</math></b>	<b>Post Hoc (Baseline- Midpoint); Cohen's d)</b>	<b>Post Hoc (Baseline- Intervention End); Cohen's d</b>	<b>Post Hoc (Midpoint- Intervention End); Cohen's d</b>
<b>Aerobic (Min)</b>	19 (8)	26 (5)	29 (3)	<0.01; 0.35	<0.01; 1.10	<0.01; 1.82	0.11; 0.68
<b>Squats</b>	5 (12)	14 (18)	36 (18)	<0.01; 0.42	0.08; 0.64	<0.01; 2.10	<0.01; 1.21
<b>Sit to Stand</b>	12 (10)	29 (17)	36 (19)	<0.01; 0.32	<0.01; 1.21	<0.005; 1.68	0.28; 0.43
<b>Step Ups</b>	14 (15)	24 (18)	31 (22)	0.04; 0.14	0.13; 0.56	<0.01; 0.98	0.29; 0.42
<b>Calf Raises</b>	19 (13)	36 (13)	42 (14)	<0.01; 0.35	<0.01; 1.21	<0.01; 1.62	0.28; 0.43

Table 3: Changes in Aerobic and Resistance Exercise Indices between Exercise Sessions ( $p < 0.05$  for Repeated Measures ANOVA and  $p < 0.0167$  for post-hoc analysis).

<b>Element</b>	<b>Fidelity Element</b>	<b>Percentage of Completion According to Protocol</b>
<b>Aerobic</b>	Duration	81%*
	Intensity	100%
	Duration and Intensity	81%
<b>Resistance</b>	Number of Exercises	62%
	Repetitions	78%
	Sets	73%
	Intensity	91%
<b>Flexibility</b>	Duration	93%
	Number of Exercises	93%

Table 4: Assessment of Exercise Fidelity (\* ≥25 minutes of total duration of aerobic exercises).