

Northumbria Research Link

Citation: Schoenmakers, Patrick P.J.M., Hettinga, Florentina and Reed, Kate E. (2019) The Moderating Role of Recovery Durations in High-Intensity Interval-Training Protocols. *International Journal of Sports Physiology and Performance*, 14 (6). pp. 859-867. ISSN 1555-0265

Published by: Human Kinetics

URL: <https://doi.org/10.1123/ijsp.2018-0876> <<https://doi.org/10.1123/ijsp.2018-0876>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/40085/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



The moderating role of recovery durations in high intensity interval training protocols

Journal:	<i>International Journal of Sports Physiology and Performance</i>
Manuscript ID	IJSPP.2018-0876.R1
Manuscript Type:	Invited Commentary
Date Submitted by the Author:	01-Mar-2019
Complete List of Authors:	Schoenmakers, Patrick; University of Essex, School of Sport, Rehabilitation and Exercise Sciences Hettinga, Florentina; Northumbria University Reed, Kate; University of Essex, School of Sport, Rehabilitation and Exercise Sciences
Keywords:	HIIT, protocol optimization, rest intervals, work: rest ratio

SCHOLARONE™
Manuscripts

Title page

Title of the article: The moderating role of recovery durations in high intensity interval training protocols

Submission type: Invited Commentary

Full names of the authors and institutional/corporate affiliations: Patrick P.J.M. Schoenmakers, Florentina J. Hettinga, Kate E. Reed,

Author's affiliation:

School of Sport, Rehabilitation and Exercise Sciences, University of Essex, Wivenhoe Park, United Kingdom, CO4 3SQ.

Contact details for the corresponding author:

Dr Katharine Reed,

Email : reedk@essex.ac.uk,

Phone: +44 (0) 1206 873326

Preferred running Head: Recovery Duration in Interval Training

Abstract word count: 242

Text-only word count: 1916

Number of figures (2) and tables (1)

1 **Abstract:**

2 **Purpose:** Over recent years, multiple studies have tried to optimize the exercise intensity and
3 duration of work intervals in high intensity interval training (HIIT) protocols. Whilst an
4 optimum work interval is of major importance to facilitate training adaptations, an optimum
5 HIIT protocol can only be achieved with an adequate recovery interval separating work
6 bouts. Surprisingly little research has focussed on the acute responses and long term impact
7 of manipulating recovery intervals in HIIT sessions. This invited commentary therefore aims
8 to review and discuss the current literature and increase the understanding of the moderating
9 role of recovery durations in HIIT protocols.

10 **Conclusion:** The acute responses to manipulations in recovery durations in repeated sprint
11 training (RST), sprint interval training (SIT) and aerobic interval training (AIT) protocols
12 have recently begun to receive scientific interest. However, limited studies have manipulated
13 only the recovery duration in RST, SIT or AIT protocols to analyze the role of recovery
14 durations on long term training adaptations. In RST and SIT, longer recovery intervals (≥ 80
15 sec) facilitate higher workloads in subsequent work intervals (compared with short recovery
16 intervals), whilst potentially lowering the aerobic stimulus of the training session. In AIT, the
17 total physiological strain endured per training protocol appears not to be moderated by the
18 recovery intervals, unless the recovery duration is too short. This invited commentary
19 highlights that further empirical evidence on a variety of RST, SIT and AIT protocols and in
20 other exercise modalities than cycling is needed.

21

22 **KEYWORDS:** HIIT, protocol optimization, rest intervals, work: rest ratio

23 Introduction:

24 High intensity interval training (HIIT) is regarded a highly effective training modality to
25 improve cardiorespiratory and metabolic functioning, and is common practice in training
26 regimes of many athletes, particularly those involved in endurance events. ¹ In HIIT, repeated
27 periods of vigorous exercise (work interval) are interspersed with recovery periods, and a
28 complex interplay between the number of intervals, the exercise intensities and the duration
29 of both the work and recovery intervals determine the workload of a HIIT session. ^{2,3} Based
30 on the duration and exercise intensities of work intervals, HIIT can be divided into multiple
31 training forms, for which many terms exist. In this invited commentary we will use and
32 discuss the terms repeated sprint training (RST), sprint interval training (SIT) and aerobic
33 interval training (AIT) as the three main subcategories of HIIT, each targeting different
34 physiological, neuromuscular and mechanical adaptations. ¹ In recent years, many studies
35 have tried to optimize the work intervals of HIIT protocols. A demanding 'work interval' is
36 needed to facilitate training adaptations, with adaptations determined at a cellular level by
37 heat shock proteins, PCG1a and other components ⁴, but a successful HIIT protocol can only
38 be achieved when work bouts are separated by an adequate recovery. Surprisingly little
39 research has explored the overall impact of recovery intervals, and a better understanding of
40 optimum exercise intensities and recovery durations in HIIT protocols is therefore timely.
41 This invited commentary will 1) review the current knowledge of the moderating role of
42 recovery duration on high intensity protocols, and 2) form a basis from which coaches and
43 sports scientists can optimise HIIT protocols according to their specific targets.
44 Characteristics of all reviewed studies are summarized in Supplementary material Table S1

45 Recovery Intervals in HIIT: How are recovery intervals usually determined?

46 A multitude of approaches are available for the prescription of recovery intervals in HIIT.
47 The most common approach is the use of a fixed work:recovery ratio (i.e., W:R = 2:1, 1:1,
48 1:8). A fixed W:R ratio separates work intervals by an *a priori* set recovery duration, for
49 instance, when W:R = 1:2, the recovery duration is twice the duration of the work interval. In
50 an attempt to individualize recovery intervals, the return of heart rate (HR) to a set threshold
51 value or to a percentage of maximum heart rate (HRmax) is used. However, the present
52 understandings of the determinants of HR recovery suggest that this practice is not
53 appropriate in the prescription of recovery durations. This was for instance evidenced by
54 Edwards et al., ⁵ who reported decreases up to ~10-15 sec for each 1000m running effort in a
55 5*1000m sequence when recovery intervals were based on HR return, compared to a W:R =
56 1:1 protocol, of which the latter resulted in ~80 sec extra recovery time between repetitions.
57 Lastly, a number of studies have used self-selected (SS) recovery durations in HIIT protocols,
58 in which athletes started subsequent work intervals when they felt 'adequately recovered to
59 exercise at the required intensity'. ⁵⁻¹⁰ While a considerable amount of variation was evident
60 in SS recovery durations across different HIIT protocols, and SS recovery time is potentially
61 dependent on maturation status ^{7,10} (see figure 1), the current understanding is that athletes
62 can adequately select recovery durations to achieve the required exercise intensities in
63 subsequent work intervals in both RST and SIT (see figure 1) and AIT (see figure 2).
64 Athletes new to the use of SS recovery intervals will likely choose a 'shorter than optimal'
65 recovery time, as common HIIT protocols typically incorporate 'short' recovery durations
66 (e.g. 1000m work : 200m recovery), which potentially compromises training effects.

67 >> figure 1 and figure 2 around here <<

68 Physiological basis of recovery.

69 The main metabolic processes that take place during recovery from intense exercise bouts are
70 the repletion of phosphocreatine stores (PCr), the removal of hydrogen ions (H⁺) and
71 restitution of the acid-base balance of the exercising muscles.^{1,11,12} These processes proceed
72 at different rates, with PCr having a much faster half-life (~30 sec) and **achieving** complete
73 restoration (~3 min),¹¹ compared with blood lactate [BLa] and pH recovery (6 - 10 min).¹² In
74 order to work at the required exercise intensity during subsequent intervals, recovery
75 intervals need to be long enough to accommodate the return to metabolic homeostasis. An
76 imbalance between the demands of the work intervals and the recovery potential of the
77 recovery intervals can lead to premature fatigue, which potentially reduces the number of
78 planned intervals, or lowers the work intensity during subsequent intervals. **An example of an**
79 **inadequate W:R is seen in the** study by Laursen et al.,¹³ who reported that two groups of well
80 trained cyclists completed only 64% of the total prescribed number of work bouts over a 4
81 week training cycle. **Participants were ‘pushed to exhaustion’ in each session, as inadequate**
82 **recovery had been prescribed given the intensity of the work interval, resulting in failure to**
83 **complete the session.** While the training intervention still improved time trial performance,
84 peak power output (PO_{peak}) and the maximum oxygen uptake (VO_{2max}),¹³ a protocol
85 involving a longer recovery interval may have evoked even greater improvements.

86 **The recovery duration during RST & SIT**

87 Repeated all-out (or sometimes labelled ‘supramaximal’⁴) sprint training has received a
88 growing research interest, as it replicates **the demands** of maximal-intensity **sprint efforts**
89 **typically performed in field-based team sports and endurance sports.** In practical terms, based
90 on the duration of the sprints **and the subsequent recovery duration**, sprint training can be
91 divided into either short (3–10 sec; RST) or long (15–30 sec; SIT) sprints.

92 In RST, a positive effect on performance in subsequent 4 – 8 sec supramaximal sprints in
93 cycling **power**^{14–17} and running **speed**^{18,19} has been reported when longer recovery durations
94 were employed. Longer recovery intervals resulted in a lower average HR and oxygen uptake
95 (VO₂) over the training session.^{14,15,17,20} Further, the fatigue index (percentage decline
96 between PO_{peak} first and last sprint), [BLa] and ratings of perceived exertion (RPE) were
97 lower when sprints were interspersed with longer recovery intervals,^{15,19} which was
98 accompanied by a greater muscular re-oxygenation.²⁰

99 **In SIT protocols similar** beneficial performance outcomes were reported across a multitude of
100 exercise modalities when recovery duration was increased between work intervals.^{8,21–23}
101 McEwan et al.,⁸ compared the acute physiological responses and running performance in
102 12 × 30 sec sprints, wherein the recovery duration was either fixed (30 sec) or self-selected
103 (SS). SS recovery time increased over the protocol (see figure 1) and averaged 51±15 sec.
104 The longer recovery intervals in SS resulted in a reduced time ≥ 90% HR_{max}, but facilitated
105 the attainment of significantly higher running speeds. In agreement with these findings,
106 Gosselin et al.,²⁴ reported a decrease in mean and peak VO₂ and mean HR in a SIT protocol
107 alternating 60 sec work intervals with 60 sec recovery, compared with 30 sec recovery
108 intervals. Less than 30 sec recovery between ‘all out’ sprints seems to have a detrimental
109 effect on power production in subsequent cycling sprints, whereas the aerobic demand in
110 sprints separated by 120 sec recovery are too low to induce endurance adaptations.^{22–24}
111 Kavaliauskas et al.,²³ therefore suggested 80 sec recovery intervals between sprints are
112 optimal when targeting both power and endurance adaptations.

113 **The recovery duration during ‘aerobic’ interval training**

114 HIIT incorporating long work intervals (up to 16 min) is typically described as ‘aerobic
115 interval training’ (AIT), as work intensities are undeniably high - but ultimately submaximal.
116 It was suggested by Thevenet et al.,²⁵ that the time athletes spend in their ‘red zone’ per AIT
117 could serve as a good criterion to judge the effectiveness of a protocol. The ‘red zone’ refers
118 to the intensity domain close to $\dot{V}O_2\text{max}$ ($\geq 90\% \dot{V}O_2\text{max}$) in which the oxygen delivery and
119 utilization systems are maximally stressed.¹ Previous research showed that trained runners
120 reach a steady state of around 90 - 95% $\dot{V}O_2\text{max}$ / HRmax across repeated 4 min work
121 intervals, independent of an increased recovery duration between bouts.^{6,9,26,27} Both Smilios
122 et al.,²⁷ and Schoenmakers⁹ reported changes in the O_2 and HR kinetics when recovery
123 durations increased (more so, mean response time was faster when intervals started from a
124 lower metabolic rate), resulting in similar time spent $\geq 90\%$ and 95% $\dot{V}O_2\text{max}$ and HRmax
125 between the different recovery durations, suggesting a comparable physiological load of the
126 AIT protocol.^{9,27} Increasing the recovery duration from 1 to 4 min did not significantly affect
127 [BLa] responses following each 4 min work intervals in runners, suggesting a balance
128 between lactate production and lactate buffering capacity.^{6,26} In a study where participants
129 were working at a greater intensity, a greater [Bla] was evident when 6 x 2 min cycling
130 intervals were separated by either 1 (AIT1) or 3 min (AIT3) passive recovery intervals.²⁸ The
131 shorter recovery intervals in AIT1 induced a lower post exercise PCr content compared with
132 AIT3, however, these larger perturbations in muscle metabolites did not result in greater
133 training adaptations in AIT1 compared with AIT3.²⁸

134 Using self-paced AIT protocols, in which work intensities were not predefined but rather
135 determined by the integrative outcome of feedback from external and internal receptors,
136 multiple research groups^{5,6,9,26,29} have evaluated running performance across work intervals.
137 In highly trained runners, increasing the recovery duration in a 10*400m set speed sequence
138 (60 vs. 120 vs. 180 sec) resulted in a lower RPE.²⁹ Trained male⁶, and recreational active
139 male and female runners²⁶ were able to increase their mean running speed in 6 x 4 min
140 intervals when the recovery duration was increased from 1 min to 2 min. A further increase in
141 recovery duration (4 min) did not provide extra performance benefits for the trained runners.
142 Conversely, Laurent et al.,²⁶ reported an additional increase in running speed when extra
143 recovery time was available. Schoenmakers et al.,⁹ reported the highest mean running speed
144 when 6 x 4 min intervals (ran on a curved non-motorized treadmill) were separated by 3 min,
145 compared to 1 min, 2 min or a SS recovery interval. These results overall indicate that
146 adequate recovery will result in the attainment of the desired work intensity within the limits
147 and requirements of a specific protocol, however, the ‘optimum’ recovery duration, most
148 likely is highly individual and depending on training status.

149 Practical Applications

150 In RST and SIT protocols, longer recovery intervals (≥ 80 sec) facilitate higher work
151 intensities in subsequent sprints and lower the fatigue index, whereas a shorter recovery
152 duration in these protocols increases the overall physiological demands of a training session.
153^{22,23} Long recovery intervals in AIT protocols allow athletes to attain higher workloads (speed
154 or power) in successive work bouts when exercise intensities are not fixed, without
155 compromising the overall physiological stimulus of a training session.^{6,9,26} When work
156 intensities are fixed in AIT protocols, the same training sessions is typically completed with a
157 lower RPE when longer recovery intervals are available, again, without compromising in the
158 physiological stimulus.²⁷⁻²⁹ Ultimately, depending on the exercise intensities of work
159 intervals, a recovery interval of 3 min is expected to be sufficient to avoid premature fatigue
160 in AIT protocols.

161 Conclusion

162 The acute responses to manipulations in recovery durations in RST, SIT and AIT protocols
163 **are receiving increasing** scientific interest. The manipulation of recovery durations in RST
164 and SIT protocols results in different acute physiological and perceptual responses, and most
165 likely in different training adaptations. The current understanding is that training at higher
166 workloads in RST and SIT protocols elicit greater adaptations in PO_{peak} and VO_{2max},
167 however, this has only been evidenced in cycling protocols. In AIT, the physiological strain
168 endured per training protocol appears not to be moderated by the recovery intervals, unless
169 the recovery interval is too short and causes premature fatigue. When adequate recovery
170 intervals are available in AIT protocols, a further increase in recovery duration is not
171 expected to provide greater physiological and/or performance adaptations when exercise
172 intensities are fixed. However, when work intensities are not predefined, longer recovery
173 durations may facilitate a higher external training load, and may therefore allow for greater
174 training adaptations. Further empirical evidence on a variety of RST, SIT and AIT protocols
175 in exercise modalities other than cycling are needed to fully determine the moderating effects
176 of recovery duration in HIIT sessions.

177

178

179

180

181

182

183

184

185

186

187 **Acknowledgements**

188 None

189

For Peer Review

190 **References**

- 191 1. Buchheit M, Laursen PB. High-intensity interval training, solutions to the
192 programming puzzle: Part I: cardiopulmonary emphasis. *Sports Med.* 2013;43(5):313-
193 338. doi:10.1007/s40279-013-0029-x
- 194 2. Åstrand I, Åstrand P-O, Christensen EH, Hedman R. Intermittent Muscular Work.
195 *Acta Physiol Scand.* 1960;48(3-4):448-453. doi:10.1111/j.1748-1716.1960.tb01879.x
- 196 3. Tschakert G, Hofmann P. High-intensity intermittent exercise: Methodological and
197 physiological aspects. *Int J Sports Physiol Perform.* 2013;8(6):600-610.
- 198 4. Gibala MJ, Little JP, MacDonald MJ, Hawley JA. Physiological adaptations to low-
199 volume, high-intensity interval training in health and disease. *J Physiol.*
200 2012;590(5):1077-1084. doi:10.1113/jphysiol.2011.224725
- 201 5. Edwards AM, Bentley MB, Mann ME, Seaholme TS. Self-pacing in interval training:
202 A teleoanticipatory approach. *Psychophysiology.* 2011;48(1):136-141.
203 doi:10.1111/j.1469-8986.2010.01034.x
- 204 6. Seiler S, Hetlelid KJ. The impact of rest duration on work intensity and RPE during
205 interval training. *Med Sci Sports Exerc.* 2005;37(9):1601-1607.
- 206 7. Gibson N, Brownstein C, Ball D, Twist C. Physiological, Perceptual and Performance
207 Responses Associated With Self-Selected Versus Standardized Recovery Periods
208 During a Repeated Sprint Protocol in Elite Youth Football Players: A Preliminary
209 Study. *Pediatr Exerc Sci.* 2017;29(2):186-193. doi:10.1123/pes.2016-0130
- 210 8. McEwan G, Arthur R, Phillips SM, Gibson N V, Easton C. Interval running with self-
211 selected recovery: Physiology, performance, and perception. *Eur J Sport Sci.* May
212 2018:1-10. doi:10.1080/17461391.2018.1472811
- 213 9. Schoenmakers PPJ, Reed KE. The Effects of Recovery Duration on Physiological and
214 Perceptual Responses of Trained Runners during four Self-Paced HIIT Sessions. *J Sci*
215 *Med Sport.* September 2018. doi:10.1016/J.JSAMS.2018.09.230
- 216 10. Brownstein CG, Ball D, Micklewright D, Gibson N V. The Effect of Maturation on
217 Performance During Repeated Sprints With Self-Selected Versus Standardized
218 Recovery Intervals in Youth Footballers. *Pediatr Exerc Sci.* July 2018:1-6.
219 doi:10.1123/pes.2017-0240
- 220 11. McMahon S, Jenkins D. Factors affecting the rate of phosphocreatine resynthesis
221 following intense exercise. *Sports Med.* 2002;32(12):761-784.
- 222 12. Menzies P, Menzies C, McIntyre L, Paterson P, Wilson J, Kemi OJ. Blood lactate
223 clearance during active recovery after an intense running bout depends on the intensity
224 of the active recovery. *J Sports Sci.* 2010;28(9):975-982.
225 doi:10.1080/02640414.2010.481721
- 226 13. Laursen PB, Shing CM, Peake JM, Coombes JS, Jenkins DG. Interval training
227 program optimization in highly trained endurance cyclists. *Med Sci Sports Exerc.*
228 2002;34(11):1801-1807. doi:10.1249/01.MSS.0000036691.95035.7D
- 229 14. Glaister M, Stone MH, Stewart AM, Hughes M, Moir GL. The influence of recovery

- 230 duration on multiple sprint cycling performance. *J Strength Cond Res.* 2005;19(4):831-
231 837.
- 232 15. Baker JS, Van Praagh E, Gelsei M, Thomas M, Davies B. High-intensity intermittent
233 cycle ergometer exercise: effect of recovery duration and resistive force selection on
234 performance. *Res Sports Med.* 2007;15(2):77-92. doi:10.1080/15438620601184190
- 235 16. Lee C-L, Cheng C-F, Lin J-C, Huang H-W. Caffeine's effect on intermittent sprint
236 cycling performance with different rest intervals. *Eur J Appl Physiol.*
237 2012;112(6):2107-2116. doi:10.1007/s00421-011-2181-z
- 238 17. Shi Q, Tong TK, Sun S, et al. Influence of recovery duration during 6-s sprint interval
239 exercise on time spent at high rates of oxygen uptake. *J Exerc Sci Fit.* 2018;16(1):16-
240 20. doi:10.1016/j.jesf.2018.01.001
- 241 18. Iaia FM, Fiorenza M, Perri E, Alberti G, Millet GP, Bangsbo J. The Effect of Two
242 Speed Endurance Training Regimes on Performance of Soccer Players. Sandbakk O,
243 ed. *PLoS One.* 2015;10(9):e0138096. doi:10.1371/journal.pone.0138096
- 244 19. Padulo J, Tabben M, Ardigo LP, et al. Repeated sprint ability related to recovery time
245 in young soccer players. *Res Sport Med.* 2015;23(4):412-423.
246 doi:10.1080/15438627.2015.1076419
- 247 20. Ohya T, Aramaki Y, Kitagawa K. Effect of Duration of Active or Passive Recovery on
248 Performance and Muscle Oxygenation during Intermittent Sprint Cycling Exercise. *Int*
249 *J Sports Med.* 2013;34(07):616-622. doi:10.1055/s-0032-1331717
- 250 21. Toubekis AG, Douda HT, Tokmakidis SP. Influence of different rest intervals during
251 active or passive recovery on repeated sprint swimming performance. *Eur J Appl*
252 *Physiol.* 2005;93(5-6):694-700. doi:10.1007/s00421-004-1244-9
- 253 22. Hazell TJ, MacPherson REK, Gravelle BMR, Lemon PWR. 10 or 30-S Sprint Interval
254 Training Bouts Enhance Both Aerobic and Anaerobic Performance. *Eur J Appl*
255 *Physiol.* 2010;110(1):153-160. doi:10.1007/s00421-010-1474-y
- 256 23. Kavaliuskas M, Aspe RR, Babraj J. High-Intensity Cycling Training: The Effect of
257 Work-to-Rest Intervals on Running Performance Measures. *J Strength Cond Res.*
258 2015;29(8):2229-2236. doi:10.1519/JSC.0000000000000868
- 259 24. Gosselin LE, Kozlowski KF, DeVinney-Boymel L, Hambridge C. Metabolic Response
260 of Different High-Intensity Aerobic Interval Exercise Protocols. *J Strength Cond Res.*
261 2012;26(10):2866-2871. doi:10.1519/JSC.0b013e318241e13d
- 262 25. Thevenet D, Tardieu M, Zouhal H, Jacob C, Abderrahman BA, Prioux J. Influence of
263 exercise intensity on time spent at high percentage of maximal oxygen uptake during
264 an intermittent session in young endurance-trained athletes. *Eur J Appl Physiol.*
265 2007;102(1):19-26. doi:10.1007/s00421-007-0540-6
- 266 26. Laurent CM, Vervaecke LS, Kutz MR, Green JM. Sex-specific responses to self-
267 paced, high-intensity interval training with variable recovery periods. *J Strength Cond*
268 *Res.* 2014;28(4):920-927. doi:10.1519/JSC.0b013e3182a1f574
- 269 27. Smilios I, Myrkos A, Zafeiridis A, Toubekis A, Spassis A, Tokmakidis SP. The effects
270 of recovery duration during high-intensity interval exercise on time spent at high rates

- 271 of oxygen consumption, oxygen kinetics and blood lactate. *J Strength Cond Res.*
272 March 2017;1. doi:10.1519/JSC.0000000000001904
- 273 28. Edge J, Eynon N, McKenna MJ, Goodman CA, Harris RC, Bishop DJ. Altering the
274 rest interval during high-intensity interval training does not affect muscle or
275 performance adaptations. *Exp Physiol.* 2013;98(2):481-490.
276 doi:10.1113/expphysiol.2012.067603
- 277 29. Zavorsky GS, Montgomery DL, Pearsall DJ. Effect of intense interval workouts on
278 running economy using three recovery durations. *Eur J Appl Physiol Occup Physiol.*
279 1998;77(3):224-230. doi:10.1007/s004210050326
- 280

For Peer Review

281 **Figure captions:**

282 Figure 1: Mean \pm SD self-selected recovery duration between 12 x 30 sec ⁸ , or 12 x 30m ^{7,10}
283 intervals

284 Figure 2: Mean \pm SD self-selected recovery duration between 6 x 4 min ^{6,9} , or 5 x 1000m ⁵
285 intervals

286

287

288

289

For Peer Review

290 **Table caption**

291 Table 1: Summary of participant and training characteristics of reviewed studies

For Peer Review




Figure 1: Mean±SD self-selected recovery duration between 12 x 30 sec 8 , or 12 x 30m 7,10 intervals
419x199mm (150 x 150 DPI)




Figure 2: Mean±SD self-selected recovery duration between 6 x 4 min 6,9 , or 5 x 1000m 5 intervals

656x313mm (96 x 96 DPI)

Table 1: Summary of participant and training characteristics of reviewed studies

Study	Sample Size, Age	Modality	High Intensity Interval Training Protocol	Recovery Duration	Key Findings
Repeated Sprint Training					
Baker et al. (2007) ¹⁵	n = 8, 26.6 ± 7.8	Cycling	Participants performed 8 × 6 sec sprint on a cycling ergometer against 0.75 g.kg ⁻¹ FFM or TBM	30 sec, 1MIN	AR: Peak power output was higher in both the FFM and TBM conditions in 1MIN vs 30 sec, accompanied by a significantly lower fatigue index. HR was higher in both 30 sec protocols, with no differences in RPE and end [Bla] measures evident.
Brownstein et al. (2018) ¹⁰	pre-PHV, n = 14, 12 ± 0.4	Running	Participants performed a repeated sprint sequence twice, comprising 10 × 30 m efforts (~5 sec)	30 sec, SS	AR: Recovery duration in SS significantly shorter (~12 sec). Mean sprint time faster in 30 sec, accompanied by smaller performance decrement. Mean and peakHR higher in SS.
	Post-PHV, n = 14, 14 ± 0.5				AR: Recovery duration in SS significantly shorter (~8 sec). Mean sprint time faster in 30 sec, accompanied by smaller performance decrement. Mean and peakHR higher in SS.
Gibson et al. (2017) ⁷	n = 11, 14 ± 1	Running	Participants performed two repeated sprint assessment of 10 × 30 m sprint efforts (~5 sec)	30 sec, SS	AR: Training sequence shorter in SS, as SS recovery duration is significantly shorter (~10 sec). Mean sprint time significantly faster in 30 sec. No differences in peakHR, [Bla] and RPE.
Glaister et al. (2005) ¹⁴	n = 25, 20.6 ± 1.5	Cycling	Participants completed 20 × 5 sec maximal sprints on a friction-braked cycle ergometer	10 sec, 30 sec	AR: Peak (~4%) and mean (~26%) power output higher in 30 sec, with lower measures of fatigue, RPE and end [Bla]. Contrary, VO ₂ , RER and HR measures were higher in 10 sec in both the work and recovery intervals.
Lee et al. (2011) ¹⁶	n = 14, 18.7 ± 0.8	Cycling	Participants completed two intermittent sprint cycling tests (ISCTs), which were composed of 12 × 4 sec sprints. Tests were separated by 4 min active recovery	20 sec, 90 sec	AR: Peak and mean sprint power in both ISCTs higher in 90 sec vs 20 sec, with a lower fatigue index and RPE score. End [Bla] higher in 20 sec.
Ohya et al. (2013) ²⁰	n = 8, 25.5 ± 2.6	Cycling	Participants performed 10 maximal 5 sec sprints interspersed with either active recovery (ACT, cycling at 40% VO ₂ max) or passive recovery (PAS, sitting)	25 sec, 50 sec, 100 sec	AR: Mean and peak power decrement over sprints was lowest in 100 sec and, independent of ACT/PAS, inversely related to recovery time. Mean VO ₂ and [Bla] were higher in 25 sec > 50 sec > 100 sec, whilst muscular reoxygenation was lower in 25 sec.
Padulo et al. (2015) ¹⁹	n = 17, 16 ± 0	Running	Participants completed three testing sessions, in which they performed six maximal 40 m shuttle sprints (20+20 m with a 180° change of direction, ~6 sec)	15 sec, 20 sec, 25sec	AR: Total sprint time was ~3% faster in 25 sec compared to 15 sec, and ~1.3% compared to 20 sec. [Bla] and fatigue index were highest in 15 sec, followed by 20 sec, and lowest in 25 sec.
Shi et al. (2018) ¹⁷	n = 13, 26.2 ± 6.2	Cycling	Participants finished three RST protocols, consisting of 40 x 6 sec all-out sprints on a cycling ergometer (with resistance equating 7.5% body mass)	15 sec, 30 sec, 1MIN	AR: Peak and mean power output was higher in 1MIN compared to 15 sec and 30 sec, with a notable lower RPE. Accumulated time ≥ 80% and 90% VO ₂ max increased as recovery time decreased, however, for HR this was only evident in time ≥ 95% HRmax.
Sprint Interval Training					
Gosselin et al. (2012) ²⁴	n = 8, 23.1 ± 2.1	Running	Participants performed 2 different training protocols, in which they exercised at a workload corresponding to 90% VO ₂ max for 60 sec	30 sec, 1MIN	AR: Mean and peak VO ₂ and HR significantly higher in 30 sec compared to 1MIN, with no differences in RPE. Both protocols failed to achieve 90% VO ₂ max.

Table 2: Continued

Study	Sample Size, Age	Modality	High Intensity Interval Training Protocol	Recovery Duration	Key Findings
Hazell et al. (2010) ²²	n = 48, 24 ± 3.2	Cycling	Participants completed 2 weeks of SIT (3 sessions a week), in which they performed 4-6 'all out' sprint of either 30 sec (G1) or 10 sec (G2 & G3), against 100 g.kg ⁻¹ . CON did not receive SIT	G1: 4MIN, G2: 4MIN, G3: 2MIN	AR: Peak and mean power output in sprints higher in G2 & G3, whilst G1 performed more total work. TA: Improvements in 5 km TT were similar between groups, whereas the increase in VO ₂ max and mean and peak Wingate power output were higher in G1 & G2 compared to G3 and CON.
Iaia et al. (2015) ¹⁸	n = 13, 18.5 ± 1	Running	Participants completed nine SIT sessions, which focussed on speed endurance production (SEP; n = 6) or speed endurance maintenance (SEM; n = 7). Both SEP and SEM consisted of 6-8 reps of 20 sec all-out sprints	SEP: 2MIN, SEM: 40 sec	AR: Mean running speed were higher in SEP sprints compared to SEM, with a lower decrement in speed across subsequent sprints. TA: SEM improved their 200-m sprint time, distance covered in Yo-Yo test increased 10.1% after SEP and 3.8% after SEM.
Kavaliuskas et al. (2015) ²³	G1, n = 8, 41 ± 12 G2, n = 8, 38 ± 7 G3, n = 8, 42 ± 6	Cycling	Participants completed a total of six SIT sessions over a two week period. The SIT protocol consisted of six 10-second "all-out" cycling efforts against a resistance equalling 7.5% of body weight. CON received no SIT	G1: 30 sec, G2: 80 sec, G3: 2MIN	AR: Average HR was greater in G1 compared with G3 for all training sessions, and was greater in G2 compared with G3 for training sessions 1 and 2. TA: All three training groups increased 3km TT to a similar extent. VO ₂ max increased in G1 & G2, but not in G3. Mean and peak Wingate power output increased after G2, whereas G3 only increased their mean power output.
McEwan et al. (2018) ⁸	N = 14, 30 ± 7	Running	Participants performed 12 × 30 sec running intervals at a target intensity of 105% MAS.	30 sec, SS	AR: Mean recovery duration longer in SS (~21 sec). Relative time ≥ 105% MAS and mean running speed greater in SS, whereas time ≥ 90% HRmax was higher in 30 sec compared to SS. No differences in end [Bla] or RPE.
Toubekis et al. (2005) ²¹	N = 16, 21.2 ± 0.6	Swimming	Participants completed eight repetitions of 25-m sprints (~15 sec), followed by a 50-m sprint test 6 min later. Recovery was either ACT or PAS.	45 sec, 2MIN	AR: Mean swimming velocity faster in PAS for both recovery durations, and faster in 2MIN compared to 45 sec with no differences in end [Bla]. 50-m sprint times were 2.4% faster in both ACT and PAS 2MIN conditions vs 45 sec.
Aerobic Interval Training					
Edge et al. (2013) ²⁸	N = 5, 21 ± 2	Cycling	Participants completed 6 × 120 sec intervals, on a power output corresponding to 92% VO ₂ max	1MIN, 3MIN	AR: Average HR in intervals higher in 1MIN vs 3MIN. 1MIN induced a greater end [Bla], H ⁺ and MLa content than 3MIN, while muscle PCr content was less after 1MIN.
Edge et al. (2013) ²⁸	G1, n = 6, 19 ± 1 G2, n = 6	Cycling	Participants performed a total 15 HIIT sessions over a 5 week period, consisting of 6 – 10 × 120 sec intervals at a workload of 92%-111% power output at VO ₂ max	G1: 1MIN, G2: 3MIN	TA: Significant increase in VO ₂ max, PPO and power output at lactate threshold, to a similar extent in both G1 and G2. Improvements in repeated sprint performance were similar.
Edwards et al. (2011) ⁵	N = 11, 26 ± 7	Running	Participants completed a series of four (5 × 1000 m) track running sessions, each at the standardized perceived exertion of RPE 17.	SS_PR1, SS_PR2, HR130, W:R = 1	AR: Recovery significantly shorter in HR130, accompanied by a significant lower mean running velocity and greater fatigue index. Similar HR and end [Bla] between all experimental conditions.

Table 3: Continued

Study	Sample Size, Age	Modality	High Intensity Interval Training Protocol	Recovery Duration	Key Findings
Laurent et al. (2014) ²⁶	G1, n = 8, 20.8 ± 2.1 G2, n = 8, 21.9 ± 3.6	Running	Trained male (G1) and female (G2) runners completed three isoeffort (maximum sustainable intensity) training sessions, each comprising six 4 min interval	1MIN, 2MIN, 4MIN	AR: SS running velocity increased in both groups when longer recovery was available. Independent of recovery duration, mean VO ₂ , HR, [Bla] and RPE were similar across conditions in both G1 & G2. Relative exercise HR and VO ₂ was higher in G2.
Laursen et al. (2002) ¹³	G1, n = 8, 26 ± 6 G2, n = 9, 24 ± 7	Cycling	Participants performed eight AIT sessions over a 4 week period, comprising 8 intervals at Pmax for the duration of 60% Tlim	G1: W:R = 0.5, G2: 65HRmax	AR: G1 had a significantly greater total mean recovery time (~110 sec) between bouts compared with G2. Both groups completed ~64% of prescribed interval bouts. TA: Improvements in VO ₂ max, PPO, and 40 km TT were similar between groups.
Seiler & Hetlelid (2005) ⁶	n = 9, 30 ± 4	Running	Participants performed three isoeffort (maximum sustainable intensity) training sessions, each comprising six 4 min intervals at a constant 5% treadmill incline.	1MIN, 2MIN, 4MIN, SS	AR: Higher running velocity in 2MIN (85% vVO ₂ max) and 4MIN (84% vVO ₂ max) vs 1MIN (83% vVO ₂ max). Higher mean VO ₂ in 2MIN and 4MIN vs 1MIN. No differences in end [Bla], HR, or RPE.
Schoenmakers & Reed (2018) ⁹	N = 12, 34 ± 11	Running	Participants performed four isoeffort (maximum sustainable intensity) training sessions, each comprising six 4 min interval on a non-motorized treadmill	1MIN, 2MIN, 3MIN, SS_PR1	AR: Running velocity significantly higher in 3MIN compared to all other protocols, and higher in ssMIN vs 2MIN. No significant differences in RPE responses, time ≥ 90% and 95% VO ₂ max, or ≥ 90% and 95% HRmax
Smilioset al. (2018) ²⁷	N = 11, 22.1 ± 1	Running	Participants executed, on three separate sessions, 4×4 min runs at 90% of MAS	2MIN, 3MIN, 4MIN	AR: Time ≥ 80 and 90% HRmax was higher in 2MIN and 3MIN compared to 4MIN, but did not differ for VO ₂ measures. Peak HR and VO ₂ were similar between conditions. RPE were higher in 2MIN and 3MIN vs 4MIN, as was 2MIN end [Bla]
Zavorsky et al. (1998) ²⁹	N = 12, 24.8 ± 5.1	Running	Participants performed three interval running workouts of 10 x 400 m on a predefined running speed	1MIN, 2MIN, 3MIN	AR: Mean HR significantly higher in 1MIN, but no differences in peakHR between conditions. RPE increased with decrease in recovery time.

Age is presented mean ± standard deviation

Note: 1MIN; 1 min recovery; 2MIN; 2 min recovery; 3MIN; 3 min recovery; 4MIN; 4 min recovery; ACT: active recovery; AIT: aerobic interval training; AR: Acute responses; [Bla]: blood lactate concentration; CON: control group; FFM: fat-free body mass; H⁺: Hydrogen; HR: heart rate; HR130: recovery duration based on HR return to 130 bpm; HRmax: maximum heart rate; ISCTs: intermittent sprint cycling tests; MAS: maximal aerobic speed; MLa: muscle lactate; PAS: passive recovery; PCr: phosphocreatine; peakHR: peak heart rate; Pmax: minimal power output to elicit VO₂max; post-PHV: post peak height velocity; PPO: peak power output; pre-PHV: pre peak height velocity; RER: respiratory exchange ratio; RPE: ratings of perceived exertion; RST: repeated sprint training SIT: sprint interval training; SS: self-selected recovery duration; SS_PR1 & SS_PR2: self-selected recovery duration based on perceived readiness scale; SEM: speed endurance maintenance; SEP: speed endurance production; TA: Adaptations to a period of training; TBM: total body mass; Tlim: time to exhaustion at Pmax; TT: time trial; VO₂: oxygen consumption; VO₂max: maximum oxygen consumption vVO₂max: minimum running velocity to elicit VO₂max; W:R = 1: recovery duration equal to work interval duration