

Northumbria Research Link

Citation: Alderdice, Fiona and Newham, James (2016) Global maternal mental health: where you live matters [Editorial]. *Journal of Reproductive and Infant Psychology*, 34 (1). pp. 1-2. ISSN 0264-6838

Published by: Taylor & Francis

URL: <https://doi.org/10.1080/02646838.2016.1126945>
<<https://doi.org/10.1080/02646838.2016.1126945>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/41724/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



**Northumbria
University**
NEWCASTLE



UniversityLibrary

Global maternal mental health: where you live matters

It is reassuring to see, in recent years, an increase in the profile of maternal mental health in the perinatal period. We know that maternal mental health has significant implications not only for the woman's ongoing physical and mental health but also that of her family both in the short and long term. But is this increased attention being matched in policy and practice internationally? Identifying women who have mental health problems and ensuring they have timely delivery of care is a universal problem. However where you live matters. Research suggests that this is not just about what services are available but what mental health services women will engage with. A better understanding of cultural perspectives is important to fully understand variability in care and levels of stigma related to mental health in the perinatal period.

In 2008, WHO reported that maternal mental health was studied in 90% of high-income countries (HIC) but data were only available in 10% of middle and lower income countries (LMIC)(WHO, 2008). This lack of data is concerning as the evidence that is available suggests that common mental health problems such as perinatal anxiety and depression are more prevalent in women from LMIC compared to HIC. For example, Atif, Lovell & Rahman (2015) report the weighted mean prevalence of depression and anxiety during the perinatal period in low-income countries was found to be 15.6% during the antenatal period and 19.8% during the postnatal period, which is higher than the rates reported in HICs (6.5% to 12.9%).

There is also high variability *within* LMIC on the prevalence of depression. Fisher et al. (2012) conducted a systematic review that showed that the prevalence of postnatal depression was high in India (23%), Nigeria (48%), Indonesia (22.4%), Bangladesh (33%), Pakistan (25%), and Ethiopia (59.5%); but relatively low in Brazil (15.9%), China (15.5%), Mongolia (9.1%), and Uganda (6.1%). While there was variability on how women were recruited to these studies (which may impact on the reliability of the findings), the disparity in these figures may reflect important cultural variations in the expression of mental illness that needs careful consideration. Such cultural variations have implications for screening and highlight the value of validation studies of the instruments used to screen for mental health problems in pregnancy to ensure their reliability and validity in different populations and ethnic communities.

Even if screening were successful, a large treatment gap has also been reported for mental illness in LMICs, accounting for 76–85% patients with mental health problems not receiving intervention (Atif, Lovell & Rahman, 2015).. Despite recommendations from WHO and the availability of low cost, community based interventions (Clarke, King & Proust, 2013), the maternal mental health agenda has not been incorporated into the primary health care system in most LMIC. The burden of communicable and non-communicable diseases along with a lack of resources has led to the marginalization of mental health within health care in LMIC settings despite its recognized impact on health (Atif, Lovell & Rahman, 2015).

Gaps in mental health screening and care are not just a problem in LMIC. In HIC

settings, women from more deprived areas experience higher levels of perinatal mental health problems and access to care is a significant problem. For example, in this issue, Rosaura, Orengo-Aguayo & Segre (2016) report on a study of the experiences of low income US women taking part in a randomized trial of depression treatment delivered at the point-of-care. In the USA 19% report clinically significant depressive symptoms and these are even more prevalent among low-income and ethnic-minority women. However low-income and ethnic-minority mothers are faced with barriers that make them less likely to receive professional mental health care. Rosaura, Orengo-Aguayo & Segre (2016) spoke with women to explore ways of removing these barriers, as part of a recent US trial of an evidence-based treatment delivered by home visitors/office nursing staff. Such studies are important if we are to implement appropriate and effective services for those most in need.

Similarly, the perinatal mental health needs of mothers in rural areas in high income countries are also poorly understood. There is research on the organization of care; for example, examining the impact that midwifery-led care in rural communities can have upon birth outcomes. However research looking at the wider health of women in pregnancy including mental health and its impact on child health and development is more limited. But there are some concerning reports that highlight that more research is needed. For example, when focusing on women of reproductive age, rates of antidepressant use were higher in women living in remote rural areas (Wemakor et al. 2014).

It is probably not surprising that the gap in both knowledge and services is highly variable depending on where you live and what resources you have available to you but that does not make it acceptable. In the absence of specialist services, other low cost, community based, psychosocial interventions have been shown to be effective in LMIC settings, which culturally may be less stigmatizing than referral to secondary care (Clarke, King & Proust, 2013). Maternal mental health is such a neglected field internationally that taking a global perspective that highlights inequalities but also identifies simple interventions of potential benefit to all. Ultimately, accumulating and comparing evidence from across countries helps us identify socio-cultural antecedents and determinants of maternal mental health that are perhaps not visible when research is population-specific. To do so, we need to ensure more access to international research on maternal mental health and over the next year we would like to see an increase in international studies, particularly from LMICs, in the Journal of Reproductive and Infant Psychology that can contribute to this neglected knowledge base.

Professor Fiona Alderdice
Dr James Newham

Atif, N, Lovell, K, Rahman A (2015) Maternal mental health: The missing “m” in the global maternal and child health agenda Seminars in Perinatology
Volume 39, Issue 5, , Pages 345–352

Clarke K, King M and Proust A (2013) Psychosocial Interventions for Perinatal Common Mental Disorders Delivered by Providers who are not mental health specialists in low and middle income countries: a systematic review and meta-analysis PLOS Medicine 10 (10).

Fisher J, De Mello MC, Patel, V et al (2012). Prevalence and determinant of common perinatal mental disorders in women in low- and lower-middle-income countries: A systematic review. *Bull World Health Organ*, 90 (8) (2012), pp. 139G–149G

Rosaura E. Orengo-Aguayo & Lisa S. Segre Depression treatment delivered at the point-of-care: a qualitative assessment of the views of low-income US mothers. *Journal of Reproductive and Infant Psychology*

Wemakor A, Casson K and D. H (2014). "Prevalence and sociodemographic patterns of antidepressant use among women of reproductive age: A prescription database study." *Journal of affective disorders* 167: 299-305.

World Health Organisation (2008). Maternal mental health and child health and development in low and middle-income countries. Report of the WHO-UNFPA meeting held in Geneva Switzerland 30th Jan-1st Feb 2008