

Northumbria Research Link

Citation: Macdonald, L., Sayers, I., Dempsey, S., Gray, William Keith, Dodds, Steve, Hand, Annette and Walker, Richard (2019) Investigating physical and cognitive changes over two years in patients with moderate to late stage Parkinson's Disease in Northumbria. In: International Parkinson and Movement Disorder Society, 22-26th September 2019, Nice, France.

URL: <https://www.mdsabstracts.org/abstract/investigating-physical-and-cognitive-changes-over-two-years-in-patients-with-moderate-to-late-stage-parkinsons-disease-in-northumbria/>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/42907/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



**Northumbria
University**
NEWCASTLE



UniversityLibrary

INVESTIGATING PHYSICAL AND COGNITIVE CHANGES OVER TWO YEARS IN PATIENTS WITH MODERATE TO LATE STAGE PARKINSON’S DISEASE IN NORTHUMBRIA

L. MCDONALD^A, I. SAYERS^A, S. DEMPSEY^A, W. GRAY^B, S. DODDS^B, A. HAND^{B,C} AND R. WALKER^{A,B}
A. Newcastle University
B. Northumbria Healthcare NHS Foundation Trust
C. Northumbria University

Background

- Parkinson’s disease (PD) patients face both motor and non-motor symptoms, with non-motor shown to have a greater impact on quality of life¹.
- Studies suggest that the Montreal Cognitive Assessment (MoCA) is more sensitive than the Mini Mental State Exam (MMSE) for detecting cognitive decline in PD².
- Grip strength has been shown to be a better predictor of decline in function compared to gait and balance³.

Results

Physical changes:

- TUG 75/162 patients were analysed:
 - 33% had a decrease
 - 15% had an increase
 - Of the patients with no change, around half had a “problematic” TUG at baseline.
- Grip strength 56/162 patients were analysed:
 - There was an overall decline in grip strength
 - Very few showed a reduction in hand movement (11% right hand, 7% left hand). Surprisingly the majority improved (47% right, 65% left).

Figure 1: Change in TUG over two years

	Increase in TUG	Decrease in TUG	No Change
Number of sample	12	25	A: 17
			B: 21

A: “Problematic” at baseline
B: Not “Problematic” at baseline

Methods

- The Northumbria Care Needs Project is investigating the care requirements of Hoehn and Yahr score III-V PD patients. Only patients with complete baseline and follow-up data were included.
- Follow up data were collected between 23/11/16 and 29/01/19, two years after patients were seen at baseline.
- We assessed changes in timed up and go (TUG), grip strength, MoCA and MMSE scores, and explored which areas of cognition were most commonly affected. TUG was defined as normal (≤10s), good (≤20s) or problematic (>20s).
- Decline in grip strength was compared against patient’s MDS-UPDRS scores for hand movements (question 3.5).


Cognitive changes:

- 64/162 patients completed cognitive assessment.

Figure 2: Average % change in total score over 2 years

MMSE	-10%
MOCA	-39%

Figure 3: % of cohort with decreasing cognition in each section of the MoCA



Conclusion/Discussion

- Cognitive decline is more apparent on testing with the MoCA than the MMSE. Visuospatial/executive function and delayed recall are most, and naming difficulty least, commonly affected.
- The majority of patients TUG rating stayed static although this may have been due to many having already progressed to “Problematic”.
- Improvements in TUG and grip strength may be due to input from physiotherapy and medication changes.
- There was no link between decline in grip strength and assessment of hand movements by MDS-UPDRS scale.

1 Martinez-Martin P, Rodriguez-Blazquez C, Kurtis MM, Chaudhuri KR. The impact of non-motor symptoms on health-related quality of life of patients with Parkinson’s disease. Movement Disorders. 2011;26(3):399–406.
2 Nazem S, Siderowf AD, et al. Montreal Cognitive Assessment Performance in Patients with Parkinsons Disease with “Normal” Global Cognition According to Mini-Mental State Examination Score. Journal of the American Geriatrics Society. 2009; 57(2):304–8.
3 Jones GR, Roland KP, Neubauer NA, Jakobi JM. Handgrip Strength related to Long-Term Electromyography: Application for Assessing Functional Decline in Parkinson’s Disease. Archives of Physical Medicine and Rehabilitation 2017;98(2):347-352