

Northumbria Research Link

Citation: Barclay, Nicola L, Rowley, Susan, Robson, Anna, Akram, Umair and Myachykov, Andriy (2020) Sleep duration, sleep variability, and impairments of visual attention. *Quarterly Journal of Experimental Psychology*, 73 (6). pp. 868-880. ISSN 1747-0218

Published by: Psychology Press

URL: <https://doi.org/10.1177/1747021819895771>
<<https://doi.org/10.1177/1747021819895771>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/43961/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

Sleep duration, sleep variability, and impairments of visual attention

Journal:	<i>Quarterly Journal of Experimental Psychology</i>
Manuscript ID	QJE-STD 17-279.R3
Manuscript Type:	Standard Article
Date Submitted by the Author:	15-Nov-2019
Complete List of Authors:	Barclay, Nicola; University of Oxford, Nuffield Department of Clinical Neurosciences Rowley, Susan; Northumbria University, Department of Psychology Robson, Anna; Sheffield Hallam University Department of Psychology Sociology and Politics Akram, Umair; Sheffield Hallam University Department of Psychology Sociology and Politics Myachykov, Andriy; Northumbria University, Department of Psychology
Keywords:	Attention, Cognition, Sleep Deprivation, Sleep Duration, Sleep Variability

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Sleep duration ~~and~~, sleep variability, ~~predict~~ and impairments ~~in~~ of visual attention

Nicola L. Barclay^{1*}, Susan Rowley², Anna Robson³, Umair Akram³ and Andriy Myachykov^{2,4}

¹*Sleep and Circadian Neuroscience Institute, Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, UK^{‡†}*

²*Department of Psychology, Faculty of Health and Life Sciences, Northumbria University, Newcastle, UK[†]*

³*Department of Psychology, Sociology and Politics, Sheffield Hallam University, Sheffield, UK[†]*

⁴*Centre for Cognition and Decision Making, National Research University Higher School of Economics, Moscow, Russian Federation*

*Corresponding author: Dr. Nicola Barclay, Sleep and Circadian Neuroscience Institute (SCNi), Nuffield Department of Clinical Neurosciences, Sir William Dunn School of Pathology, University of Oxford, South Parks Road, Oxford, OX1 3RE, United Kingdom.

(T) 01865 618667 (E) nicola.barclay@ndcn.ox.ac.uk

[‡]Institution where data analysis and manuscript preparation were performed

[†]Institution where recruitment and data collection were carried out

Running head title: Sleep duration and visual attention

Disclosure of Interest: The authors report no conflicts of interest

Word count: ~~5475~~5564

Sleep duration, ~~sleep~~ and variability, and impairments predict impairments ~~in~~of visual attention

Abstract

Attentional networks are sensitive to sleep deprivation. However, variation in attentional performance as a function of normal sleep parameters is under-studied. We examined whether attentional performance is influenced by 1) individual differences in sleep duration; 2) sleep duration variability; and/or 3) their interaction. Fifty-seven healthy participants (61.4% female; mean age=32.37 years; SD=8.68) completed questionnaires, wore wrist actigraphy for one week, and subsequently completed the Attention Network Test. Sleep duration and sleep duration variability did not predict orienting score, executive control score or error rates. Sleep duration variability appeared to moderate the association between sleep duration with overall reaction time ($\beta = -.34$, $t = -2.13$, $p = .04$) and alerting scores ($\beta = -.43$, $t = 2.94$, $p = .01$), though further inspection of the data suggested that these were spurious findings. Time of testing was a significant predictor of alerting score ($\beta = .35$, $t = 2.96$, $p = .01$), chronotype of orienting ($\beta = .31$, $t = 2.28$, $p = .03$) and age of overall reaction time ($\beta = .35$, $t = 2.70$, $p = .01$). Our results highlight the importance of examining the associations between variations in sleep-wake patterns and attentional networks in samples with greater variation in sleep, as well as the importance of rigorously teasing apart mechanisms of the sleep homeostat from those related to the circadian rhythm in studies examining cognition.

Keywords

Attention, Cognition, Sleep Deprivation, Sleep Duration, Sleep Variability

Introduction

Insufficient sleep, in either quantity or quality, initiates a cascade of physiological events, which alter brain neurochemistry, and in consequence – our health and behaviour. Perhaps one of the most noticeable consequences of insufficient sleep is the inability to successfully focus attention and maintain an alert state. Numerous studies have highlighted the detrimental effects of experimentally induced sleep restriction or total sleep deprivation on vigilant attention (Basner & Dinges, 2011; Basner, Mollicone, & Dinges, 2011; Lim & Dinges, 2008); while other studies show that response speeds decrease in a dose-response manner as sleep duration decreases (Belenky et al., 2003; Van Dongen, Maislin, Mullington, & Dinges, 2003).

Some reports suggest that contemporary society imposes a state of chronic sleep deprivation in the Western world (Webb & Agnew, 1975). Today it is common to impose restrictions on our sleep to optimise our time spent awake, but this comes at a cost to our health, mood, occupational safety, and cognition (Cappuccio, D'Elia, Strazzullo, & Miller, 2010; Fernandez-Mendoza et al., 2010; Kalmbach, Arnedt, Song, Guille, & Sen, 2017). An epidemiological study demonstrated that the typical sleep duration of young adults during weekdays is around 6.7 hours (Breslau, Roth, Rosenthal, & Andreski, 1997), which is below the 8 hours recommended for this age group (Hirshkowitz et al., 2015). In addition to suboptimal sleep duration, adherence to consistent and biologically compatible sleep schedules are often impinged by external environmental factors such as work commitments, social engagements and 24/7 electronic stimulation (Wittmann, Dinich, Mellow, & Roenneberg, 2006). Indeed, it is common to see a mismatch between sleep obtained on work-days and that obtained on free-days, which is indicative of a state of “social jetlag”, imposed by our social world (Wittmann et al., 2006). On work-days, constrained sleep offset times often result in drastically curtailed sleep duration - a sleep “debt” that is paid off on free-days

1
2
3 by extending sleep where the opportunity permits. However, employing compensatory
4 behaviours, such as extension of time in bed on free-days, and going to bed earlier than what
5 would be dictated by the endogenous circadian rhythm on work-days, is maladaptive (Buysse
6 et al., 2010). This coupling of short sleep duration and self-imposed variability in sleep
7 behaviours may disrupt the homeostatic drive for sleep, making it difficult to get to sleep
8 when desired, and potentially contributing to a delayed circadian phase. Consequently, we
9 may develop a sleep schedule at conflict with the constraints of the social world.
10
11
12
13
14
15
16
17
18
19

20 Several studies have shown that intra-individual variability in sleep duration is often
21 greater than that observed between individuals (Knutson, Rathouz, Yan, Liu, & Lauderdale,
22 2007; Tworoger, Lee, Schernhammer, & Grodstein, 2006) and it is possible that sleep
23 duration variability contributes to the development or maintenance of sleep disorders
24 (Billiard, Alperovitch, Perot, & Jammes, 1987; Buysse et al., 2010; Spielman, Caruso, &
25 Glovinsky, 1987). Indeed, individuals with insomnia often experience night-to-night
26 variability across various domains assessed subjectively by sleep diaries, and objectively – by
27 actigraphy (Buysse et al., 2010; Vallières, Ivers, Bastien, Beaulieu-Bonneau, & Morin, 2005;
28 Vallières, Ivers, Beaulieu-Bonneau, & Morin, 2011). Difficulty sleeping on one night may
29 increase sleep pressure such that the subsequent sleep period is longer, deeper and more
30 restorative. In some cases, several nights of poor sleep may be followed by a night of
31 relatively good sleep (Vallières et al., 2005; Vallières et al., 2011), though this pattern is not
32 predictably consistent (Buysse et al., 2010).
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50 Whilst most research on the consequences of insufficient sleep have focussed on short
51 sleep duration or sleep disruption, the importance of *consistent sleep schedules* is often
52 overlooked. Sleep variability has been broadly defined as inconsistencies in various sleep
53 indices including sleep onset and offset timing, sleep latency, sleep quality, wake after sleep
54 onset, sleep duration and efficiency. It is typically calculated as the variability from weekdays
55
56
57
58
59
60

1
2
3 to weekends (Lemola, Schwarz, & Siffert, 2012), or as the within-subject standard deviation
4
5 for the corresponding variables over the course of several days (Whiting & Murdock, 2016)
6
7 to weeks (Lemola, Ledermann, & Friedman, 2013; Lev Ari & Shulman, 2012;
8
9 Sánchez-Ortuño & Edinger, 2012).

10
11
12
13 Greater sleep variability has been negatively associated with health (e.g. higher body
14
15 mass index: Moore et al., 2011), mood (e.g. poorer subjective well-being: Lemola et al.,
16
17 2013; and increased severity of depression: Suh et al., 2012), and behaviour (e.g. adolescent
18
19 aggression: Lemola et al., 2012). Of particular relevance to the current study, Whiting and
20
21 Murdock (2016) demonstrated that sleep duration variability moderates the association
22
23 between sleep duration and attentional disengagement; i.e., the capacity to withdraw attention
24
25 from the currently focused stimulus (attentional switch). Shorter sleep duration was
26
27 associated with poorer ability to switch attention when coupled with low sleep duration
28
29 variability. Therefore, consistently short sleep duration was associated with greatest deficits
30
31 in attentional disengagement, and consistently longer sleep duration was associated with
32
33 better performance.
34
35
36
37
38

39
40 The majority of work on visual attention has used the ‘gold standard’ measure of the
41
42 Psychomotor Vigilance Test (PVT), which has consistently been shown to be sensitive to
43
44 sleep deprivation (see Tkachenko & Dinges 2018, for a review). However, visual attention is
45
46 not limited to our ability to switch focus between stimuli; it is a complex system of
47
48 functionally and anatomically distinct brain networks, which support our ability to (i)
49
50 maintain an “alert” state (alerting network), (ii) “orient” attention to stimuli (orienting
51
52 network), and (iii) resolve conflict when numerous stimuli simultaneously compete for
53
54 attention (executive control network) (Fernandez-Duque & Posner, 2001; Petersen & Posner,
55
56 2012). These networks are governed by distinct neurobiological pathways (Fan & Posner,
57
58 2004) and they are individually supported by network-specific genetic modulators (Fossella
59
60

1
2
3 et al., 2002), making it likely that they are also differentially affected by sleep. Indeed,
4
5 previous research has registered differential effects of sleep deprivation on the functioning of
6
7 these distinct networks (see Tkachenko & Dinges, 2018). For example, one study compared
8
9 ANT performance at baseline vs. following 24-hours of sleep deprivation within subjects and
10
11 observed longer reaction times, poorer accuracy, and diminished P3 event-related potential
12
13 response following sleep deprivation (Trujillo, Kornguth, & Schnyer, 2009). Similarly,
14
15 another study demonstrated longer reaction times and impaired functioning of the orienting
16
17 and executive control networks following sleep deprivation compared to baseline (Martella,
18
19 Casagrande, & Lupianez, 2011); though these effects were only partially replicated by some
20
21 researchers (Jugovac & Cavallero, 2012) and not replicated by others (Muto et al., 2012;
22
23 Roca et al., 2012).

24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Despite this growing body of research on the relationships between *experimentally*
induced sleep deprivation and the performance of attentional networks, a dearth of research
focuses on what happens in the real world. Our own previous work demonstrated that
impairments in attentional performance emerge after 18-hours of sustained wakefulness – a
period of wakefulness, which is typical of many adults during the working week (Barclay &
Myachykov, 2017). Given that today’s society may foster inconsistent sleep patterns due to
the changing responsibilities on work- and free-days, it appears pertinent to examine in detail
the influence of sleep duration variability on attention, and to determine whether sleep
duration and its variability have differential effects on distinct attentional networks.

Thus, the present study investigated associations between sleep duration over one
week assessed by actigraphy and the efficiency of the attentional networks, as well as the
possibility that these associations are moderated by sleep duration variability. We
hypothesised that poorer attentional performance would be predicted by 1) shorter sleep
duration and greater sleep duration variability, independently; and 2) consistently

1
2
3 shorter/inconsistently longer sleep duration across the week. Given previously observed
4
5 chronotype and time-of-day effects on attentional performance (e.g. Barclay & Myachykov,
6
7 2017; Matchock & Mordkoff, 2009), analyses accounted for their potential direct and
8
9 interactive effects where relevant.
10

11 12 13 **Method**

14 15 16 *Participants*

17
18
19 Participants were recruited from the general population of England through poster
20
21 advertisements, emails to staff and students of Northumbria University, Sheffield Hallam
22
23 University and University of Oxford, and through social media. To be eligible to participate,
24
25 potential participants had to be between 22-50 years of age; not taking any medications that
26
27 may affect their sleep; not have a history of/or current medical, neurological or psychiatric
28
29 illness (including anxiety or depression); not have a sleep disorder (treated or untreated); and
30
31 not be a shift worker. Fifty-eight participants meeting these eligibility criteria initially
32
33 enrolled in the study; 57 provided complete data (61.4% female; mean age = 32.37 years; SD
34
35 = 8.68; 61.4% female).
36
37
38
39
40

41 *Measures*

42 43 44 *Screening questionnaire*

45
46
47 An in-house screening questionnaire assessing sleep, sleep disorders, general health and
48
49 demographic information assessed eligibility for participation.
50
51

52 53 *Sleep quality*

54
55
56 Sleep quality over the past month was assessed using the Pittsburgh Sleep Quality Index
57
58 (PSQI: Buysse, Reynolds, Monk, Berman, & Kupfer, 1989). The PSQI is an 18-item self-
59
60

1
2
3 report questionnaire, which assesses 7 components of sleep quality: subjective sleep quality,
4 sleep latency, sleep duration, sleep efficiency, sleep disturbances, use of sleeping medication,
5 and daytime dysfunction. Component scores are summed to produce a global sleep quality
6 score ranging from 0 to 21. Higher scores represent poorer sleep quality. Global sleep quality
7 score was considered as a covariate in the statistical analyses if it was shown to be associated
8 with the dependent variable of interest.
9

10 11 12 13 14 15 16 17 18 *Chronotype*

19
20
21 Chronotype was assessed using the Morningness-Eveningness Questionnaire (MEQ: Horne
22 & Östberg, 1976). The MEQ is a 19-item self-report questionnaire, which assesses preferred
23 timing of daytime activities, sleep habits, hours of peak performance and times of ‘feeling
24 best’ and maximum alertness. Responses are combined to provide a total score ranging from
25 16 to 86. Higher scores represent a tendency towards morningness. Chronotype score was
26 considered as a covariate in the statistical analyses if it was shown to be associated with the
27 dependent variable of interest.
28
29
30
31
32
33
34
35
36
37

38 *Sleepiness*

39
40
41 State sleepiness was assessed using the Stanford Sleepiness Scale (SSS: Hoddes, Zarcone, &
42 Dement, 1972). The SSS is a 1-item measure assessing momentary alertness. Responses
43 range from 1 (wide awake) to 7 (sleep onset soon). Sleepiness was considered as a covariate
44 in the statistical analyses if it was shown to be associated with the dependent variable of
45 interest.
46
47
48
49
50
51
52
53

54 *Actigraphy*

55
56
57 Actilife GT3X and CamNtech Ltd. MW8 actiwatches were used to detect movement bouts,
58 worn for seven consecutive nights on the non-dominant wrist. Data collection started
59
60

1
2
3 immediately following the initial laboratory visit and concluded on the morning of day 8.
4
5 Actilife 5 (Actigraph Corp, 2011) and Motionware 1.1.25 (CamNtech Ltd. 2009) software
6
7 was used to analyse the actigraphy data to provide summary statistics of the participants'
8
9 weekly sleep. We used the Sadeh Algorithm (Sadeh, Sharkey, & Carskadon, 1994) to
10
11 determine sleep/wake status for each 60 second (actilife) or 30 second (CamNtech) epoch
12
13 with 30Hz sample rate, as per device recommendations for optimal sleep recording (note that
14
15 there were no differences between device type on any of the independent variables or
16
17 outcomes of interest). Participants also kept daily sleep diaries, indicating bed-time and rise-
18
19 time and these times were entered into the software as start times for actigraphic analysis (i.e.
20
21 to indicate intention to sleep). Sleep duration was averaged for each participant across the
22
23 seven nights. Variability in sleep duration was calculated as the within-participant standard
24
25 deviation in sleep duration across the seven nights.
26
27
28
29
30

31 *Attention*

32
33
34
35 The Attention Network Test (ANT: Fan, McCandliss, Sommer, Raz, & Posner, 2002) is a
36
37 computerised reaction time task used to examine the individual attentional networks'
38
39 performance (see Figure 1). In the ANT, participants perform on trials containing centre,
40
41 double, spatial cues, or no cues (100msec) between two central fixation events. At the second
42
43 central fixation (400msec), the target arrow (left or right) is presented either above or below
44
45 the fixation cross, and is either presented alone (neutral condition); with 2 flankers either side
46
47 pointing in the same direction (congruent condition); or with 2 flankers either side pointing in
48
49 the opposite direction (incongruent condition) (lasting no longer than 1700msec). Upon
50
51 presentation of the target, participants are required to indicate by pressing designated keys on
52
53 a computer keyboard whether the corresponding arrows point leftwards or rightwards. As
54
55 outlined by us previously (Barclay & Myachykov, 2017), the ANT provides a raw reaction
56
57
58
59
60

1
2
3 time (RT) measure for each of the conditions (cue type: no cue, centre cue, double cue,
4 spatial cue; flanker type: neutral, congruent, incongruent) as well as error rates. Additionally,
5
6 the ANT provides specific measures of alerting, orienting and conflict resolution (executive
7 control). The alerting score is calculated by subtracting the mean RT of the double-cue
8 conditions (which alerts the participant to the imminent target, but provides no information
9 on its location either above or below the cross) from the mean RT of the no-cue conditions.
10
11 The orienting score is calculated by subtracting the mean RT of the spatial cue conditions
12 (which alerts participants to the imminent target and provides information on its location)
13 from the mean RT of the centre cue conditions (which only alerts participants to the
14 imminent target at one location). The conflict (executive control) score is calculated by
15 subtracting the mean RT of all congruent flanked conditions from all incongruent flanked
16 conditions (from all cue types). Lower scores typically indicate difficulty: a) maintaining
17 alertness without a cue (alerting); b) disengaging from the centre cue (orienting); or c)
18 resolving conflict (executive control) (Fan & Posner, 2004).
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

35
36 [{\(Insert Figure 1 here\)}](#)
37
38

39 ***Procedure***

40
41
42 Interested participants completed the screening questionnaire to confirm eligibility. Eligible
43 participants met with the researcher to provide their informed consent, and to receive an
44 actiwatch and sleep diary as well as information regarding their use each day/night.
45
46 Participants wore actiwatches for seven consecutive nights, and they were informed that the
47 week should be 'typical' for them. Participants returned to the laboratory on day 8 to hand
48 back the actiwatch and sleep diary; to complete the PSQI, MEQ and SSS; and to participate
49 in the ANT. Note that all participants completed the ANT between 8:00am and 8:00pm (57%
50 completed the ANT before 12:00pm and 89% before 6:00pm). Time of testing was included
51
52
53
54
55
56
57
58
59
60

1
2
3 in statistical analyses as a covariate. All procedures complied with the ethical principles laid
4
5 out by the American Psychological Association and the Declaration of Helsinki. Approval
6
7 was granted by Oxford Central University Research Ethics Committee and Northumbria
8
9 University Psychology Department Ethics Committee.
10
11
12
13
14
15

16 *Statistical Analyses*

17
18
19 Previous researchers observed a large effect size for change in R^2 for sleep duration
20
21 variability predicting one facet of attention (Whiting & Murdock, 2016). In the present study,
22
23 a power calculation using G*Power indicated that to identify large effects as hypothesised
24
25 (i.e. F^2 for $\Delta R^2 \geq .35$), 47 participants would be required to achieve 90% power with an $\alpha < .01$
26
27 (Cohen, 1988). Our final sample of 57 participants provided sufficient power to address our
28
29 hypotheses. Descriptive statistics were first derived followed by t-tests to examine possible
30
31 sex differences in the dependent and independent variables. Correlations between dependent
32
33 variables, independent variables and potential covariates were then examined. A repeated-
34
35 measures within-subjects ANOVA was performed to examine the sensitivity of the ANT in
36
37 the current sample. A series of hierarchical regression models were then run for each
38
39 dependent variable (overall mean reaction time $\{RT\}$, alerting, orienting and executive
40
41 control scores, and overall error rates). Covariates that were significantly correlated with the
42
43 outcome variables were incorporated in regression models in the first steps (age, chronotype
44
45 and time of testing). In each regression model, predictor variables and covariates were grand
46
47 mean centred. A reciprocal transformation was applied to the error rate data to reduce
48
49 positive skew. Simultaneous regression models were also run to derive individual coefficients
50
51 for each predictor/covariate for significant models.
52
53
54
55
56
57
58
59
60

1
2
3 We examined five models in separate blocks as follows: 1) covariate 1 (if applicable); 2)
4 covariate 2 (if applicable); 3) sleep duration; 4) sleep duration variability; and 5) sleep
5 duration X sleep duration variability interaction. Moderation models were run using the
6 PROCESS tool for SPSS (Hayes, 2013) in order to further examine significant interactions.
7
8 Of note, linear regression models were also run with covariates entered in the last steps in
9 order to examine potential mediation effects. Mediation effects were not indicated for any
10 models (models not reported, available upon request). Significance of each model, and each
11 predictor within each model, was considered at a Bonferroni corrected level of $p < .01$ (.05/5)
12 to account for the multiple testing of ANT-derived dependent variables.
13
14
15
16
17
18
19
20
21
22
23

24 **Results**

25
26
27 Mean sleep quality score was 4.39 (SD=2.24) indicating good sleep, and chronotype score
28 was 54.47 (SD=10.43) indicating a largely ‘neither’ type chronotype. Mean actigraphically
29 measured sleep duration across the week was 421.45 mins (SD=39.70; range 336.71 mins to
30 541.71 mins), and mean variability in sleep duration across the week was 60.99 mins
31 (SD=38.42; range 16.22 mins to 256.14 mins). Mean ANT reaction times were pooled from
32 all correct trials for all participants. Incorrect trials accounted for 3.65% of the total trials.
33
34 Additionally, trials with RT <200ms and >4 absolute deviations from the median (MAD, see
35 Leys, Ley, Klein, Bernard & Licata, 2013, for information) were excluded from analysis
36 (2.87% of correct trials). In total, 6.41% of trials were excluded. Table 1 shows the mean RT
37 and SD, and Table 2 the mean error rates and SD, for each of the experimental conditions (4
38 x cue type; 3 x flanker type).
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53

54 [{\(Insert Tables 1 and 2 here\)}](#)

55
56
57 There were sex differences in actigraphically derived mean sleep durations ($t(55) =$
58 -2.89 , $p < .05$, [Hedges’ \$g = .79\$](#)). Males had significantly shorter mean sleep duration
59
60

(mean=403.44 mins; SD=34.39) than females (mean=432.76 mins, SD=39.03). There were also significant sex differences in orienting, ($t(55) = -2.35$, $p < .05$, Hedges' $g = .64$). Orienting score for males was significantly lower (mean=22.11; SD=17.63) than females (mean=33.94; SD=19.00) suggesting that females were faster to orient their attention. There were no sex differences in any of the remaining key actigraphically assessed sleep variables or key ANT variables (sleep duration variability: $t(55) = .95$, $p = .35$; overall RT: $t(55) = -.54$, $p = .59$; alerting: $t(55) = .55$, $p = .59$; executive control: $t(55) = .16$, $p = .87$; and overall error rates: $t(55) = .04$, $p = .97$). There were several significant correlations that are fundamental to the analyses (see Table 3): 1) older age and longer overall RTs; 2) lower sleep duration variability and poorer alerting scores; 3) older age and poorer alerting scores; 4) earlier time of testing and poorer alerting scores; 5) younger age and poorer orienting scores; and 6) increasing tendency towards eveningness and poorer orienting scores.

{(Insert Table 3 here)}

A repeated-measures ANOVA was performed on mean overall RTs, with cue (no cue, centre, double and spatial cues) and flanker type (neutral, congruent and incongruent) as within-subject factors. Assumptions of sphericity were not met for the main effects of cue and flanker type, and the interaction between cue and flanker type. Consequently, Greenhouse-Geisser correction to degrees of freedom was employed. There were significant main effects of cue, ($F(2.50, 137.52) = 176.27$, $p < .01$, $\eta_p^2 = .76$) and flanker type, ($F(1.45, 79.82) = 1040.91$, $p < .01$, $\eta_p^2 = .95$) on mean overall RT, and a significant interaction between cue and flanker type, ($F(4.37, 240.51) = 12.92$, $p < .05$, $\eta_p^2 = .19$). Longer reaction times were registered in trials with no cue vs. all other cues (relevant to alerting and orienting); and incongruent vs. both other flanker types (relevant to executive control).

1
2
3 A series of hierarchical linear regression models were run separately for each outcome
4 variable (see Tables 4-6). For mean overall RT as the dependent variable, model 1 (including
5 age) was significant, ($F(1,56)=9.03, p=.00$), and model 4 (including age, sleep duration,
6 sleep duration variability and the sleep duration X sleep duration variability interaction)
7 significantly improved model fit, ($\Delta F(1,52)=4.52, p=.04$). The final model explained 22%
8 of variance in overall RT (see Table 4). From the simultaneous regression model, age was a
9 significant predictor of overall RT, such that older age predicted longer overall RTs ($\beta=.35,$
10 $t=2.70, p=.01$), as was the sleep duration X sleep duration variability interaction ($\beta=-.34, t=-$
11 $2.13, p=.04$). Moderation analysis in PROCESS revealed that the slowest RTs were
12 predicted by shorter sleep duration and high sleep duration variability, and that the fastest
13 RTs were predicted by longer sleep duration and high sleep duration variability ([t values for](#)
14 [each moderator level: low: \$t=.36, p=.72\$; medium: \$t=-1.11, p=.27\$; high: \$t=-2.21, p=.03\$](#)) (see
15 Figure 2a).

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34 [{\(Insert Table 4 here\)}](#)
35
36

37 For alerting score as the dependent variable, Models 2 (including age and time of
38 testing) and 5 (including age, time of testing, sleep duration, sleep duration variability and the
39 sleep duration X sleep duration variability interaction) significantly improved model fit,
40 ($\Delta F=10.32, p=.00$; and $\Delta F=8.62, p=.01$, respectively). The final model explained 37%
41 of variance in alerting score. From the simultaneous regression model, time of testing and the
42 sleep duration X sleep duration variability interaction significantly predicted alerting score
43 ($\beta=-.22, t=-2.96, p=.01$; and $\beta=.43, t=2.94, p=.01$). Moderation analysis in PROCESS
44 revealed that the poorest alerting score was predicted by longer sleep duration and low sleep
45 duration variability, and that the best alerting score was predicted by longer sleep duration
46 and high sleep duration variability ([t values for each moderator level: low: \$t=-1.15, p=.25\$;](#)
47 [medium: \$t=.53, p=.60\$; high: \$t=2.17, p=.03\$](#)) ($t=2.54, p=.01$) (see Figure 2b). Because of the
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 strength of the correlation between sleep duration variability and alerting score (Table 3), we
4 decided to examine the extent to which sleep duration moderates this association. We
5 performed a median split on the sleep duration variable and performed moderation analysis in
6 PROCESS. It appeared that long sleep duration drives the association between sleep duration
7 variability and alerting score (t value for short and long sleep duration: short: $t=-.57=2.03$,
8 $p=.57$; long: $t=3.26$, $p=.0004$) (see Figure 2c).

9
10
11
12
13
14
15
16
17
18 [{\(Insert Table 5 and Figures 2a, 2b and 2c here\)}](#)
19
20

21 Due to the unexpected direction of the interaction effect (i.e. that long sleep duration
22 with low sleep duration variability predicted poorest alerting), we further visualised the
23 distribution of sleep duration variability and alerting score (see Figure 3a). The presence of
24 an outlier experiencing high sleep duration variability and high alerting score is present in the
25 data (256.14 mins and alerting score of 110.8). Indeed, inspection of the nightly sleep
26 duration data for this individual revealed nightly sleep durations ranging from 142 mins to
27 924 mins, the latter of which occurred the night prior to testing. Inspection of the distribution
28 of sleep duration the night prior to testing and alerting score also revealed the contribution of
29 this outlier to the association (Figure 3b). The average sleep duration experienced by this
30 participant was not unusual (501.43 mins); hence the distribution of mean sleep duration and
31 alerting appears normal (Figure 3c). Due to the presence of this outlier, we re-ran the
32 correlations after excluding this participant. The association between sleep duration
33 variability and alerting reduced from .34 ($p<.01$) to -.03 ($p=.82$); and the association between
34 sleep duration the night prior to testing and alerting reduced from .30 ($p<.05$) to -.06 ($p=.65$).
35 The regression analyses were re-run after exclusion of this participant, and the pattern of
36 results for overall RT as the dependent variable changed such that the interaction term was no
37 longer significant (though age remained a significant predictor of overall RT ($\beta=.36$,
38 $t=2.65$, $p=.01$); likewise for alerting as the dependent variable the interaction between sleep
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 duration X sleep duration variability reduced to non-significance, whilst time of testing
4
5 remained a significant predictor ($\beta=.37, t=2.85, p=.01$).
6
7

8
9 For orienting score as the dependent variable, model 1 (including age) and model 2
10 (including age and chronotype) significantly improved model fit ($\Delta F=4.73, p=.03$; and
11 $\Delta F=5.54, p=.02$, respectively). Model 2 explained 16% of variability in orienting score.
12
13 From the simultaneous regression model, only chronotype significantly predicted orienting
14
15 score ($\beta=.31, t=2.28, p=.03$) such that a tendency towards morningness was associated with
16
17 better orienting. These results remained after excluding the anomalous participant.
18
19
20
21
22

23 [{\(Insert Table 6 here\)}](#)
24
25

26 For executive control scores and overall error rates, all regression analyses were non-
27
28 significant (all model p 's $>.05$) (both with the inclusion and exclusion of the anomalous
29
30 participant).
31
32

33 [{\(Insert Figure 3 here\)}](#)
34
35
36

37 **Discussion**

38
39

40 The results presented here expand previous research investigating the importance of
41
42 sleep for optimum cognitive functioning. Here we extend research from experimental sleep
43
44 deprivation studies, to demonstrate the influence of normal sleep duration variability in the
45
46 general population on various components of visual attention. There are four key findings
47
48 stemming from this study: 1) neither sleep duration, nor sleep duration variability, nor their
49
50 interaction, were predictive of orienting, executive control or error rates; 2) earlier time of
51
52 testing was associated with poorer efficiency of the alerting network; 3) the associations
53
54 between sleep duration with overall reaction time and efficiency of the alerting network
55
56 *appeared* to be moderated by sleep duration variability; however, the presence of an outlier
57
58
59
60

1
2
3 experiencing high sleep duration variability and extremely long sleep duration the night prior
4 to testing appeared to solely account for the interactive effects of sleep duration and sleep
5 duration variability on overall reaction time/ alerting score; and 4) older age was associated
6 with longer overall reaction times, and this was independent of sleep duration and sleep
7 duration variability. Below we explain these findings in relation to our initial hypotheses and
8 in the context of the emerging field.
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

24
25 ***Sleep duration and sleep duration variability did not predict orienting score, executive***
26 ***control score or error rates***
27
28
29

30 Contrary to our hypotheses, we did not find that sleep duration or sleep duration
31 variability predicted components of attention (the efficiency of the orienting network, the
32 efficiency of the executive control network, or proportion of errors made). Likewise, the
33 interaction between sleep duration and sleep duration variability was not a significant
34 predictor of these measures of attention. Other experimental studies show a general detriment
35 to sustained attention under varying degrees of sleep deprivation, fragmentation and
36 restriction (see Tkachenko & Dinges, 2018 for a review) while observational studies similarly
37 show that consistently short sleep duration is associated with attentional disengagement
38 (Whiting & Murdock, 2016). Our data does not show a similar effect, at least in terms of
39 orienting, executive control and accuracy.
40
41
42
43
44
45
46
47
48
49
50
51
52
53

54 There are a couple of plausible explanations for this. First, it is possible that these
55 particular ANT-derived variables are insensitive to natural variation in sleep duration and
56 intra-individual sleep duration variability. The experimental sleep-deprivation literature
57
58
59
60

indeed suggests that whilst sustained attention (which would be analogous to our measure of overall reaction time) and alerting are particularly sensitive to sleep loss, other attentional functions, including orienting and executive function, may remain spared (Cunningham, Jones Eskes & Rusak, 2018; Roca et al., 2012; Tkachenko & Dinges, 2018, but see Martella, Casagrande, & Lupianez, 2011). Second, it is possible that there was not enough variability in our data for such effects to emerge. Indeed, we sampled the average sleep duration over the course of 7 days in healthy good sleepers, and asked them to maintain their ‘habitual’ sleep patterns. The standard deviation of our ‘mean sleep duration’ variable was 39.90 minutes, and 86% of our sample had a mean sleep duration between 6-8 hours. Thus, it is possible that meaningful effects on attentional performance occur at far greater extremes of sleep duration than the ones exhibited by our sample. Studies examining consequences of sleep duration variation often focus on greater extremes, and demonstrate a U-shaped curve: highlighting the detrimental effects of both short (conceptualised as <6 hours per night) and long (usually conceptualised as >8 hours per night) sleep duration on various indicators of health (Patel & Hu, 2008), all-cause mortality (Cappuccio et al., 2010), and cognition (Kronholm et al., 2009). Our intention was to show that in the ‘real-world’ consistently short sleep or inconsistently long sleep results in deficits in attention, yet it is possible that our selection process yielded a sample of healthy good sleepers that failed to allow these patterns to emerge. Future research needs to encapsulate more variable sleep-wake patterns to truly describe the heterogeneity of sleep (by including those with sleep disturbances/more variable sleep-wake patterns such as shift workers) that exists in the general population and examine concomitant associations with cognition.

Third, effects of sleep duration and sleep duration variability may have been obscured by the effect of time of testing. Whilst the majority of our participants completed the ANT before midday, we did not strictly control time of testing, and so testing was performed up

1
2
3 until 8pm. Had time of testing been consistent, we may have seen an effect of sleep duration
4 emerge. Matchock and Mordkoff (2014) for example, observed increased reaction times on
5 the ANT *upon awakening* under conditions of restricted sleep durations of 3-hours and 6-
6 hours, compared to a baseline condition of normal sleep duration. However, the results of
7 Matchock & Mordkoff may actually reflect “sleep inertia” or result from testing at
8 suboptimal circadian phase in the early morning, rather than purely the effect of short sleep
9 duration. We chose not to restrict our testing time to *upon awakening* as we wanted to rule
10 out the effect of sleep inertia, and we wanted to examine potential effects of time of testing.
11
12
13
14
15
16
17
18
19
20
21

22
23 Nevertheless, our data revealed a reliable effect of time of testing on the efficiency of
24 the alerting network. The alerting network efficiency was progressively better as the time of
25 testing was later in the day. This finding is partially consistent with the studies that observed
26 significant time-of-day effects on the alerting, orienting and executive control networks
27 (Matchock & Mordkoff, 2009; Fimm, Brand & Spijkers, 2016), and corroborates what is
28 typically observed in relation to the circadian rhythm of neurobehavioural functioning: it is
29 poor upon awakening in the morning, and then steadily improves across the waking day to a
30 peak around early evening, followed by a progressive decline into the night (Goel et al.,
31 2011; Mollicone et al., 2010). Taken together, these findings highlight the importance of
32 systematically isolating variables relating to the sleep homeostat and the circadian rhythm.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Associations between sleep duration with overall reaction time and efficiency of the alerting network appeared to be moderated by sleep duration variability

Contrary to our hypotheses, shorter sleep duration coupled with high sleep duration variability predicted the longest overall reaction times, and longer sleep duration coupled

1
2
3 with high sleep duration variability predicted the shortest reaction times. Concerning alerting,
4 longer sleep duration coupled with low sleep duration variability appeared to be associated
5
6 with poorer efficiency of the alerting network, whilst on the other hand longer sleep duration
7
8 coupled with high sleep duration variability appeared to be associated with better efficiency
9
10 of the alerting network. However, further inspection of our data revealed that these effects
11
12 were accounted for by an individual exhibiting extremely high sleep duration variability, and
13
14 extremely long sleep duration the night prior to testing – with this participant excluded, these
15
16 interaction effects were reduced to non-significance. This finding highlights the importance
17
18 of considering the nightly variability in sleep duration when taking an average of such data.
19
20 Many research studies rely on taking an average of sleep variables across a pre-specified time
21
22 period (3 nights, a week), but such crude measures fail to capture the nightly variability in
23
24 sleep duration. With this participant excluded, the standard deviation of our sleep duration
25
26 variability predictor reduced from 38.41 to 28.24. Thus, our data were perhaps too
27
28 homogenous for important effects on reaction time and alerting to emerge.
29
30
31
32
33
34
35

36 *Association between Age And Overall Reaction Time:*

37
38 Perhaps less surprising in our study was the finding that older age was associated with
39
40 longer overall reaction time. It appears that the relative deficit in attention shown here stems
41
42 from age-related factors rather than short sleep duration per se. The efficiency of the alerting
43
44 network is significantly impaired in older adults comparative to younger adults (Gamboz,
45
46 Zamarian, & Cavallero, 2010; Jennings, Dagenbach, Engle, & Funke, 2007). Given that sleep
47
48 patterns dramatically change across the lifespan (Ohayon, Carskadon, Guilleminault, &
49
50 Vitiello, 2004), it is pertinent to further investigate the interactive effects of aging and sleep
51
52 on cognition. Others have hypothesised that increased reaction time following sleep
53
54 restriction results from diminished oculomotor functioning (Zils, Sprenger, Heide, Born, &
55
56
57
58
59
60

1
2
3 Gais, 2005), and this effect may become more pronounced as we age, given the natural
4 progression of oculomotor functioning across the lifespan (Katsanis, Iacono, & Harris, 1998).
5
6
7

8 ***Limitations and Conclusions:***

9
10 It is worth noting a couple of key limitations of the present study: 1) Our analyses focus on
11 sleep duration and sleep duration variability rather than other indices of sleep, such as sleep
12 onset latency, sleep efficiency, wake after sleep onset and number of awakenings. However,
13 our sample size did not allow us to assess these variables at an acceptable level of power.
14
15 Additionally, we wanted to focus on a variable that is to some extent under voluntary control,
16 i.e., we can control the time we go to bed and the time we get out of bed, and by consequence
17 our sleep duration, whereas we cannot control our sleep onset latency. We wanted to address
18 a variable that can be behaviourally manipulated, and thus be the potential target of
19 interventions to improve sleep and, by consequence, cognition. Further investigation of sleep
20 under normal conditions (rather than the artificial setting of experimental sleep restriction/
21 deprivation), investigating an array of sleep characteristics in a more heterogeneous sample
22 will further shed light on the role of sleep duration and its variability on attention. 2) Our
23 objective measure of sleep using actigraphy is not completely free of subjective biases, as the
24 calculation of sleep parameters is partially dependent on subjectively reported bed times and
25 rise times. Measuring sleep by using polysomnography would not only address this issue, but
26 would also provide us with the opportunity to examine architectural properties of sleep
27 macro- and micro-structure to identify the residual neural underpinnings of attention.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50 To conclude, this work did not demonstrate an effect of actigraphically measured
51 sleep duration or sleep duration variability on measures of visual attention. However, this
52 conclusion is derived from data focussing on the healthy population, experiencing good sleep.
53 We do not rule out the possibility that poor sleep and inconsistencies in sleep patterns
54 contribute to cognitive impairments in populations for whom sleep is disturbed.
55
56
57
58
59
60

Acknowledgements

The authors thank the participants for their participation. A part of this study was carried out as an MSc project by author SR. The study was supported by Northumbria University Psychology Department, internal funds from the Sleep and Circadian Neuroscience Institute, University of Oxford and by the Russian Academic Excellence Project ‘5-100’.

References

- Actigraph Corp. (2011). ActiLife 5 - User's Manual. *Prepared by ActiGraph R&D and Software Departments.*
- Barclay, N. L., & Myachykov, A. (2017). Sustained wakefulness and visual attention: moderation by chronotype. *Experimental Brain Research, 235*, 57-68.
- Basner, M., & Dinges, D. F. (2011). Maximizing sensitivity of the psychomotor vigilance test (PVT) to sleep loss. *Sleep, 34*, 581-591.
- Basner, M., Mollicone, D., & Dinges, D. F. (2011). Validity and sensitivity of a brief psychomotor vigilance test (PVT-B) to total and partial sleep deprivation. *Acta Astronautica, 69*, 949-959.
- Belenky, G., Wesensten, N. J., Thorne, D. R., Thomas, M. L., Sing, H. C., Redmond, D. P., et al. (2003). Patterns of performance degradation and restoration during sleep restriction and subsequent recovery: a sleep dose-response study. *Journal of Sleep Research, 12*, 1-12.
- Billiard, M., Alperovitch, A., Perot, C., & Jammes, A. (1987). Excessive daytime somnolence in young men: prevalence and contributing factors. *Sleep, 10*, 297-305.
- Breslau, N., Roth, T., Rosenthal, L., & Andreski, P. (1997). Daytime sleepiness: an epidemiological study of young adults. *American Journal of Public Health, 87*, 1649-1653.
- Buysse, D. J., Cheng, Y., Germain, A., Moul, D. E., Franzen, P. L., Fletcher, M., et al. (2010). Night-to-night sleep variability in older adults with and without chronic insomnia. *Sleep Medicine, 11*, 56-64.
- Buysse, D. J., Reynolds, C. F., Monk, T. H., Berman, S. R., & Kupfer, D. J. (1989). The Pittsburgh Sleep Quality Index: A new instrument for psychiatric practice and research. *Psychiatry Research, 28*, 192-213.

- 1
2
3 Cappuccio, F. P., D'Elia, L., Strazzullo, P., & Miller, M. A. (2010). Sleep duration and all-
4
5 cause mortality: a systematic review and meta-analysis of prospective studies. *Sleep*,
6
7 33, 585-592.
8
9
10 Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed.). New
11
12 Jersey: Lawrence Erlbaum.
13
14 Cunningham, J. E., Jones, S. A., Eskes, G. A., & Rusak, B. (2018). Acute sleep restriction
15
16 has differential effects on components of attention. *Frontiers in Psychiatry*, 9.
17
18 Fan, J., McCandliss, B. D., Sommer, T., Raz, A., & Posner, M. I. (2002). Testing the
19
20 efficiency and independence of attentional networks. *Journal of Cognitive*
21
22 *Neuroscience*, 14, 340-347.
23
24
25 Fan, J., & Posner, M. (2004). Human attentional networks. *Psychiatrische Praxis*, 31, 210-
26
27 214.
28
29
30 Fernandez-Duque, D., & Posner, M. I. (2001). Brain imaging of attentional networks in
31
32 normal and pathological states. *Journal of Clinical and Experimental*
33
34 *Neuropsychology*, 23, 74-93.
35
36
37 Fernandez-Mendoza, J., Calhoun, S., Bixler, E. O., Pejovic, S., Karataraki, M., Liao, D., et al.
38
39 (2010). Insomnia with objective short sleep duration is associated with deficits in
40
41 neuropsychological performance: a general population study. *Sleep*, 33, 459-465.
42
43
44 Fimm, B., Brand, T., & Spijkers, W. (2016). Time-of-day variation of visuo-spatial attention.
45
46 *British Journal of Psychology*, 107(2), 299-321.
47
48
49 Fossella, J., Sommer, T., Fan, J., Wu, Y. H., Swanson, J. M., Pfaff, D. W., et al. (2002).
50
51 Assessing the molecular genetics of attention networks. *BMC Neuroscience*, 3, 14.
52
53
54 Gamboz, N., Zamarian, S., & Cavallero, C. (2010). Age-related differences in the attention
55
56 network test (ANT). *Experimental Aging Research*, 36, 287-305.
57
58
59
60

- 1
2
3 Goel, N., Van Dongen, H., & Dinges, D. (2011). Circadian rhythms in sleepiness, alertness,
4 and performance. In M. H. Kryger, T. Roth, & W. C. Dement (Eds.), *Principles and*
5 *Practice of Sleep Medicine, Fifth edition* (pp. 445-455). Philadelphia: W. B. Saunders.
6
7
8
9
10 Hayes, A. F. (2013). *Introduction to mediation, moderation, and conditional process*
11 *analysis: A regression-based approach*. Guilford Press.
12
13
14
15 Hirshkowitz, M., Whiton, K., Albert, S. M., Alessi, C., Bruni, O., DonCarlos, L., et al.
16
17 (2015). National Sleep Foundation's sleep time duration recommendations:
18 methodology and results summary. *Sleep Health, 1*, 40-43.
19
20
21
22 Hoddes, E., Zarcone, V., & Dement, W. (1972). Development and use of Stanford Sleepiness
23 scale (SSS). In *Psychophysiology* (9) (p.150). New York: Cambridge University
24 Press.
25
26
27
28
29 Horne, J. A., & Östberg, O. (1976). A self-assessment questionnaire to determine
30 morningness-eveningness in human circadian rhythms. *International Journal of*
31 *Chronobiology, 4*, 97-110.
32
33
34
35
36 Jennings, J. M., Dagenbach, D., Engle, C. M., & Funke, L. J. (2007). Age-related changes
37 and the attention network task: An examination of alerting, orienting, and executive
38 function. *Aging, Neuropsychology, and Cognition, 14*, 353-369.
39
40
41
42
43 Jugovac, D., & Cavallero, C. (2012). Twenty-four hours of total sleep deprivation selectively
44 impairs attentional networks. *Experimental Psychology, 59*, 115-123.
45
46
47
48
49 Kalmbach, D. A., Arnedt, J. T., Song, P. X., Guille, C., & Sen, S. (2017). Sleep Disturbance
50 and Short Sleep as Risk Factors for Depression and Perceived Medical Errors in First-
51 Year Residents. *Sleep, 40*, <https://doi.org/10.1093/sleep/zsw1073>.
52
53
54
55
56
57
58
59
60 Katsanis, J., Iacono, W. G., & Harris, M. (1998). Development of oculomotor functioning in
preadolescence, adolescence, and adulthood. *Psychophysiology, 35*, 64-72.

- 1
2
3 Knutson, K. L., Rathouz, P. J., Yan, L. L., Liu, K., & Lauderdale, D. S. (2007). Intra-
4 individual daily and yearly variability in actigraphically recorded sleep measures: the
5 CARDIA study. *Sleep, 30*, 793.
6
7
8
9
10 Kronholm, E., Sallinen, M., Suutama, T., Sulkava, R., Era, P., & Partonen, T. (2009).
11 Self-reported sleep duration and cognitive functioning in the general population.
12 *Journal of Sleep Research, 18*, 436-446.
13
14
15
16
17 Lemola, S., Ledermann, T., & Friedman, E. M. (2013). Variability of sleep duration is related
18 to subjective sleep quality and subjective well-being: an actigraphy study. *PLoS One*,
19 8, e71292.
20
21
22
23
24 Lemola, S., Schwarz, B., & Siffert, A. (2012). Interparental conflict and early adolescents'
25 aggression: Is irregular sleep a vulnerability factor? *Journal of Adolescence, 35*, 97-
26 105.
27
28
29
30
31 Lev Ari, L., & Shulman, S. (2012). Pathways of sleep, affect, and stress constellations during
32 the first year of college: Transition difficulties of emerging adults. *Journal of Youth*
33 *Studies, 15*, 273-292.
34
35
36
37
38 Leys, C., Ley, C., Klein, O., Bernard, P., & Licata, L. (2013). Detecting outliers: Do not use
39 standard deviation around the mean, use absolute deviation around the
40 median. *Journal of Experimental Social Psychology, 49*(4), 764-766.
41
42
43
44
45 Lim, J., & Dinges, D. F. (2008). Sleep deprivation and vigilant attention. *Annals of the New*
46 *York Academy of Sciences, 1129*, 305-322.
47
48
49
50
51 Martella, D., Casagrande, M., & Lupianez, J. (2011). Alerting, orienting and executive
52 control: the effects of sleep deprivation on attentional networks. *Experimental Brain*
53 *Research, 210*, 81-89.
54
55
56
57
58
59
60

- 1
2
3 Matchcock, R. L., & Mordkoff, J. T. (2009). Chronotype and time-of-day influences on the
4 alerting, orienting, and executive components of attention. *Experimental Brain*
5 *Research, 192*, 189-198.
6
7
8
9
10 Matchcock, R. L., & Mordkoff, J. T. (2014). Effects of sleep stage and sleep episode length on
11 the alerting, orienting, and conflict components of attention. *Experimental brain*
12 *research, 232*, 811-820.
13
14
15
16
17 Mollicone, D. J., Van Dongen, H., Rogers, N. L., Banks, S., & Dinges, D. F. (2010). Time of
18 day effects on neurobehavioral performance during chronic sleep restriction. *Aviation,*
19 *Space, and Environmental Medicine, 81*, 735-744.
20
21
22
23
24 Moore, M., Kirchner, H. L., Drotar, D., Johnson, N., Rosen, C., & Redline, S. (2011).
25 Correlates of adolescent sleep time and variability in sleep time: the role of individual
26 and health related characteristics. *Sleep Medicine, 12*, 239-245.
27
28
29
30
31 Muto, V., Shaffii-le Bourdiec, A., Matarazzo, L., Foret, A., Mascetti, L., Jaspar, M., et al.
32 (2012). Influence of acute sleep loss on the neural correlates of alerting, orientating
33 and executive attention components. *Journal of Sleep Research, 21*, 648-658.
34
35
36
37
38 Ohayon, M. M., Carskadon, M. A., Guilleminault, C., & Vitiello, M. V. (2004). Meta-
39 analysis of quantitative sleep parameters from childhood to old age in healthy
40 individuals: developing normative sleep values across the human
41 lifespan. *Sleep, 27*(7), 1255-1273.
42
43
44
45
46
47 Patel, S. R., & Hu, F. B. (2008). Short sleep duration and weight gain: a systematic review.
48 *Obesity, 16*, 643-653.
49
50
51
52 Petersen, S. E., & Posner, M. I. (2012). The attention system of the human brain: 20 years
53 after. *Annual Review of Neuroscience, 35*, 73-89.
54
55
56
57
58
59
60

- 1
2
3 Roca, J., Fuentes, L. J., Marotta, A., Lopez-Ramon, M. F., Castro, C., Lupianez, J., et al.
4
5 (2012). The effects of sleep deprivation on the attentional functions and vigilance.
6
7 *Acta Psychologica, 140*, 164-176.
8
9
10 Sadeh, A., Sharkey, K. M., & Carskadon, M. A. (1994). Activity-Based Sleep—Wake
11
12 Identification: An Empirical Test of Methodological Issues. *Sleep, 17*, 201-207.
13
14 Sánchez-Ortuño, M. M., & Edinger, J. D. (2012). Internight sleep variability: its clinical
15
16 significance and responsiveness to treatment in primary and comorbid insomnia.
17
18 *Journal of Sleep Research, 21*, 527-534.
19
20
21 Spielman, A. J., Caruso, L., & Glovinsky, P. B. (1987). A behavioral perspective on insomnia
22
23 treatment. *Psychiatric Clinics of North America, 10*, 541-553.
24
25
26 Suh, S., Nowakowski, S., Bernert, R. A., Ong, J. C., Siebern, A. T., Dowdle, C. L., et al.
27
28 (2012). Clinical significance of night-to-night sleep variability in insomnia. *Sleep*
29
30 *Medicine, 13*, 469-475.
31
32
33 Tkachenko, O., & Dinges, D. F. (2018). Interindividual variability in neurobehavioral
34
35 response to sleep loss: A comprehensive review. *Neuroscience & Biobehavioral*
36
37 *Reviews*.
38
39
40 Trujillo, L. T., Kornguth, S., & Schnyer, D. M. (2009). An ERP examination of the different
41
42 effects of sleep deprivation on exogenously cued and endogenously cued attention.
43
44 *Sleep, 32*, 1285-1297.
45
46
47 Tworoger, S. S., Lee, S., Schernhammer, E. S., & Grodstein, F. (2006). The association of
48
49 self-reported sleep duration, difficulty sleeping, and snoring with cognitive function in
50
51 older women. *Alzheimer Disease & Associated Disorders, 20*, 41-48.
52
53
54 Vallières, A., Ivers, H., Bastien, C. H., Beaulieu-Bonneau, S., & Morin, C. M. (2005).
55
56 Variability and predictability in sleep patterns of chronic insomniacs. *Journal of Sleep*
57
58 *Research, 14*, 447-453.
59
60

- 1
2
3 Vallières, A., Ivers, H., Beaulieu-Bonneau, S., & Morin, C. M. (2011). Predictability of sleep
4
5 in patients with insomnia. *Sleep, 34*, 609-617.
6
7
8 Van Dongen, H. P. A., Maislin, G., Mullington, J. M., & Dinges, D. F. (2003). The
9
10 cumulative cost of additional wakefulness: Dose-response effects on neurobehavioral
11
12 functions and sleep physiology from chronic sleep restriction and total sleep
13
14 deprivation. *Sleep, 26*, 117-126.
15
16
17 Webb, W., & Agnew, H. (1975). Are we chronically sleep deprived? *Bulletin of the*
18
19 *Psychonomic Society, 6*, 47-48.
20
21
22 Whiting, W. L., & Murdock, K. K. (2016). Emerging adults' sleep patterns and attentional
23
24 capture: the pivotal role of consistency. *Cognitive Processing, 17*, 155-162.
25
26
27 Wittmann, M., Dinich, J., Mellow, M., & Roenneberg, T. (2006). Social jetlag: misalignment
28
29 of biological and social time. *Chronobiology International, 23*, 497-509.
30
31
32 Zils, E., Sprenger, A., Heide, W., Born, J., & Gais, S. (2005). Differential effects of sleep
33
34 deprivation on saccadic eye movements. *Sleep, 28*, 1109-1115.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table Captions

Table 1. Mean and SD RT (msec) for each experimental condition of the ANT (correct trials only and *4 absolute deviations from the median* excluded)

Table 2. Percentage of errors (SD) for each experimental condition of the ANT

Table 3. Correlations between key dependent variables, independent variables and potential covariates

Table 4. Multiple regression analyses of mean sleep duration, sleep duration variability and covariates predicting overall reaction time

Table 5. Multiple regression analyses of mean sleep duration, sleep duration variability and covariates predicting alerting

Table 6. Multiple regression analyses of mean sleep duration, sleep duration variability and covariates predicting orienting

Figure Captions

Figure 1. Attention Network Test procedure. (a) The four cue conditions; (b) The flanker types; and (c) An example of the procedure. Reprinted from Fan et al (2002) with permission granted by Prof. Posner.

Figure 2. Moderation models in PROCESS of a) association between sleep duration and overall reaction times moderated by sleep duration variability; b) association between sleep duration and alerting score moderated by sleep duration variability. Separate lines represent low (22.58 mins), medium (61 mins) and high (99.41 mins) sleep duration variability; and c) association between sleep duration variability and alerting score moderated by sleep duration

1
2
3 derived by median split (short sleepers <419.14 minutes; long sleepers ≥ 419.14 minutes).

4
5 Higher alerting scores indicate better efficiency of the alerting network.
6
7

8
9 Figure 3. Scatterplots of a) the association between sleep duration variability and alerting
10 score, showing the presence of an outlier experiencing high sleep duration variability; b) the
11 association between sleep duration the night before testing and alerting score, showing the
12 presence of an outlier experiencing long sleep duration the night before testing; and c) the
13 association between mean sleep duration and alerting score. Higher alerting scores indicate
14 presence of an outlier experiencing long sleep duration the night before testing; and c) the
15 association between mean sleep duration and alerting score. Higher alerting scores indicate
16 better efficiency of the alerting network.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1

Mean and SD RT (msec) for each experimental condition of the ANT (correct trials only and outliers 4 absolute deviations from the median excluded)

Flanker Types	Cue			
	No cue	Centre	Double	Spatial
Neutral	557.46 (47.40)	514.37 (52.41)	515.71 (56.85)	492.07 (51.01)
Congruent	609.17 (57.44)	572.16 (58.00)	569.46 (69.30)	547.95 (55.85)
Incongruent	716.57 (60.14)	710.17 (62.59)	703.26 (67.64)	664.84 (63.83)

Note. ANT = Attention Network Test; msec = milliseconds; SD = standard deviation

Table 2

Percentage of errors (SD) for each experimental condition of the ANT

Flanker Types	Cue			
	No cue	Centre	Double	Spatial
Neutral	3.93% (19.44%)	3.53% (18.46%)	3.38% (18.08%)	2.53% (15.72%)
Congruent	1.43% (11.88%)	0.85% (9.21%)	1.24% (11.09%)	1.81% (13.35%)
Incongruent	7.11% (25.71%)	8.02% (27.17%)	8.13% (27.34%)	6.75% (25.10%)

Note. ANT = Attention Network Test; SD = standard deviation

 Predictor Variables

6. Mean Sleep -.20 .16 -.07 .00 .00 /

Duration

7. Sleep -.14 .34** -.10 .10 .18 .17 /

Duration

Variability

 8. Sleep -.19 .30* -.10 .07 .18 .66** .47** /

Duration

Night

before

testing

 Potential Covariates

9. Age .38** -.26* .28* .18 -.03 -.28* -.20 -.21 /

10. Time ANT	-.25	.40**	.19	.16	-.10	.16	.27*	.16	-.02	/			
tested													
11. Sleepiness	-.19	.19	.08	.05	-.03	.11	-.03	.11	-.12	.13	/		
12. Sleep Quality	-.10	.14	.12	.13	-.07	-.13	-.07	-.04	.15	.23	.35**	/	
13. Chronotype	.23	-.04	.36**	.23	-.11	-.13	-.19	-.20	.28*	.11	-.03	.29*	/

Note. * $p < .05$; ** $p < .01$ (Bonferroni corrected). ANT = Attention Network Test. All variables are raw scores.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Table 4

Multiple regression analyses of mean sleep duration, sleep duration variability and covariates predicting overall reaction time

Covariates/ Predictors	<u>Hierarchical Regression</u>				<u>Regression Coefficients of Predictors/Covariates (Simultaneous Entry)</u>					
	ΔR^2	R^2	Adj R^2	$p \Delta F$	B	SE B	β	T	p	
Age	.14	.14	.13	.00	2.09	.77	.35	2.70	.01	
Sleep Duration	.01	.15	.12	.44	-.08	.17	-.06	-.45	.65	
Sleep Duration Variability	.00	.15	.11	.65	.20	.22	.15	.94	.35	
Sleep Duration X Sleep Duration Variability	.07	.22	.16	.04	-.01	.00	-.34	-2.13	.04	

Note. Analyses were performed with centred predictors and covariates.

Table 5

Multiple regression analyses of mean sleep duration, sleep duration variability and covariates predicting alerting

Covariates/ Predictors	<u>Hierarchical Regression</u>				<u>Regression Coefficients of Predictors/Covariates (Simultaneous Entry)</u>				
	<u>ΔR^2</u>	<u>R^2</u>	<u>Adj R^2</u>	<u>$p \Delta F$</u>	<u>B</u>	<u>SE B</u>	<u>β</u>	<u>T</u>	<u>p</u>
Age	.07	.07	.05	.05	-.53	.28	-.22	-1.87	.07
Time of Testing	.15	.22	.19	.00	2.06	.70	.35	2.96	.01
Sleep Duration	.00	.22	.18	.82	-.02	.06	-.03	-.28	.78
Sleep Duration	.04	.26	.20	.13	-.04	.08	-.07	-.47	.64
Variability									
Sleep Duration X Sleep	.11	.37	.30	.01	.00	.00	.43	2.94	.01
Duration Variability									

Note. Analyses were performed with centred predictors and covariates.

Table 6

Multiple regression analyses of mean sleep duration, sleep duration variability and covariates predicting orienting

Covariates/ Predictors	<u>Hierarchical Regression</u>				<u>Regression Coefficients of</u>					
	<u>Indices</u>					<u>Predictors/Covariates (Simultaneous</u>				
	ΔR^2	R^2	Adj R^2	p	B	SE B	β	T	p	
Age	.08	.08	.06	.03	.46	.31	.21	1.50	.14	
Chronotype	.08	.16	.13	.02	.56	.25	.31	2.28	.03	
Sleep Duration	.00	.16	.12	.81	.02	.07	.05	.33	.74	
Sleep Duration Variability	.00	.16	.10	.93	.03	.08	.06	.34	.74	
Sleep Duration X Sleep Duration Variability	.01	.17	.10	.51	-.00	.00	-.11	-.66	.51	

Note. Analyses were performed with centred predictors and covariates.




Figure 1. Attention Network Test procedure. (a) The four cue conditions; (b) The flanker types; and (c) An example of the procedure. Reprinted from Fan et al (2002) with permission granted by Prof. Posner.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2a. Moderation model in PROCESS of association between sleep duration and overall reaction times moderated by sleep duration variability

224x131mm (96 x 96 DPI)




Figure 2b. Moderation model in PROCESS of association between sleep duration and alerting score moderated by sleep duration variability. Separate lines represent low (22.58 mins), medium (61 mins) and high (99.41 mins) sleep duration variability. Higher alerting scores indicate better efficiency of the alerting network.

219x131mm (96 x 96 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2c. Moderation model in PROCESS of association between sleep duration variability and alerting score moderated by sleep duration derived by median split (short sleepers <419.14 minutes; long sleepers ≥419.14 minutes). Higher alerting scores indicate better efficiency of the alerting network.

202x133mm (96 x 96 DPI)




Figure 3. Scatterplots of a) the association between sleep duration variability and alerting score, showing the presence of an outlier experiencing high sleep duration variability; b) the association between sleep duration the night before testing and alerting score, showing the presence of an outlier experiencing long sleep duration the night before testing; and c) the association between mean sleep duration and alerting score. Higher alerting scores indicate better efficiency of the alerting network.

466x339mm (96 x 96 DPI)