

Northumbria Research Link

Citation: Lofrano-Prado, Mara C., do Prado, Wagner Luiz, Botero, Joao Paulo, Cardel, Michelle L., Farah, Breno Q., Oliveira, Max D., Cucato, Gabriel, Correia, Marilia A. and Ritti-Dias, Raphael M. (2021) The same storm but not the same boat: Effects of COVID -19 stay-at-home order on mental health in individuals with overweight. *Clinical Obesity*, 11 (1). e12425. ISSN 1758-8103

Published by: Wiley-Blackwell

URL: <https://doi.org/10.1111/cob.12425> <<https://doi.org/10.1111/cob.12425>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/45025/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

- What is already known about this subject?

- Overweight has been associated with increased risks of biological health-related complications by COVID-19.
- Women with obesity/overweight have an increased risk to develop mental struggles, such as depression and anxiety.

- What this study adds?

- Women with overweight are more likely to report higher levels of anxiety, depression, and sadness, and lower self-esteem relative to women with normal weight.
- The present data provide the opportunity to make meaningful change at the policy level, and demonstrate need for increasing mental health resources during these difficult times.

ABSTRACT

Objective: To describe the effects of stay-at-home orders and social distancing during the COVID-19 outbreak on mental health and to compare these outcomes between individuals with normal weight and overweight.

Methods: This cross-sectional study included 1857 Brazilian adults, who were invited through social media to answer an online questionnaire from May 05th and 17th 2020. The instrument included questions related to health behavior, mental health (anxiety, depression, self-esteem, sadness and stress) and overall health. Overweight was defined as body mass index (BMI) ≥ 25 Kg/m². Multiple logistic regression was conducted to identify whether overweight is associated with mental health variables.

Results: Women reported increased anxiety (36.5% vs. 22.2%, $p < 0.01$), depression (16.2% vs. 8.8%, $p < 0.01$), low self-esteem (19.8% vs. 10.6%, $p < 0.01$), sadness (17.7% vs. 10.2%, $p < 0.01$), and stress (29.5% vs. 19.3%, $p < 0.01$) relative to men. Women with overweight are more likely to report higher feeling of anxiety (OR 1.62, CI95% 1.22-2.14), depression (OR 1.79, CI95% 1.25-2.55), low self-esteem (OR 1.82, CI95% 1.28-2.58) and sadness (OR 1.51, CI95% 1.08-2.10), adjusted for age, social isolation days, educational level, chronic diseases, smoke, alcohol intake and physical activity.

Conclusion: Women, specially those with overweight are more vulnerable to the deleterious effects of stay-at-home orders on mental health during the COVID-19 pandemic.

INTRODUCTION

The 2019 new coronavirus disease (COVID-19) pandemic is the largest outbreak in decades.(1) With the lack of specific pharmacological treatment and vaccine options to prevent the spread of COVID-19, both “stay-at-home” measures and social isolation has been proposed by the World Health Organization in order to flatten the curve of COVID-19 infection and reduce the spread of the virus. While studies suggest a mitigating effect of social distancing on the disease spread,(2, 3) boredom, frustration, and a sense of isolation may lead to unintended consequences on mental health(4, 5)

Sex and weight status have been associated with adverse mental health outcomes.(6) Specifically, women and individuals with overweight have an increased risk to develop mental struggles, such as depression and anxiety.(7) Therefore, it is plausible to speculate that these groups are more likely to suffer deleterious effects of living through a pandemic and social isolation on mental health. In this study, we describe the effects of stay-at-home orders and social distance during the COVID-19 outbreak on mental health and to compare these outcomes between individuals with normal weight and overweight. We posit that women and individuals with overweight will be negatively affected by stay-at-home orders and social isolation resulting from COVID-19.

MATERIALS AND METHODS

Sample and Ethics

This survey was conducted in Brazil between May 05th and 17th 2020. Participants were invited through social media ([Facebook](#), [Twitter](#), [Instagram](#), and [WhatsApp](#)) to answer an online questionnaire. This study was approved by the Universidade Nove de Julho’ Ethics Committee (CAAE #30890220.4.0000.5511). Consent was provided and participants did not identify themselves in the survey. All procedures follow the

Declaration of Helsinki. Inclusion criteria was limited to adults ages 18 or older and completing all responses within the survey.

Procedures

After ethics approval, a questionnaire (in Portuguese) in Google Forms was presented to participants with 70 questions divided into 7 domains: (a) personal information; (b) COVID-19 personal care; (c) physical activity; (d) eating behavior; (e) health risk habits; (f) mental health; (g) overall health. For the purpose of the present study relevant questions in the following domains were included: (a) personal information, (b) Covid-19 personal care, (c) physical activity, (e) health risk habits, (f) mental health, and (g) overall health domains.

Domains

Mental health: This domain is composed of 6-self-reported items to identify individuals' feelings related to the COVID-19 pandemic. The following questions were used: (1) Due to COVID-19, are you feeling more anxious than usual?, (2) Due to COVID-19, are you feeling more sad than usual?, (3) Due to COVID-19, are you feeling more stressed than usual?, (4) Due to COVID-19, are you feeling more depressed than usual?, and (5) Due to COVID-19, are you feeling lower self-esteem than usual? Responses included a Likert scale with responses ranging from 'No, A little, Sometimes, Very often, or Always'.

Personal information: From this domain three self-reported information was collected including: (1) Sex (possible answers: 'Woman or Man'), (2) Date of birth (DD/MM/YYYY), and (3) Educational level (open question).

Overall health: This domain assesses the presence of diagnosed diseases. From the list of diseases, the participant was asked to mark all that apply. (possible answers: Hypertension, diabetes, high cholesterol, high triglycerides, depression, arthritis/osteoarthritis/rheumatism, asthma, cardiopathy, or other). In this domain, we also asked “What is your weight (in kilograms)?” and “What is your height (in centimeters)?”

COVID-19 personal care: The question “How long have you (in days) experienced social isolation?” was included as an open question.

Physical activity: The questions included: (1) How many times are you exercising a week? (possible answers: ‘None, 1, 2, 3, 4, 5, 6, 7’), (2) For how long are you exercising daily? (possible answers: ‘None, less than 30 min, between 30 to 60 min, more than 60 min’), and (3) What is the intensity of the physical activity? Low: Standing work, light household chores; Medium: Walking and other moderate activities; High: Jogging, running or other intense activities (possible answers: ‘Low, Medium / Moderate, High, I am not exercising’).

Health risk behavior: This domain aimed identify health-related behaviors. Questions included: (1) Do you smoke? (possible answers were: ‘yes or no’), and (2) Do you drink alcohol? (possible answers were: ‘yes or no’);

Overweight definition: Height and weight were self-reported (overall health domain) and used for standard BMI calculation (kg/m^2). Following the WHO recommendations,(8) overweight was defined as $\text{BMI} \geq 25 \text{ Kg/m}^2$.

Statistical Analysis

Frequency and mean (95% confidence interval) were used for descriptive analyses. Multiple logistic regression was conducted to identify the extent to which weight status is associated with mental health variables during COVID-19 (changes in of

anxiety, depression, self-esteem, sadness and stress). Multiple logistic regression was adjusted for confounders (age, social isolation days, educational level, **pre-existing depression**, chronic disease status, smoker, alcoholic beverages and physical activity level). In multivariate models we also tested for interactions of sex for the studied variables. The Hosmer-Lemeshow test was used to assess the model's goodness-of-fit. The significance level was set at $p < 0.05$ for all analyses. All statistical analyses were made in the Statistical Package for the Social Sciences software – SPSS/PASW version 20 (IBM Corp, New York, USA).

RESULTS

Participants include 1895 adults, but 46 did not answer all questions and were not included in the present analyses. The final sample was composed of 1854 adults living in Brazil. Table 1 includes participant characteristics and psychological variables of subjects included in this study according to sex and weight status. **Overweight women and men were significantly older ($p < 0.001$), higher body mass index ($p < 0.001$), had spent less days in social isolation ($p = 0.026$), were more likely to chronic diseases ($p < 0.001$) compared to normal weight. Overweight women were more likely be physically active and to report increases in anxiety, depression, low self-esteem, sadness, and stress related to COVID-19 than normal weight women ($p < 0.001$). Consume alcohol was higher in overweight men than normal ones ($p < 0.001$).**

Please insert table 1

There was an interaction between sex and overweight status for mental health variables ($P < 0.001$). Among women, overweight was positively associated with

increased anxiety, depression, and sadness, and lower self-esteem ($P < 0.050$ for all). Stress was not significantly associated with weight status. Among men, overweight was not significantly associated with mental health variables. For all regressions, the Hosmer-Lemeshow test was not significant.

*** Please insert Table 2***

DISCUSSION

In this study, we examine the extent to which stay-at-home order and social isolation during the COVID-19 pandemic affected self-reported mental health changes among men and women with varying weight status. We observed that during the COVID-19 pandemic (1) women reported worse mental health compared to men, (2) women with overweight are more likely to report higher levels of anxiety, depression, and sadness, and lower self-esteem relative to women with normal weight, and 3) weight status is not associated with mental health among men.

A systematic review showed the overall prevalence of depressive symptoms among Brazilian adults prior to the initiation of COVID-19 was 14% (95%CI 13-16; $I^2 = 99.5\%$). The prevalence of 1-year major depressive disorder among Brazilian adults was 8% (95%CI 7-10; $I^2 = 86.7\%$), and the prevalence was higher among Brazilian woman in both depressive symptoms and major depressive disorder (9). A 2019 study among adults in Sao Paulo ($n=5,037$) found 19.9% presented with an anxiety disorder, and anxiety disorders were more frequent in women(10).

From a biological perspective, studies have demonstrated that men are facing more health complications and hospitalization,(11) as well as higher risk to death related to the COVID-19 compared to women.(12) However, our data suggests that women are

more vulnerable than men to being adversely affected in mental health outcomes resulting from COVID-19. Our results are consistent with findings from a survey conducted with 52,730 individuals in China during the COVID-19 pandemic, which found that women had significantly higher psychological distress than their male counterparts did. One potential explanation is that women may suffer more due the dramatic changes in lifestyle habits during the pandemic, such as isolation, disproportionate burden on home and childcare demands, resulting in impaired mental health outcomes.(13, 14)

Overweight status has been associated with increased risks of biological health-related complications by COVID 19.(11, 15) Noteworthy, individuals with obesity are more likely to present with adverse mental health problems.(16) **A recent online survey conducted with a local clinical sample of adult patients with obesity (n=123; 87% female) demonstrated COVID-19 stay-at-home-orders adversely affected several health-related outcomes, including anxiety (72.8%) and depression (83.6%) (17).** Our results add new data to the current literature showing that women with overweight present 62% more chance to present feelings of anxiety, 79% more chance to present feeling of depression, 82% more chance to present feelings of sadness and 51% more chance to present feelings of decreased self-esteem than their normal weight peers as a result of COVID-19.

The quarantine and social isolation have a differential negative impact on mental health by sex. Though all participants were in the same storm of COVID-19, women, particularly those with overweight, appear to be in a different boat, and are more susceptible to mental health suffering during the outbreak. Clinicians may better serve by being aware of potential vulnerabilities patients are facing in order to provide the adequate support to minimize the side-effects of stay-at-home order and social isolation due to the COVID-19 pandemic.

Remote strategies to tackle psychological consequences during COVID-19 pandemic, especially among women with overweight could be useful. Evidence demonstrates telemedicine including psychosocial interventions, particularly cognitive behavioural therapy via telephone or videoconference, can improve quality of life, depression, and anxiety in clinical populations.(1, 18) Even with insufficient data for the effectiveness of eHealth interventions for individuals with overweight, this can be considered as an alternative strategy during the COVID-19 pandemic.(19) Also, we can suggest that family and friends must be aware of these issues and try to support their beloved ones.

It is important to note that our use of self-reported height and weight can lead to an underestimation of BMI. Additionally, we assessed self-reported changes in depression, anxiety, self-esteem and sadness, but this data is not diagnostic, nor can it be related to diseases or symptoms of mental health disorders. The use of non-validated questionnaires to assess the psychological outcomes, the lack of information about mental health status for participants prior to stay-at-home orders, and the lack of data related to household size and socio-economic status are limitations of the present study. Moreover, the population recruited for this study is not a representative sample, thus results may not be widely generalizable.

Our data suggest that women are more susceptible to the deleterious effects of stay-at-home orders and social isolation and women with overweight are at a greater risk to face mental health problems than normal weight peers.

CONFLICTS OF INTEREST STATEMENT: The authors declared no conflict of interest.

ACKNOWLEDGEMENTS: The authors thank to the volunteers and the health professionals and all the essential workers who are fighting for us during the covid-19 pandemic.

AUTHOR CONTRIBUTIONS: MCLP and WLP designed the study. WLP, RMRD, MAC, GC and JPB contributed to the design and implementation of the research. BQF and MDO performed the analysis and designed the figures. MCLP, WLP and MC analysed the data. MCLP and WLP wrote the paper with input from all authors. All authors contributed to manuscript revision, read and approved the submitted version.

REFERENCES

1. Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS, *et al.* Immediate Psychological Responses and Associated Factors during the Initial Stage of the 2019 Coronavirus Disease (COVID-19) Epidemic among the General Population in China. *International journal of environmental research and public health* 2020;**17**.
2. Koo JR, Cook AR, Park M, Sun Y, Sun H, Lim JT, *et al.* Interventions to mitigate early spread of SARS-CoV-2 in Singapore: a modelling study. *The Lancet Infectious diseases* 2020.
3. Luo H, Lie Y, Prinzen FW. Surveillance of COVID-19 in the General Population Using an Online Questionnaire: Report From 18,161 Respondents in China. *JMIR public health and surveillance* 2020;**6**: e18576.
4. Rubin GJ, Wessely S. The psychological effects of quarantining a city. *BMJ (Clinical research ed)* 2020;**368**: m313.
5. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, *et al.* The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet (London, England)* 2020;**395**: 912-920.
6. Pan L, Li X, Feng Y, Hong L. Psychological assessment of children and adolescents with obesity. *The Journal of international medical research* 2018;**46**: 89-97.
7. Lazarevich I, Irigoyen Camacho ME, Velázquez-Alva MDC, Zepeda Zepeda M. Relationship among obesity, depression, and emotional eating in young adults. *Appetite* 2016;**107**: 639-644.
8. WHO. Obesity and overweight. <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight> 20 may 2020.
9. Silva M, Galvão T, Martins S, Pereira MJRbdp. Prevalence of depression morbidity among Brazilian adults: a systematic review and meta-analysis. 2014;**36 3**: 262-270.
10. Andrade LH, Wang YP, Andreoni S, Silveira CM, Alexandrino-Silva C, Siu ER, *et al.* Mental disorders in megacities: findings from the São Paulo megacity mental health survey, Brazil. *PloS one* 2012;**7**: e31879.
11. Richardson S, Hirsch JS, Narasimhan M, Crawford JM, McGinn T, Davidson KW, *et al.* Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *Jama* 2020.

12. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, *et al.* Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet (London, England)* 2020;**395**: 1054-1062.
13. Cardel MI, Dhurandhar E, Yarar-Fisher C, Foster M, Hidalgo B, McClure LA, *et al.* Turning Chutes into Ladders for Women Faculty: A Review and Roadmap for Equity in Academia. *Journal of women's health (2002)* 2020;**29**: 721-733.
14. Jolly S, Griffith KA, DeCastro R, Stewart A, Ubel P, Jagsi R. Gender differences in time spent on parenting and domestic responsibilities by high-achieving young physician-researchers. *Annals of internal medicine* 2014;**160**: 344-353.
15. Petrakis D, Margină D, Tsarouhas K, Tekos F, Stan M, Nikitovic D, *et al.* Obesity - a risk factor for increased COVID-19 prevalence, severity and lethality (Review). *Molecular medicine reports* 2020;**22**: 9-19.
16. Rajan TM, Menon V. Psychiatric disorders and obesity: A review of association studies. *Journal of postgraduate medicine* 2017;**63**: 182-190.
17. Almandoz JP, Xie L, Schellinger JN, Mathew MS, Gazda C, Ofori A, *et al.* Impact of COVID-19 stay-at-home orders on weight-related behaviours among patients with obesity. *Clinical obesity* 2020;**10**: e12386.
18. David LA, Sijercic I, Cassin SE. Preoperative and post-operative psychosocial interventions for bariatric surgery patients: A systematic review. *Obesity reviews : an official journal of the International Association for the Study of Obesity* 2020;**21**: e12926.
19. Patel ML, Wakayama LN, Bass MB, Breland JY. Motivational interviewing in eHealth and telehealth interventions for weight loss: A systematic review. *Preventive medicine* 2019;**126**: 105738.

Table 1. Participant Characteristics and Psychological Variables Stratified by Sex.

Variables	Men (n=769)		p	Women (n=1085)		P
	Normal	Overweight		Normal	Overweight	
Age (years old)	34.8±13.3	42.0±13.2	<0.001	35.4±11.8	40.1±13.0	<0.001
Social isolation (days)	41.9±16.3	43.3±14.3	0.243	44.7±13.5	44.1±15.7	0.500
Body index mass (kg/m ²)	23.0±1.6	29.3±3.7	<0.001	22.0±1.9	29.1±4.5	<0.001
Pre-existing Chronic disease (% Yes)	20.2	35.5	<0.001	25.6	43.6	<0.001
Pre-existing Depression (% Yes)	2.3	3.0	0.590	5.4	8.3	0.053
Pre-existing Diabetes (% Yes)	1.1	1.8	0.504	0.8	1.9	0.152
Smoker (% Yes)	8.0	6.9	0.574	3.5	5.1	0.203
Alcohol intake (% Yes)	65.4	77.4	<0.001	60.7	60.5	0.943
Physically inactive (% Yes)	64.9	69.0	0.244	69.3	81.2	<0.001
Weight status						
Overweight (% Yes)	-	67.3	-	-	69.8	
Class 1 Obesity (% Yes)	-	25.7	-	-	23.2	-
Class 2 Obesity (% Yes)	-	5.4	-	-	4.4	-
Class 3 Obesity (% Yes)	-	1.6	-	-	2.6	-
Mental Health Variables in Relations to COVID-19						
Increased Anxiety (% Yes)	21.4	22.9	0.635	33.7	40.3	0.029
Increased Depression (% Yes)	9.2	8.9	0.902	14.0	19.7	0.013
Lower Self-esteem (% Yes)	9.9	10.5	0.819	14.7	21.8	0.003
Increased Sadness (% Yes)	11.8	10.3	0.505	17.5	22.7	0.035
Increased Stress (% Yes)	19.5	19.5	0.984	29.0	30.2	0.690

Data presented as mean ± standard deviation or frequency

Table 2. Crude and adjusted odds ratios (OR) and confidence interval (CI 95%) for the association between overweight and mental health variables in men and women.

		Men		Women	
		Crude OR (CI95%)	Adjusted OR (CI95%)	Crude OR (CI95%)	Adjusted OR (CI95%)
Increased Anxiety (% Yes)	Normal weight	1	1	1	1
	Overweight	1.09 (0.76 – 1.57)	1.29 (0.87 – 1.93)	1.32 (1.03 – 1.70)	1.63 (1.22 – 2.16)
Increased Depression (% Yes)	Normal weight	1	1	1	1
	Overweight	0.97 (0.58 – 1.63)	1.18 (0.67 – 2.10)	1.51 (1.09 – 2.08)	1.88 (1.30 – 2.70)
Lower self-esteem (% Yes)	Normal weight	1	1	1	1
	Overweight	1.06 (0.65 – 1.74)	1.51 (0.88 – 2.60)	1.61 (1.18 – 2.21)	1.90 (1.34 – 2.69)
Increased Sadness (% Yes)	Normal weight	1	1	1	1
	Overweight	0.85 (0.53 – 1.37)	1.13 (0.68 – 1.90)	1.39 (1.02 – 1.87)	1.57 (1.12 – 2.10)
Increased Stress (% Yes)	Normal weight	1	1	1	1
	Overweight	1.00 (0.69 – 1.46)	1.30 (0.86 – 1.97)	1.06 (0.81 – 1.38)	1.20 (0.90 – 1.61)

OR – odds-ratio; CI95% - 95% confidence intervals. Adjusted for age, social isolation days, educational level, chronic diseases, smoker, alcoholic intake, pre-existing depression, and physical activity level.