

Northumbria Research Link

Citation: Angelova, Maia, Hertz, Anaëlle and Hussin, Veronique (2012) Squeezed coherent states and the one-dimensional Morse quantum system. *Journal of Physics A: Mathematical and Theoretical*, 45 (24). p. 244007. ISSN 1751-8113

Published by: IOP Publishing

URL: <http://dx.doi.org/10.1088/1751-8113/45/24/244007>
<<http://dx.doi.org/10.1088/1751-8113/45/24/244007>>

This version was downloaded from Northumbria Research Link:
<https://nrl.northumbria.ac.uk/id/eprint/6372/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

Squeezed coherent states and the one-dimensional Morse quantum system

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2012 J. Phys. A: Math. Theor. 45 244007

(<http://iopscience.iop.org/1751-8121/45/24/244007>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 88.104.157.164

The article was downloaded on 21/06/2012 at 00:27

Please note that [terms and conditions apply](#).

Squeezed coherent states and the one-dimensional Morse quantum system

M Angelova¹, A Hertz² and V Hussin³

¹ Mathematical Modelling Lab, School of Computing, Engineering and Information Sciences, Northumbria University, Newcastle NE2 1XE, UK

² Département de Physique, Université de Montréal, Montréal, Québec H3C 3J7, Canada

³ Département de Mathématiques et de Statistique, Université de Montréal, Montréal, Québec H3C 3J7, Canada

E-mail: maia.angelova@northumbria.ac.uk, anaelle.hertz@umontreal.ca and veronique.hussin@umontreal.ca

Received 3 November 2011, in final form 21 February 2012

Published 30 May 2012

Online at stacks.iop.org/JPhysA/45/244007

Abstract

The Morse potential one-dimensional quantum system is a realistic model for studying vibrations of atoms in a diatomic molecule. This system is very close to the harmonic oscillator one. We thus propose a construction of squeezed coherent states similar to the one of the harmonic oscillator using ladder operators. The properties of these states are analysed with respect to the localization in position, minimal Heisenberg uncertainty relation, the statistical properties and illustrated with examples using the finite number of states in a well-known diatomic molecule.

This article is part of a special issue of *Journal of Physics A: Mathematical and Theoretical* devoted to ‘Coherent states: mathematical and physical aspects’.

PACS numbers: 03.65.Aa, 03.65.–w

(Some figures may appear in colour only in the online journal)

1. Introduction

Coherent and squeezed states are known to be very important in many fields of physics. Coherent states (CS) were discovered in 1926 by Schrödinger [1], while squeezed states were introduced by Kennard in 1927 [2]. However, these works were, in the main, ignored or forgotten until the 1960s, when these states became very popular and received a lot of attention from both fields, mathematics and physics. Among many important papers, let us mention the works of Glauber [3], Klauder [4, 5] and Nieto [6]. In the particular field of quantum optics, the books of Walls and Milburn [7], Gazeau [8] and Rand [9] make very good reading and also consider applications. The study of squeezed states for systems admitting an infinite discrete spectrum, obtained as a generalization of CS, has recently been the centre of much attention (see, for example, [10–14]).

In modern developments, CS are standardly defined in three equivalent ways: displacement operator method, ladder (annihilation) operator method and minimum uncertainty method (for a review, see for example [6]). Initially defined for the case of the harmonic oscillator, CS have been generalized for other systems. We can use, for example, the definition of Klauder [15] saying that they are obtained as the following superposition of energy eigenstates $\{|\psi_n\rangle, n \in \mathbb{N}\}$:

$$\psi(z) = \frac{1}{\sqrt{\mathcal{N}(|z|^2)}} \sum_{n \in I} \frac{z^n}{\sqrt{\rho(n)}} |\psi_n\rangle. \quad (1)$$

The sum is taken over all the discrete values of n and the set I is usually infinite. The parameter z is a complex variable in general, \mathcal{N} is a normalization factor and $\{\rho(n), n \in \mathbb{N}\}$ is a set of strictly positive parameters, usually depending on the energy of the system under consideration. The last quantities correspond to a moment problem (see [15] for details).

For a quantum system which admits an infinite discrete spectrum $\{|\psi_n\rangle, n = 0, 1, \dots\}$, and ladder operators A^- and A^+ acting on the energy eigenstates as

$$A^- |\psi_n\rangle = \sqrt{k(n)} |\psi_{n-1}\rangle, \quad A^+ |\psi_n\rangle = \sqrt{k(n+1)} |\psi_{n+1}\rangle, \quad (2)$$

these CS are defined as eigenstates of A^- . We thus obtain [15]

$$\rho(n) = \prod_{i=1}^n k(i), \quad \rho(0) = 1. \quad (3)$$

(Note that the quantity $k(i)$ is not unique and can be chosen to impose additional constraints on the ladder operators. In particular, for the harmonic oscillator, we have $k(i) = i$,

$$[A^-, A^+] = \mathbb{I}, \quad H_{\text{ho}} = \hbar\omega(A^+ A^- + \frac{1}{2}), \quad (4)$$

and expression (1) gives the usual CS.)

Now, for a quantum system which admits a finite discrete spectrum like the one which involves the Morse potential, various constructions of CS have been adapted [16–19]. In a recent paper [20], we have used ladder operators [21, 22] to construct different types of CS of the Morse potential and have compared them with the so-called Gaussian CS [23]. In particular, such a construction has been inspired by the approach mentioned above (see formula (1)) but where the set I of values of n is now finite. The CS are not exactly eigenstates of the annihilation operator A^- but we have shown [20] that, in practice, the last terms on the right-hand side of the sum in (1) do not contribute significantly. In some approaches (see, for example, [24]), these states are called pseudo-CS.

To our knowledge, squeezed coherent states (SCS) for the Morse potential have not been constructed. The aim of this paper is thus to show that such a construction can be closely related to the one for infinite spectrum systems. In fact, these states would be almost eigenstates of a linear combination of the ladder operators. Moreover, we demonstrate using numerical evaluations that one set of these states, the energy-like, behave properly as squeezed states.

In section 2, we give a review of relevant results on SCS and minimal uncertainty relations for a quantum system with an infinite spectrum. In section 3, starting with the definition of the Morse model and its ladder operators, we define the corresponding SCS. We thus obtain two types of states called oscillator-like and energy-like. In section 4, we show that the energy-like states have stable trajectories in the phase space and minimize the uncertainty relation. Due to the complexity of the expressions of those states, numerical calculations are used. We conclude the paper in section 5.

2. SCS for a quantum system with an infinite spectrum

As in the case of the harmonic oscillator, general SCS [6], for a quantum system with an infinite discrete energy spectrum, may be constructed as solutions of the eigenvalue equation:

$$(A^- + \gamma A^+) \psi(z, \gamma) = z \psi(z, \gamma), \quad z, \gamma \in \mathbb{C}. \quad (5)$$

The mixing of A^- and A^+ is said to be controlled by a squeezing parameter γ and z is called the coherent parameter. The CS are special solutions when $\gamma = 0$. Conditions on γ must be imposed for the states to be normalizable.

SCS based on $su(2)$ or $su(1, 1)$ algebras [12, 13] and also direct sums of these algebras with the algebra $h(2)$ [14] have been constructed using group theoretical methods. This involves, in particular, the operator displacement D and squeezing S similar to the ones of the harmonic oscillator. In fact, for $su(2)$ or $su(1, 1)$ algebras, $k(n)$ is a quadratic function of n .

More generally, equation (5) may be solved by using a direct expansion of $\psi(z, \gamma)$ in the form

$$\psi(z, \gamma) = \frac{1}{\sqrt{\mathcal{N}_g(z, \gamma)}} \sum_{n=0}^{\infty} \frac{Z(z, \gamma, n)}{\sqrt{\rho(n)}} |\psi_n\rangle, \quad (6)$$

with

$$\mathcal{N}_g(z, \gamma) = \sum_{n=0}^{\infty} \frac{|Z(z, \gamma, n)|^2}{\rho(n)}, \quad (7)$$

where $\rho(n)$ is given by (3). Indeed, for the case $\gamma \neq 0$, inserting (6) into (5), we obtain a three-term recurrence relation,

$$Z(z, \gamma, n+1) - z Z(z, \gamma, n) + \gamma k(n) Z(z, \gamma, n-1) = 0, \quad n = 1, 2, \dots, \quad (8)$$

and without restriction, we take $Z(z, \gamma, 0) = 1$ and thus $Z(z, \gamma, 1) = z$.

For the harmonic oscillator, $\rho(n) = n!$ and we obtain explicitly [14, 25]

$$Z_{ho}(z, \gamma, n) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \frac{n!}{i!(n-2i)!} \left(-\frac{\gamma}{2}\right)^i z^{(n-2i)} = \left(\frac{\gamma}{2}\right)^{\frac{n}{2}} \mathcal{H}\left(n, \frac{z}{\sqrt{2\gamma}}\right). \quad (9)$$

It is well known that $|\gamma| < 1$ for the states to be normalizable in this case. In (9), we see that the Hermite polynomials $\mathcal{H}(n, w)$ take values on \mathbb{C} . These polynomials have interesting properties in terms of orthogonality, measure and resolution of the identity [26].

For the harmonic oscillator, these states minimize the Schrödinger–Robertson uncertainty relation [27] which becomes the usual Heisenberg uncertainty relation for γ real. Indeed, we obtain

$$(\Delta \hat{x})^2 = \frac{1}{1+\gamma} - \frac{1}{2}, \quad (\Delta \hat{p})^2 = \frac{1}{1-\gamma} - \frac{1}{2}, \quad (10)$$

and the uncertainty becomes

$$\Delta(z, \gamma) = (\Delta \hat{x})^2 (\Delta \hat{p})^2 = \frac{1}{4}. \quad (11)$$

We see that this implies the reduction of the ‘quantum noise’ on one of the observables while increasing it on the other. In the following, we will treat the case when the quantum noise is reduced on the observable x because we want a good localization in the position.

3. The Morse potential and different types of SCS

The Morse potential quantum system is a realistic model for studying vibrations of atoms in a diatomic molecule. Since this system is very close to the harmonic oscillator, the SCS will be constructed following the procedure given for the harmonic oscillator, but we will deal with a finite number of eigenstates.

3.1. The model

The one-dimensional Morse model is given by the energy eigenvalue equation (see, for example, [21])

$$\hat{H} \psi(x) = \left(\frac{\hat{p}^2}{2m_r} + V_M(x) \right) \psi(x) = E \psi(x), \quad (12)$$

where m_r is the reduced mass of the oscillating system composed of two atoms of masses m_1 and m_2 , i.e. $\frac{1}{m_r} = \frac{1}{m_1} + \frac{1}{m_2}$. The potential is $V_M(x) = V_0(e^{-2\beta x} - 2e^{-\beta x})$, where the space variable x represents the displacement of the two atoms from their equilibrium positions, V_0 is a scaling energy constant representing the depth of the potential well at equilibrium $x = 0$ and β is the parameter of the model (related to the characteristics of the well, such as its depth and width).

The finite discrete spectrum is known as

$$E_n = -\frac{\hbar^2}{2m_r} \beta^2 \epsilon_n^2, \quad (13)$$

where

$$\epsilon_n = \frac{\nu - 1}{2} - n = p - n, \quad \nu = \sqrt{\frac{8m_r V_0}{\hbar^2 \beta^2}}, \quad (14)$$

and $\{n = 0, 1, 2, \dots, [p]\}$, with $[p]$ the integer part of $p = \frac{\nu-1}{2}$. The following shifted energies

$$e(n) = \frac{2m_r}{\hbar^2 \beta^2} (E_n - E_0) = \epsilon_0^2 - \epsilon_n^2 = n(2p - n) \quad (15)$$

are useful for the construction of SCS. Using the change of variable $y = \nu e^{-\beta x}$, we obtain the energy eigenfunctions, for the discrete spectrum, in terms of the associated Laguerre polynomials, denoted by $L_n^{2\epsilon_n}$, as

$$\psi_n^\nu(x) = \mathcal{N}_n e^{-\frac{y}{2}} y^{\epsilon_n} L_n^{2\epsilon_n}(y), \quad (16)$$

where \mathcal{N}_n is a normalization factor given by

$$\mathcal{N}_n = \sqrt{\frac{\beta(\nu - 2n - 1)\Gamma(n + 1)}{\Gamma(\nu - n)}} = \sqrt{\frac{2\beta(p - n)\Gamma(n + 1)}{\Gamma(2p - n + 1)}}. \quad (17)$$

Since p is related to physical parameters (see (14)), it is not an integer in practice and \mathcal{N} is never zero as expected.

For many applications, it is convenient to introduce the number operator \hat{N} such that

$$\hat{N} \psi_n^\nu(x) = n \psi_n^\nu(x) \quad (18)$$

and we see from (13) that \hat{H} can in fact be written as $\hat{H} = -\frac{\hbar^2}{2m_r} \beta^2 (p - \hat{N})^2$.

3.2. Ladder operators

The ladder operators of the Morse system are defined as in (2); however, the set of eigenfunctions $\{|\psi_n\rangle\}$ is finite. As mentioned in the introduction, the quantity $k(n)$ in (2) is not unique and we consider two natural choices [20]. The first choice, the ‘oscillator-like’, corresponds to $k(n) = n$. The ladder operators satisfy an $h(2)$ algebra. The second choice, the ‘energy-like’, corresponds to $k(n) = e(n)$, as given in (15). Ladder operators satisfy an $su(1, 1)$ algebra. In what follows, the subscripts o and e will be used to refer to these choices.

Although our future calculations do not need the explicit form of the ladder operators, we give them for completeness [19, 21, 28, 29]. For example, we obtain [21]

$$A^- = - \left[\frac{d}{dy} (v - 2N) - \frac{(v - 2N - 1)(v - 2N)}{2y} + \frac{v}{2} \right] \sqrt{K(N)}, \quad (19)$$

$$A^+ = (\sqrt{K(N)})^{-1} \left[\frac{d}{dy} (v - 2N - 2) + \frac{(v - 2N - 1)(v - 2N - 2)}{2y} - \frac{v}{2} \right], \quad (20)$$

where $K(n)$ is related to $k(n)$ by

$$k(n) = \frac{n(v - n)(v - 2n - 1)}{v - 2n + 1} K(n). \quad (21)$$

These relations are valid for any integer n in the interval $[0, [p] - 1]$. Note that, for $n = [p]$, we obtain a permitted energy eigenstate $\psi_{[p]}^v(x)$ of the Morse potential but the action of the creation operators on this state does not give zero in general. It gives a state which may not be normalizable with respect to our scalar product. This problem has already been mentioned in some contributions (see, for example, [21, 28]). For arbitrary p , the special choice $k(n) = n([p] + 1 - n)$ leads to $A_+ \psi_{[p]}^v(x) = 0$ and $A_+ \psi_{[p]-1}^v(x) = \sqrt{[p]} \psi_{[p]}^v(x)$.

The ‘oscillator-like’ ladder operators are thus obtained by taking

$$K_o(n) = \frac{v - 2n + 1}{(v - n)(v - 2n - 1)}, \quad (22)$$

while we see that $K_e(n) = K_o(n)(v - 1 - n)$ for the ‘energy-like’ ladder operators.

3.3. The harmonic oscillator limit

The harmonic oscillator limit [22] is obtained by first shifting the Morse potential by $V_0 = \frac{k'}{2\beta^2}$ to obtain

$$V_1 = V_0(1 - e^{-\beta x})^2 = V_M + V_0, \quad (23)$$

and then taking $\beta \rightarrow 0$ so that $V_1 \rightarrow \frac{1}{2}k'x^2$ where k' is the force constant. Note that the new Hamiltonian with potential V_1 has thus the energy levels shifted and we obtain

$$E_n^1 = - \frac{\hbar^2}{2m_r} \beta^2 \left[\left(\frac{v - 1}{2} - n \right)^2 - \left(\frac{v}{2} \right)^2 \right]. \quad (24)$$

Since v is given by (14), we obtain here $v = \frac{2\sqrt{m_r k'}}{\beta^2 \hbar}$. The oscillator limit is obtained when $v \rightarrow \infty$ giving, as expected, an infinite spectrum and the good limit for the energies,

$$\lim_{v \rightarrow \infty} E_n^1 = \hbar \sqrt{\frac{k'}{m_r}} \left(n + \frac{1}{2} \right).$$

Second, we have to take the limit on the ladder operators. We replace β by its expression in terms of v and define $c = \sqrt{\frac{4m_r k'}{\hbar^2}}$. The annihilation operator A^- , given in (19), thus takes the form

$$A^- = \sqrt{K(n)} \left[\frac{e^{\sqrt{\frac{c}{v}}x}}{\sqrt{c}v} (v - 2n) \frac{d}{dx} + \frac{e^{\sqrt{\frac{c}{v}}x}}{2v} (v - 2n - 1)(v - 2n) - \frac{v}{2} \right]. \quad (25)$$

Since $K(n)$ depends also on v , we have to take the limit carefully. To solve it, just take the Taylor expansion of the exponential to the first order. We then see that $K(n)$ must behave as v^{-1} , which is exactly what we obtain from (22) and we find

$$\lim_{v \rightarrow \infty} A^- = \frac{1}{\sqrt{c}} \left(\frac{d}{dx} + \frac{c}{2}x \right). \quad (26)$$

A similar calculation gives the expected limit for A^+ .

3.4. SCS and their time evolution

The SCS of the Morse Hamiltonian are now defined as the finite sum

$$\Psi^v(z, \gamma, x) = \frac{1}{\sqrt{\mathcal{N}^v(z, \gamma)}} \sum_{n=0}^{[p]-1} \frac{Z(z, \gamma, n)}{\sqrt{\rho(n)}} \psi_n^v(x), \quad (27)$$

where $\rho(n)$ is given in (3), $Z(z, \gamma, n)$ satisfies (8) and

$$\mathcal{N}^v(z, \gamma) = \sum_{n=0}^{[p]-1} \frac{|Z(z, \gamma, n)|^2}{\rho(n)}. \quad (28)$$

Such a definition is relevant since we have seen in the preceding subsection that the ‘oscillator-like’ ladder operators tend to the ones of the harmonic oscillator when $k(n) = n$ and the appropriate limit is taken. Moreover, these states are ‘almost’ eigenstates of a linear combination of the generic ladder operators A^- and A^+ which can be written as

$$(A^- + \gamma A^+) \Psi^v(z, \gamma, x) \approx z \Psi^v(z, \gamma, x). \quad (29)$$

In fact, the correction can be computed using the recurrence relation (8) and we find

$$\chi^v(z, \gamma, [p], x) = \Lambda_1(z, \gamma, [p]) \psi_{[p]-1}^v(x) + \Lambda_0(z, \gamma, [p]) \psi_{[p]}^v(x), \quad (30)$$

where

$$\begin{aligned} \Lambda_1(z, \gamma, [p]) &= \frac{1}{\sqrt{\rho_{[p]-1}}} Z(z, \gamma, [p]), \\ \Lambda_0(z, \gamma, [p]) &= \frac{1}{\sqrt{\rho_{[p]}}} \gamma k([p]) Z(z, \gamma, [p] - 1). \end{aligned} \quad (31)$$

In practice, the last two terms of the sum in (27) have a very weak contribution which justifies thus the term ‘almost’ eigenstates used above.

Other constructions of SCS have been considered (see, for example, [19, 22]). They implicitly use the displacement operator D . It must be questioned first because we deal with a finite number of eigenstates in (27). Indeed, the action of this operator is not well defined even if we take a finite development of the exponentials. Moreover, the Baker–Campbell–Hausdorff formulae for expanding D (as products of exponentials of simple operators) are not necessarily valid (see, for example [29]). Secondly, only one parameter is involved in this displacement operator, which is the reason why they are called CS by these authors [19, 22]. They are, in fact, special cases of our SCS where z and γ are not independent ($\gamma \neq 0$).

Since our SCS are closely related to the ones of the harmonic oscillator, we are interested in the behaviour of these states in the physical observable position x and observable momentum p . Here these observables are not obtained as linear combinations of the ladder operators (as we can see from (19) and (20)) and we must compute the mean values explicitly.

For an arbitrary observable θ , we have $\langle \theta \rangle(z, \gamma; t) = \langle \Psi^v(z, \gamma, x; t) | \theta | \Psi^v(z, \gamma, x; t) \rangle$, where the time evolution of our SCS is given by

$$\Psi^v(z, \gamma, x; t) = \frac{1}{\sqrt{\mathcal{N}^v(z, \gamma)}} \sum_{n=0}^{[p]-1} \frac{Z(z, \gamma, n)}{\sqrt{\rho(n)}} e^{-\frac{iE_n}{\hbar}t} \psi_n^v(x) \quad (32)$$

and we obtain explicitly

$$\begin{aligned} \langle \theta \rangle(z, \gamma; t) &= \frac{1}{\mathcal{N}^v(z, \gamma)} \left(\sum_{n=0}^{[p]-1} \frac{|Z(z, \gamma, n)|^2}{\rho(n)} \langle \theta \rangle_{n,n} + \sum_{n=0}^{[p]-1} \sum_{k=1}^{[p]-1-n} \frac{Z^*(z, \gamma, n+k)}{\sqrt{\rho(n+k)}} \frac{Z(z, \gamma, n)}{\sqrt{\rho(n)}} \right. \\ &\quad \left. \times \left(e^{-\frac{i(E_{n+k}-E_n)}{\hbar}t} \langle \theta \rangle_{n,n+k} + e^{\frac{i(E_{n+k}-E_n)}{\hbar}t} \langle \theta \rangle_{n+k,n} \right) \right), \end{aligned} \quad (33)$$

where

$$\langle \theta \rangle_{m,n} = \langle \psi_m^v | \theta | \psi_n^v \rangle. \tag{34}$$

In the following developments, we consider observables which are such that $\langle \theta \rangle_{m,n}$ are symmetric or skew-symmetric with respect to the exchange of m and n . We are thus led to two different cases. If $\langle \theta \rangle_{n+k,n} = \langle \theta \rangle_{n,n+k}$, we obtain

$$\begin{aligned} \langle \theta \rangle(z, \gamma; t) = & \frac{1}{\mathcal{N}^v(z, \gamma)} \left(\sum_{n=0}^{[p]-1} \frac{|Z(z, \gamma, n)|^2}{\rho(n)} \langle \theta \rangle_{n,n} \right. \\ & \left. + 2 \sum_{n=0}^{[p]-1} \sum_{k=1}^{[p]-1-n} \frac{Z^*(z, \gamma, n+k) Z(z, \gamma, n)}{\sqrt{\rho(n+k)} \sqrt{\rho(n)}} \cos(\alpha(n, k)t) \langle \theta \rangle_{n+k,n} \right), \end{aligned} \tag{35}$$

where

$$\alpha(n, k) = \frac{\hbar\beta^2}{2m_r} k(2(p-n) - k). \tag{36}$$

If $\langle \theta \rangle_{n+k,n} = -\langle \theta \rangle_{n,n+k}$, we obtain

$$\langle \theta \rangle(z, \gamma; t) = \frac{2i}{\mathcal{N}^v(z, \gamma)} \sum_{n=0}^{[p]-1} \sum_{k=1}^{[p]-1-n} \frac{Z^*(z, \gamma, n+k) Z(z, \gamma, n)}{\sqrt{\rho(n+k)} \sqrt{\rho(n)}} \sin(\alpha(n, k)t) \langle \theta \rangle_{n+k,n}. \tag{37}$$

We obtain, from [30], the mean values of \hat{x} and \hat{p} (with $\beta = 1$),

$$\langle \hat{x} \rangle_{n+k,n} = (-1)^{k+1} \mathcal{N}_{n+k} \mathcal{N}_n \frac{\Gamma(v-k-n)}{k(v-k-1-2n)n!}, \quad k \neq 0, \tag{38}$$

$$\langle \hat{x} \rangle_{n,n} = \ln v - \Phi(0, v-1-2n) + \sum_{j=1}^n \frac{1}{v-n-j}, \tag{39}$$

where the function $\Phi(0, z) = \frac{d}{dz} \ln \Gamma(z)$ and

$$\langle \hat{p} \rangle_{n+k,n} = i\hbar(-1)^{k+1} \mathcal{N}_{n+k} \mathcal{N}_n \frac{\Gamma(v-k-n)}{2n!} (1 - \delta_{k0}). \tag{40}$$

We also obtain after some calculations

$$\langle \hat{p}^2 \rangle_{n+k,n} = \hbar^2 (-1)^{k+1} \mathcal{N}_{n+k} \mathcal{N}_n \frac{\Gamma(v-k-n)}{4n!} ((k-1)v - k(k+2n+1)), \quad k \neq 0 \tag{41}$$

and

$$\langle \hat{p}^2 \rangle_{n,n} = -\hbar^2 \frac{(2n+1)(2n+1-v)}{4}. \tag{42}$$

The computation of the mean values of \hat{x}^2 is more tricky since it involves the functions $\Phi(0, z)$ and $\Phi(1, z)$. We do not have an analytic expression but will be able to compute explicitly $\langle \hat{x}^2 \rangle_{n+k,n}$ and $\langle \hat{x}^2 \rangle_{n,n}$ since we have a finite number of these expressions to plug in $\langle \hat{x}^2 \rangle(z, \gamma; t)$.

3.5. Oscillator-like SCS

In this case, in accordance with the expression of $\rho(n)$ given in (3), we take $k(i) = i$ and $\rho(n) = n!$ and we obtain $Z_o(z, \gamma, n) = Z_{ho}(z, \gamma, n) = (9)$. For the special case where $\gamma = 0$, we obtain $Z_{ho}(z, 0, n) = z^n$, while for the squeezed vacuum $z = 0$, we obtain

$$Z_o(0, \gamma, 2n) = \frac{(2n)!}{n!} \left(-\frac{\gamma}{2}\right)^n, \quad Z_o(0, \gamma, 2n+1) = 0. \tag{43}$$

Moreover, in those states, we obtain the same probability distribution as for the harmonic oscillator:

$$P_o(z, \gamma, n) = |\langle \psi_n^v(x) | \psi(z, \gamma, x) \rangle|^2 = \frac{1}{\mathcal{N}_o^v(z, \gamma)} \left(\frac{|\gamma|}{2} \right)^n \frac{|\mathcal{H}(n, \frac{z}{\sqrt{2\gamma}})|^2}{n!}, \quad (44)$$

with

$$\mathcal{N}_o^v(z, \gamma) = \sum_{n=0}^{[p]-1} \left(\frac{|\gamma|}{2} \right)^n \frac{|\mathcal{H}(n, \frac{z}{\sqrt{2\gamma}})|^2}{n!}. \quad (45)$$

The mean value and dispersion of the number operator \hat{N} are now given by

$$\langle \hat{N} \rangle_o = \sum_{n=0}^{[p]-1} n P_o(z, \gamma, n), \quad (\Delta \hat{N})_o^2 = \sum_{n=0}^{[p]-1} n^2 P_o(z, \gamma, n) - \left(\sum_{n=0}^{[p]-1} n P_o(z, \gamma, n) \right)^2. \quad (46)$$

The statistical properties of these states are similar to the ones of the harmonic oscillator since we obtain essentially the same quantity for Mandel's Q -parameter [31] given in general by

$$Q(z, \gamma) = \frac{(\Delta \hat{N})^2 - \langle \hat{N} \rangle}{\langle \hat{N} \rangle}. \quad (47)$$

The only difference is that, in the calculation of the dispersion and mean values in \hat{N} , the sums are now finite. In particular, it is well known (see, for example, [7]) that the probability density is a Poisson distribution in the special coherent case ($\gamma = 0$).

3.6. Energy-like SCS

In this case, in accordance with the expression of $\rho(n)$ given in (3), we take $k(i) = i(2p - i)$ and $\rho(n) = (-1)^n n!(1 - 2p)_n$ where $(a)_n$ is the usual notation for the Pochhammer symbol

$$(a)_n = a(a + 1)(a + 2) \cdots (a + n - 1) = \frac{\Gamma(a + n)}{\Gamma(a)}. \quad (48)$$

The exact (infinite) recurrence relation (8) can be solved directly in terms of hypergeometric functions. The solution is

$$Z_e(z, \gamma, n) = (-1)^n \gamma^{\frac{n}{2}} \frac{\Gamma(2p)}{\Gamma(2p - n)} {}_2F_1 \left(\begin{matrix} -n, -\frac{z}{2\sqrt{\gamma}} + \frac{1-2p}{2} \\ 1 - 2p \end{matrix}; 2 \right), \quad n = 1, 2, \dots, [p] - 1. \quad (49)$$

Since this result is far from being trivial, we give some details of the proof and also the expressions of few first polynomials of this sequence.

Let us first set

$$k(n) = n(A - n), \quad A \in \mathbb{R}, \quad (50)$$

so that the recurrence relation (8) becomes

$$Z(z, \gamma, n + 1) - z Z(z, \gamma, n) + \gamma n(A - n) Z(z, \gamma, n - 1) = 0, \quad n = 1, 2, \dots, \quad (51)$$

with $Z(z, \gamma, 0) = 1$ and thus $Z(z, \gamma, 1) = z$. Since we know the solution for the harmonic oscillator (i.e. when $k(n) = n$), we follow the same lines to solve (51) for an infinite sequence of values of n . We introduce the new complex variable $w = \frac{z}{\sqrt{2\gamma}}$ and take

$$Z(z, \gamma, n) = \left(\frac{\gamma}{2} \right)^{\frac{n}{2}} f(n, w). \quad (52)$$

We thus obtain a new recurrence relation on the functions $f(n, w)$:

$$\begin{aligned} f(n+1, w) - 2w f(n, w) + 2n(A-n) f(n-1, w) &= 0, \\ f(1, w) = 2w, \quad f(0, w) = 1, \quad n = 1, 2, \dots \end{aligned} \tag{53}$$

It is easy to see that $f(n, w)$ is in fact a polynomial of degree n in w . Moreover, it can be expressed in terms of hypergeometric functions of the type ${}_2F_1$. We explicitly obtain

$$f(n, w) = 2^{\frac{n}{2}} (-A+1)_n {}_2F_1 \left(\begin{matrix} -n, -\frac{w}{\sqrt{2}} + \frac{1-A}{2} \\ 1-A \end{matrix}; 2 \right) \tag{54}$$

and the hypergeometric function is in fact a polynomial in w since we have

$${}_2F_1 \left(\begin{matrix} -n, -v \\ 1-A \end{matrix}; 2 \right) = \sum_{k=0}^n \frac{2^k (-n)_k (-v)_k}{k! (-A+1)_k}. \tag{55}$$

The original function (52) thus takes the form (49) when $A = 2p$ as expected. It is valid for any real value of A and in fact, we see that the first polynomials of the sequence are given by

$$\begin{aligned} Z_e(z, \gamma, 0) &= 1, \quad Z(z, \gamma, 1) = z, \\ Z_e(z, \gamma, 2) &= z^2 - (A-1)\gamma, \\ Z_e(z, \gamma, 3) &= z^3 - (3A-5)\gamma z, \\ Z_e(z, \gamma, 4) &= z^4 - 2(3A-7)z^2\gamma + 3(A-1)(A-3)\gamma^2. \end{aligned}$$

To be complete, let us mention that for the special case where A is an integer, we see that the recurrence relation (53) splits into two different ones. Indeed, we first obtain a finite sequence of $f(n, w)$ satisfying (53) for $n = 1, 2, \dots, A-1$ and second, an infinite sequence of $f(n, w)$ for $n = A, A+1, \dots$ satisfying the recurrence relation

$$f(A+k+1, w) - 2w f(A+k, w) - 2k(A+k) f(A+k-1, w) = 0, \quad k = 0, 1, 2, \dots \tag{56}$$

Since for $k = 0$ we obtain $f(A+1, w) = 2w f(A, w)$, we can write $f(A+k, w) = h(k, w) f(A, w)$ where $h(k, w)$ is a polynomial of degree k in w satisfying the recurrence relation

$$\begin{aligned} h(k+1, w) - 2w h(k, w) + 2k(-A-k) h(k-1, w) &= 0, \\ h(1, w) = 2w, \quad h(0, w) = 1, \quad k = 1, 2, \dots, \end{aligned} \tag{57}$$

which is (53) where A has been replaced by $-A$. The polynomials $h(k, w)$ are thus given by

$$h(k, w) = 2^{\frac{k}{2}} (A+1)_k {}_2F_1 \left(\begin{matrix} -k, -\frac{w}{\sqrt{2}} + \frac{1+A}{2} \\ 1+A \end{matrix}; 2 \right), \quad k = 0, 1, 2, \dots \tag{58}$$

The solutions $f(n, w)$ satisfying (53) for $n = 1, 2, \dots, A-1$ are in fact associated with a finite sequence of Krawtchouk polynomials, while the solutions $h(n, w)$ for $n = 0, 1, \dots$ are associated with Meixner polynomials [32]. They both satisfy discrete orthogonality relations on the variable w but these are not relevant in our context since w is a continuous parameter.

When $\gamma = 0$, we obtain $Z_e(z, 0, n) = z^n$ leading to CS, while for the squeezed vacuum ($z = 0$), we obtain

$$Z_e(0, \gamma, 2n) = 4^{(n-1)} (1-2p)(3/2)_{n-1} (3/2-p)_{n-1} \gamma^n, \quad Z_e(0, \gamma, 2n+1) = 0. \tag{59}$$

Now the probability distribution, denoted by $P_e(z, \gamma, n)_e$, is given by

$$P_e(z, \gamma, n) = \frac{1}{\mathcal{N}_e^v(z, \gamma)} \frac{\Gamma(2p - n)}{\Gamma(2p)n!} |Z(z, \gamma, n)|^2, \quad (60)$$

where

$$\mathcal{N}_e^v(z, \gamma) = \sum_{n=0}^{\lfloor p \rfloor - 1} \frac{\Gamma(2p - n)}{\Gamma(2p)n!} |Z(z, \gamma, n)|^2. \quad (61)$$

Similar expressions for $\langle \hat{N} \rangle_e$ and $(\Delta \hat{N})_e^2$ are obtained as in (46).

4. Behaviour of the SCS for the case of the hydrogen chloride molecule

For the hydrogen chloride molecule $^1\text{H}^{35}\text{Cl}$ (already considered in a previous paper [20]), we will fix the values of the physical parameter ν given in (14) with published values of m_r , β and V_0 , or as most often in practice, using $\nu = \omega_e/\omega_e x_e$, the ratio between the experimentally measured molecular harmonicity ω_e and anharmonicity $\omega_e x_e$ constants (see, for example, [33–36]). For the ground state, $X^1\Sigma^+$, we have $\nu \approx 57.44$ and $[p] = 28$. We also choose the units such that $\frac{\hbar}{2m_r} = 1$ and $\beta = 1$.

Let us start with some general facts. First, the squeezing parameter must be restricted to $|\gamma| < 1$ for the states to be normalizable (as for the harmonic oscillator). Second, the minimization of the Heisenberg uncertainty relation depends on the values of z when γ is real (and less than 1). Finally, we will see that the squeezing effect (reduction of the dispersion of one observable at the price of increasing it in the other) is always present even for $\gamma = 0$. Moreover, the dispersion in \hat{x} may be chosen smaller for the energy-like than for the oscillator-like states leading to a better localization for the energy-like states.

In order to compare our results with the well-known ones for the harmonic oscillator, we restrict ourselves to the positive real values of z and γ with $\gamma < 1$. In fact, no dramatic difference appears when z and/or γ are negatives.

4.1. Coherent system of states ($\gamma = 0$)

The oscillator-like and energy-like states differ only by the denominator (and the normalization factor) in development (27). In figure 1, we show the trajectories for both states with $z = 2$ and $t \in [0, 1]$ and we see that the oscillator-like states are less stable than the energy-like ones for the same value of z . This is true when $z < 20$. We also see that, even if $\gamma = 0$, squeezing always appears in the coherent system of states for the Morse potential.

At $t = 0$, the uncertainty product $\Delta(z, 0) = (\Delta x)^2(\Delta p)^2$ and the dispersion $(\Delta x)^2$ show a very stable behaviour in z for the energy-like states (see figure 2). These states are mostly minimal uncertainty states and the dispersion in x is very small leading to a very good localization (at least for $z < 20$). This last fact is confirmed by the large eccentricity (in fact, it is very close to 1) of the ellipses obtained before. Interestingly, if we want to minimize the dispersion in p , we see, again in figure 2, that $\Delta(z, 0)$ is no longer minimized. Also, reducing the dispersion in p increases the dispersion in x . Moreover, we can easily see that the best minimization of the dispersion in p is observed in this special case, i.e. for $\gamma = 0$. The oscillator-like states have almost the same behaviour but $\Delta(z, 0)$ increases faster for smaller values of z (around $z = 3$).

Finally, the density probabilities of both types of states have been computed to confirm the preceding results. In figure 3, the energy-like states show again a better behaviour than the oscillator-like states as time evolves.




Figure 1. Phase-space trajectories for oscillator-like (dashed line) and energy-like (plain line) states when $(z, \gamma) = (2, 0)$ and $t \in [0, 1]$.




Figure 2. Uncertainty and dispersions for the energy-like states at $\gamma = 0$, Δ and $(\Delta \hat{x})^2$ with $z \in [0, 25]$ (left) and Δ , $(\Delta \hat{x})^2$ and $(\Delta \hat{p})^2$ with $z \in [10, 27]$ (right).

4.2. Squeezed vacuum ($z = 0$)

In this case, both types of states are polynomials in γ and only even combinations of eigenfunctions appear (see (43) and (59)). Again, the energy-like states have more stable trajectories but for small values of γ . In figure 4, we have compared the trajectories for the energy-like states when $\gamma = 0.2, 0.5$ and 0.7 .




Figure 3. Density probability $|\Psi_e^v(1, 0, x; t)|^2$ (left) and $|\Psi_o^v(1, 0, x; t)|^2$ (right) for $x \in [-1, 2]$ and $t \in [0, 1]$.




Figure 4. Phase-space trajectories for energy-like states in the vacuum with $\gamma = 0.2, 0.5, 0.7$ and $t \in [0, 1]$.

At $t = 0$, the uncertainty product $\Delta(0, \gamma) = (\Delta x)^2(\Delta p)^2$ and the dispersion $(\Delta x)^2$ take almost the same values for both types of states. Minimal uncertainty is satisfied for $\gamma \in [0, 0.2]$ and we obtain very small values of the dispersion in x for the same values of γ . The behaviour is similar to the one in figure 2.

With respect to the statistical properties, figure 5 shows bunching effects of $Q(0, \gamma) > 0$ both for the energy-like and oscillator-like states. It confirms that both types of states have similar statistical behaviour. Finally, the density probability has been computed for the energy-like states. In figure 6, we see that the best localization occurs when γ is smaller than 0.2. We see also, by observing the time evolution, that the vacuum states are less stable than the CS. Similar behaviour is observed for the oscillator-like states.




Figure 5. Comparison of the Mandel parameter $Q(0, \gamma)$ in the vacuum for the energy-like and oscillator-like squeezed states as a function of r such that $\gamma = \tanh r$.




Figure 6. Density probability $|\Psi_e^v(0, \gamma, x; t)|^2$ in the vacuum for energy-like CS for $x \in [-1, 2]$ as a function of $\gamma \in [0, 1]$ with $t = 0$ (left) and a function of $t \in [0, 1]$ with $\gamma = 0.2$ (right).




Figure 7. Comparison between the Mandel parameter for the energy-like and oscillator-like squeezed states as a function of r such that $\gamma = \tanh r$ for $z = 2$.

4.3. General system of states

In this case, we observe that the trajectories deviate quickly from closed curves while the minimum uncertainty is still realized but when $\gamma < 0.2$. The graphs for the Mandel parameter for $z = 2$ are given in figure 7. Bunching and anti-bunching are observed for both types

of states; the bunching is more prominent for energy-like states for all values of r . A more significant anti-bunching effect is observed for oscillator-like states showing a steady effect for $r < 0$ and a minimum for $r > 0$ (when $\gamma \approx 0.2$).

The calculations of observables for other diatomic molecules can be performed in a similar way. For example, we have performed such calculations for the molecule $^{133}\text{Cs}_2$ which has a larger value for ν . Indeed, we have $\nu \approx 524.55$ and thus $[p] = 261$. We have obtained similar behaviour for both types of states that is not relevant to produce here.

5. Conclusions

In this paper, we have introduced squeezed coherent states of a quantum system with a finite discrete energy spectrum described by the Morse potential. These states are almost eigenstates of linear combination of ladder operators and are characterized by two continuous parameters z and γ . We have considered two different types of ladder operators and constructed the corresponding oscillator-like and energy-like squeezed coherent states.

We have investigated the behaviour of these states regarding localization and minimum uncertainty. The calculation of the dispersions and mean values has been performed analytically except for the mean value of \hat{x}^2 for which the analytical form is not known. We have computed the Mandel parameter to investigate the statistical properties of our states.

The oscillator-like squeezed coherent states are closely related to the similar states of the harmonic oscillator. However, they do not have a very good localization except for the vacuum case for γ small and they exhibit a certain deviation from the minimal uncertainty principle. The energy-like squeezed coherent states minimize better the uncertainty relation for the case $\gamma = 0$ and we get a good localization in position. They are more stable in time than the oscillator-like states. Both types of states have similar statistical properties.

Acknowledgments

VH acknowledges the support of research grants from NSERC of Canada. AH acknowledges the support of a NSERC research fellowship. This work was started while VH visited Northumbria University (as a visiting professor and on sabbatical leave). This institution is acknowledged for hospitality. The authors thank J Van der Jeugt for helpful discussions on special functions and orthogonal polynomials.

References

- [1] Schrödinger E 1926 *Naturwiss* **14** 664–6
- [2] Kennard E H 1927 *Z. Phys.* **44** 326–52
- [3] Glauber R J 1963 *Phys. Rev.* **130** 2529–39
Glauber R J 1963 *Phys. Rev.* **131** 2766–88
- [4] Klauder J R 1960 *Ann. Phys.* **11** 123–68
- [5] Klauder J R and Skagerstam B S 1985 *Coherent States—Applications in Physics and Mathematical Physics* (Singapore: World Scientific)
- [6] Nieto M M 1997 arXiv:[quant-ph/9708012](https://arxiv.org/abs/quant-ph/9708012)
- [7] Walls D F and Milburn G J 2008 *Quantum Optics* 2nd edn (Berlin: Springer)
- [8] Gazeau J P 2009 *Coherent States in Quantum Physics* (New York: Wiley)
- [9] Rand S C 2010 *Lectures on Light: Nonlinear and Quantum Optics Using Density Matrix* (Oxford: Oxford University Press)
- [10] Braunstein S L and McLachlan R I 1987 *Phys. Rev. A* **35** 1659–67
- [11] Hillery M 1987 *Phys. Rev. A* **36** 3796–802
- [12] Bergou J A, Hillery M and Yu D 1991 *Phys. Rev. A* **43** 515–20

- [13] Fu H-C and Sasaki R 1996 *Phys. Rev. A* **53** 3836–44
- [14] Alvarez N and Hussin V 2002 *J. Math. Phys.* **43** 2063–85
- [15] Klauder J R, Penson K A and Sixderniers J M 2001 *Phys. Rev. A* **64** 013817
- [16] Dong S H 2002 *Can. J. Phys.* **80** 129–39
- [17] Roy B and Roy P 2002 *Phys. Lett. A* **296** 187–91
- [18] Récamier J and Jäuregui R 2003 *J. Opt. B: Quantum Semiclass. Opt.* **5** S365–70
- [19] Daoud M and Popov D 2004 *Int. J. Mod. Phys. B* **18** 325–36
- [20] Angelova M and Hussin V 2008 *J. Phys. A: Math. Theor.* **41** 304016
- [21] Dong S H, Lemus R and Frank A 2002 *Int. J. Quantum Chem.* **86** 433–9
- [22] Dong S-H 2008 *Factorization Method in Quantum Mechanics (Fundamental Theories in Physics vol 150)* (Dordrecht: Springer)
- [23] Fox R F and Choi M F 2001 *Phys. Rev. A* **64** 042104
- [24] Draganescu G E, Messina A and Napoli A 2009 *J. Mod. Opt.* **56** 508–15
- [25] Yuen H P 1976 *Phys. Rev. A* **13** 2226–43
- [26] Szafraniec F H 1998 *Contemp. Math.* **212** 269–76
- [27] Merzbacher E 1998 *Quantum Mechanics* (New York: Wiley)
- [28] Singh A C and Babynanda D O 2006 *Int. J. Quantum Chem.* **106** 415–25
- [29] Odake S and Sasaki R 2006 *J. Math. Phys.* **47** 102102
- [30] Sage M L 1978 *Chem. Phys.* **35** 375–80
- [31] Mandel L 1979 *Opt. Lett.* **4** 205–7
- [32] Koekoek R and Swarttouw R F 1998 The Askey-scheme of hypergeometric orthogonal polynomials and its q -analogue *Technical Report 98-17* (Delft: Delft University of Technology) (<http://fa.its.tudelf.nl/koekoek/askey/contents.html>)
- [33] Herzberg G 1950 *Molecular Spectra and Molecular Structure: Vol I. Spectra of Diatomic Molecules* 2nd edn (Princeton, NJ: Van Nostrand-Reinhold)
- [34] Guelachvili G, Noah P and Bedace P 1981 *J. Mol. Spectrom.* **85** 271–81
- [35] Lide D R (ed) 2010 *CRC Handbook of Chemistry and Physics* 90th edn (London: Taylor and Francis) section 9 (Molecular structure and spectroscopy)
- [36] Angelova M and Frank A 2005 *Phys. At. Nuclei* **68** 1625