

Northumbria Research Link

Citation: Mantso, Theodora, Sfakianos, Aristeidis, Atkinson, Aithne, Anastopoulos, Ioannis, Mitsiogianni, Melina, Botaitis, Sotirios, Perente, Sebachedin, Simopoulos, Constantinos, Vasileiadis, Stavros, Franco, Rodrigo, Pappa, Aglaia and Panagiotidis, Mihalis (2016) Development of a Novel Experimental In Vitro Model of Isothiocyanate-induced Apoptosis in Human Malignant Melanoma Cells. *Anticancer Research*, 36 (12). pp. 6303-6310. ISSN 0250-7005

Published by: International Institute of Anticancer Research

URL: <https://doi.org/10.21873/anticanres.11226>
<<https://doi.org/10.21873/anticanres.11226>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/29218/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

Development of a Novel Experimental *in Vitro* Model of Isothiocyanate-Induced Apoptosis
in Human Malignant Melanoma Cells

THEODORA MANTSO^{1*}, ARIS SFAKIANOS^{1*}, AITHNE ATKINSON¹, IOANNIS
ANESTOPOULOS², MELINA MITSIOGIANNI³, SOTIRIOS BOTAITIS⁴, SEBACHEDIN
PERENTE⁴, CONSTANTINOS SIMOPOULOS⁴, STAVROS VASILEIADIS², RODRIGO
FRANCO^{5,6}, AGLAIA PAPPA² and MIHALIS I. PANAYIOTIDIS^{1,3}

*These Authors contributed equally to this work

¹School of Life Sciences, Heriot Watt University, Edinburgh, Scotland, UK;

²Department of Molecular & Cell Biology, Democritus University of Thrace,
Alexandroupolis, Greece;

³Department of Applied Sciences, Northumbria University, Newcastle Upon Tyne, UK;

⁴Second Department of Surgery, Democritus University of Thrace, Alexandroupolis, Greece;

⁵Redox Biology Centre, University of Nebraska, Lincoln, NE, USA;

⁶School of Veterinary Medicine & Biomedical Sciences, University of Nebraska, Lincoln,
NE, USA

Running title: Isothiocyanates-induced apoptosis in malignant melanoma

Correspondence to: Professor Mihalis I. Panayiotidis, Department of Applied Sciences,
Northumbria University, Ellison Building A516, Newcastle Upon Tyne, NE1 8ST, UK, tel:
+44 01912274305, e-mail: m.panagiotidis@northumbria.ac.uk

Key Words: Isothiocyanates, sulforaphane, phenethyl isothiocyanate, benzyl isothiocyanate, malignant melanoma, apoptosis

Abstract. Background. Isothiocyanates are constituents of cruciferous vegetables which have been associated with reduced cancer risk partially through their ability to induce apoptosis in malignant cells including melanoma. Materials and Methods: We have utilized human malignant melanoma (A375), epidermoid carcinoma (A431) and immortalized keratinocyte (HaCaT) cells exposed to various isothiocyanates, under different experimental conditions.

Results: An experimental *in vitro* model utilizing low isothiocyanate concentrations (0.1-5 μ M for 48h with all treatments being refreshed after 24h) was shown to be (i) most efficient in exerting an anti-cancer effect when compared to higher concentrations (5-100 μ M for 24 or 48h added as a single bolus) and (ii) specific to A375 cells while A431 and HaCaT cells remained unaffected. Such effect involved the activation of several caspases including (iii) initiator caspases 8, 9, 4 (indicating the involvement of intrinsic, extrinsic and endoplasmic reticulum-based pathways) and (iv) effector caspases 3, 7 and 6.

Conclusions: Utilization of low isothiocyanate concentrations (under conditions described herein) exerts an anti-cancer effect specific to human malignant melanoma cells and thus providing a therapeutic basis for their utilization in management of the disease.

Malignant melanoma is the fifth most common cancer in UK with its incidence rates being continuously rising faster than any other malignancy (1). Epidemiological studies suggest that an increased dietary consumption of cruciferous vegetables can reduce cancer incidence. These effects can be attributed to the high levels of glucosinolates (GSLs) present which are sulphur-containing glycosides and precursors for a group of compounds called isothiocyanates (ITCs) (2). Briefly, upon mechanical disruption of the plant cell wall (*e.g.* by

chewing), the enzyme myrosinase is released which then catalyses the hydrolysis of GSLs to ITCs, with subsequent release of HSO_4^- and D-glucose (3). Different GSLs can form different ITCs, since glucoraphanin acts as the precursor for sulforaphane (SFN), gluconasturtin for phenethyl isothiocyanate (PEITC) and glucotropaeolin for benzyl isothiocyanate (BITC) (4). There is much speculation as to how ITCs may exhibit their chemotherapeutic effects, but the likelihood is that multiple molecular events are responsible. Potential biochemical mechanisms include (i) inhibition of carcinogen activity *via* suppression of phase I enzymes in xenobiotic metabolism, (ii) stimulation of phase II enzymes and (ii) induction of apoptosis (5, 6). Despite many reports demonstrating ITCs' effectiveness against different cancers there have been a limited number of studies investigating their ability to induce apoptosis in human malignant melanoma cells (7) which their results are dependent on the utilization of high concentrations of ITCs.

In this study, we have aim to develop a novel experimental *in vitro* model based on the utilization of low ITC concentrations capable of inducing apoptosis in human malignant melanoma (A375) cells. For this reason, our objectives were to (i) establish an experimental model of exposure to various ITCs and, furthermore, to (ii) characterize ITCs' anti-cancer activities by determining the induction of a number of caspases descriptive of various apoptotic pathways including those of intrinsic, extrinsic and ER-dependent origin.

Materials and Methods

Chemicals. SFN was obtained from Abcam (Cambridge, UK) whereas PEITC and BITC from Sigma-Aldrich (St. Louis, MO, USA). All compounds were dissolved in DMSO (Sigma-Aldrich) and stored at -20°C . DMED medium, trypsin, PBS, FBS, L-Glutamine and Penicillin/Streptomycin were obtained from Labtech International Ltd (East Sussex, UK). Resazurin sodium salt was supplied by Sigma-Aldrich. All chemicals were of analytical

grade and purchased from Sigma-Aldrich, Applichem (Darmstadt, Germany) and Invitrogen (Carlsbad, CA, USA). BSA was obtained from Biosera (Boussens, France). Protease inhibitor cocktail was obtained from Roche (Basel, Switzerland). PVDF membranes (0.45 and 0.2 μm) were purchased from Millipore (Bedford, MA, USA). The pre-stained pink protein ladder was from Nippon Genetics Europe (Dueren, Germany). Primary (anti-PARP, anti-procaspase 3, anti-procaspase 6, anti-procaspase 7 and anti-lamin A/C) and horseradish peroxidase-conjugated secondary antibodies were purchased from Cell Signalling Technology (Danvers, MA, USA) whereas anti- β -actin from Sigma-Aldrich and enhanced chemi-luminescence western blotting substrate kit from Pierce Thermo Scientific (Waltham, MA, USA).

Cell culture and treatment protocols. A375 and A431 cells were purchased from Sigma-Aldrich (St. Louis, MO, USA). HaCaT cells were a kind gift from Dr Sharon Broby (Dermal Toxicology and Effects Group; Centre for Radiation, Chemical and Environmental Hazards; Public Health England, UK). All cell lines were cultured in DMEM medium (10% FBS, 2mM L-Glutamine and 1% Penicillin/Streptomycin) and maintained in a humidified atmosphere at 37°C and 5% CO₂. SFN, PEITC, BITC or vehicle were added either as (i) a single bolus concentration ranging between 5-100 μM for 24 to 48h or (ii) at 0.1-5 μM for 48h with all treatments being refreshed after 24h.

Cell viability assay. Cells (5×10^3) were seeded in 100 μl of complete medium into 96-well plates and kept overnight in the incubator before they were treated with either one of the treatment protocols. Cell viability was assessed by using the Alamar blue assay where in brief, resazurin sodium salt was dissolved in PBS (1mg/ml final concentration) and 10 μl of reagent were added into each well of the 96-well plates. After 4h of incubation, fluorescence

intensity was measured at 560_{Exc}/590_{Em} nm by using a Spectramax M5 multi-mode plate reader (Sunnyvale, CA, USA).

Caspase activity assay. The caspase family fluorometric substrate kit II plus from Abcam (Cambridge, UK) was used according to the manufacturer's instructions. Cell lysates were prepared and 150-200 µg of protein extracts were incubated with each respective substrate for 1h at 37°C. Fluorescence intensity was monitored at 400_{Exc}/505_{Em}nm by using a Spectramax M5 multi-mode plate reader

Preparation of cell lysates and protein determination. Cells (0.8×10^6) were plated in 100mm dishes and cultured overnight at 37°C. Next day, cells were treated with 0.1-5 µM of each ITC for 48h (with all treatments being refreshed after 24h) and then trypsinized, collected in micro-centrifuge tubes and washed twice with PBS. Cell pellets were lysed in lysis buffer (10mM HEPES at pH 7.9, 10mM KCl, 0.1mM EDTA, 1.5mM MgCl₂, 0.2% NP40) and supplemented with a protease inhibitors' cocktail (20mM β-glycerophosphate and 0.2mM Na₃VO₄). Then, they were left on ice while periodically vortexed over a 30 min period. Cell lysates were centrifuged at 14,000 x g for 15min at 4°C and supernatants were transferred in fresh tubes. Protein content was determined by utilizing the BCA protein assay from Thermo Scientific (Waltham, MA, USA) according to manufacturer's instructions. Protein extracts were stored at -20°C until usage.

Western immunoblotting. Forty micrograms (40 µg) of cellular protein extracts were separated by SDS-polyacrylamide gels and transferred electrophoretically onto PVDF membranes (either 0.45 or 0.2 µm). The blots were blocked in 5% non-fat milk powder in TBST buffer (50mM Tris-HCl, 150mM NaCl at pH 7.6 and 0.1% Tween-20) for 2h at room

temperature. The membranes were incubated with the appropriate primary antibody (1:1000 for anti-PARP, anti-procaspase 3, anti-procaspase 6, anti-procaspase 7, anti-lamin A/C and 1:2000 for anti- β -actin) overnight at 4°C under agitation. Next day, the membranes were incubated with the appropriate (mouse or rabbit) horseradish peroxidase-conjugated secondary antibody (1:2000) for 1h at room temperature under agitation, after being washed three times with TBST. Membranes were washed three times with TBST and once with distilled water. Labeled protein bands were detected by utilizing enhanced chemiluminescence.

Statistical analysis. Measurements for cell viability assays were performed in quintuplicate (n=5) while those for caspase activity assays in triplicates (n=3) and all values are expressed as mean \pm standard error. Statistical analysis was performed by means of a Student *t*-test. A value of $P < 0.05$ was considered statistically significant.

Results

The effect of three different ITCs (SFN, PEITC and BITC) was evaluated at a range of single bolus concentrations (5-100 μ M) in both human A375 malignant melanoma (Figure 1A-C) and HaCaT immortalized keratinocyte cells (Figure 1D-F) after 24h and 48h of treatment. Addition of a single bolus of each ITC concentration statistically significantly decreased ($P < 0.05$) both A375 and HaCaT cell viability levels in a concentration-dependent manner. More specifically, SFN decreased cell viability at 10 μ M onwards (24-48h) in A375 (Figure 1A) and HaCaT (Figure 1D) cells. PEITC also decreased cell viability levels but at a concentration of 25 μ M onwards (24-48h) for A375 (Figure 1B) and 10 μ M onwards (24-48h) for HaCaT (Figure 1E) cells. BITC exerted the highest potency, as it induced a

significant reduction in cell viability levels at 5 μM onwards (24-48h) in both A375 (Figure 1C) and HaCaT (Figure 1F) cells.

When utilized an *in vitro* model of ITC treatment by subjecting human malignant melanoma (A375) and non-melanoma (A431) cells as well as human normal keratinocyte (HaCaT; control) cells to a range of low ITC concentrations (0.1-5 μM for 48h but being refreshed after an initial 24h of treatment) our data revealed that SFN and PEITC statistically significantly inhibited ($P<0.05$) cell viability levels at 5 μM in A375 cells only (Figure 2A and 2B respectively). However, at this concentration, cells exposed to SFN showed a more profound inhibitory effect on viability levels compared to PEITC (approximately 60% vs. 80% respectively). Finally, treatment with BITC also, statistically significantly decreased ($P<0.05$) the viability levels at concentrations as low as 2.5 μM onwards in A375 and A431 cells only. However, this effect was more profound in A375 cells where a 2.5 μM inhibition of cell viability was recorded to levels of ~70% which was further decreased to ~ 30% after treatment with 5 μM of BITC (Figure 2C).

Finally, in order to investigate whether treatment with ITCs induces an apoptotic response in A375 cells, the activity levels of the initiator caspases 4, 8 and 9 were evaluated. Activation of these caspases is indicative of the involvement of various apoptotic pathways in a manner where caspase 8 involves the extrinsic (29), caspase 9 the intrinsic (30) and caspase 4 the endoplasmic reticulum (ER)-dependent (31) pathways. Our data showed that treatment with BITC statistically significantly increased ($P<0.05$) the activity levels of all initiator caspases whereas this was the case only for caspase 9 in PEITC treated cells. Alternatively, treatment with SFN showed no statistically significant changes in the activity levels of any of the above-mentioned caspases (Figure 3A-C). Similarly, western blot analysis data indicate that each of the SFN, PEITC and BITC were responsible for inducing a statistically significant decrease ($P<0.05$) in protein expression levels of the effector procaspases 3

(Figure 3D) and 6 (Figure 3E) while for procaspase 7 this was the case only for BITC treatment (Figure 3F). Consequently, when these procaspases become activated they cleave specific cellular proteins including poly(ADP-ribose) polymerase (PARP) (Figure 3D) and lamins A and C (Figure 3E) such as in the case of procaspase 3 and 6, respectively. Overall, our results demonstrate that ITCs induce caspase activation in a manner which treatment with BITC results in the largest decrease in procaspase levels, followed by PEITC and SFN.

Discussion

The majority of various studies have utilized non-melanoma (8-14) and melanoma (15-19) experimental models based on a single bolus addition of large ITC concentrations where a wide range of biological effects were documented but without provision of appropriate control (non-tumorigenic) cells. Similarly, we have been able to document such anti-cancer activity in melanoma cells but also in control keratinocyte cells suggesting a non-specific potency towards any type of cell line (tumorigenic or not).

To overcome the barrier of non-specific potency, we have developed a novel experimental *in vitro* model of ITC treatment by subjecting human malignant melanoma (A375) and non-melanoma (A431) cells as well as human normal keratinocyte (HaCaT; control) cells to a range of low ITC concentrations (0.1-5 μ M) for 48h, that were refreshed after an initial 24h of treatment. Overall, our results highlight the novelty of the *in vitro* experimental model developed not only of being capable in exerting a potent anti-cancer activity (at low ITC concentrations) but also of being specific to human malignant melanoma (A375) cells while leaving unaffected human non-melanoma (A431) and normal keratinocyte (HaCaT) cells. Moreover, our results indicate that BITC (among all the three ITCs) exerts the highest anti-cancer potency and thus highlights its effectiveness as a potential anti-cancer strategy.

Finally, since ITCs have shown to exert a plurality of anti-cancer effects by targeting various cellular processes such as, apoptosis, autophagy, cell cycle, angiogenesis, epigenetic pathways, antioxidant defenses, *etc.* (20-28)], we focused on apoptosis in order to elucidate which underlined pathways were involved in accounting for the observed reduction of viability levels in A375 cells. Overall, ITCs induced apoptosis by activating a number of initiator (8, 9, 4) and effector (3, 6, 7) caspases indicating the involvement of various apoptotic pathways including but not limited, the extrinsic (caspase 8), intrinsic (caspase 9) and ER-dependent (caspase 4) pathways. In our mind, this is the first report that (i) describes a novel experimental *in vitro* model based on the utilization of low ITC concentrations in a manner described herein and (ii) how it exerts an anti-cancer effect uniquely to melanoma cells (without affecting other non-melanoma and normal keratinocyte cells) by inducing apoptosis at concentrations substantially lower than those utilized in current bibliography.

Acknowledgements

This work was supported, in part, by a Heriot Watt University (HWU) and Northumbria University (NNU) start-up funds (Professor Mihalis Panayiotidis) including a HWU (Mrs Theodora Mantso) and a NNU (Mrs Melina Mitsiogianni) PhD studentship and by a LLP Erasmus Programme (Mr Aris Sfakianos and Dr Aglaia Pappa). Furthermore, the Authors would like to thank Dr Sharon Broby (Dermal Toxicology and Effects Group; Centre for Radiation, Chemical and Environmental Hazards; Public Health England, UK) for kindly providing the human keratinocyte (HaCaT) cell line.

References

1. Markovic SN, Erickson LA, Rao RD, Weenig RH, Pockaj BA, Bardia A, Vachon CM, Schild SE, McWilliams RR, Hand JL, Laman SD, Kottschade LA, Maples WJ, Pittelkow MR, Pulido JS, Cameron JD and Creagan ET: Malignant melanoma in the 21st century, part

- 1: Epidemiology, risk factors, screening, prevention, and diagnosis. *Mayo Clin Proc* 82(3): 364-380, 2007.
2. Abdull AF and Noor NM: Cruciferous vegetables: dietary phytochemicals for cancer prevention. *Asian Pac J Cancer Prev* 14(3): 1565-1570, 2013.
3. Bones AM and Rossiter JT: The myrosinase-glucosinolate system, its organisation and biochemistry. *Physiol Plantar* 97(1): 194-208, 1996.
4. Zhang YS: Cancer-preventive isothiocyanates: measurement of human exposure and mechanism of action. *Mutat Res* 555(1-2): 173-190, 2004.
5. Clarke JD, Dashwood RH and Ho E: Multi-targeted prevention of cancer by sulforaphane. *Cancer Lett* 269(2): 291-304, 2008.
6. Yu R, Mandlekar S, Harvey KJ, Ucker DS and Kong AN: Chemopreventive isothiocyanates induce apoptosis and caspase-3-like protease activity. *Cancer Res* 58(3): 402-408, 1998.
7. Nikhil K, Sharan S, Singh AK, Chakraborty A and Roy P: Anticancer Activities of Pterostilbene-Isothiocyanate Conjugate in Breast Cancer Cells: Involvement of PPAR gamma. *PLoS One* 9(8): e104592, 2014.
8. Cheng YM, Tsai CC and Hsu YC: Sulforaphane, a Dietary Isothiocyanate, Induces G₂/M Arrest in Cervical Cancer Cells through CyclinB1 Downregulation and GADD45 β /CDC2 Association. *Int J Mol Sci* 17(9): e1530, 2016.
9. Lee CS, Cho HJ, Jeong YJ, Shin JM, Park KK, Park YY, Bae YS, Chung IK, Kim M, Kim CH, Jin F, Chang HW and Chang YC: Isothiocyanates inhibit the invasion and migration of C6 glioma cells by blocking FAK/JNK-mediated MMP-9 expression. *Oncol Rep* 34(6): 2901-2908, 2015.
10. Su JC, Lin K, Wang Y, Sui SH, Gao ZY and Wang ZG: In vitro studies of phenethyl isothiocyanate against the growth of LN229 human glioma cells. *Int J Clin Exp Pathol* 8(4): 4269-4276, 2015.
11. Lawson AP, Long MJ, Coffey RT, Qian Y, Weerapana E, El Oualid F and Hedstrom L: Naturally Occurring Isothiocyanates Exert Anticancer Effects by Inhibiting Deubiquitinating Enzymes. *Cancer Res* 75(23): 5130-5142, 2015.
12. Chen CY, Yu ZY, Chuang YS, Huang RM and Wang TC: Sulforaphane attenuates EGFR signaling in NSCLC cells. *J Biomed Sci* 22: 38, 2015.
13. Wang L, Tian Z, Yang Q, Li H, Guan H, Shi B, Hou P and Ji M: Sulforaphane inhibits thyroid cancer cell growth and invasiveness through the reactive oxygen species-dependent pathway. *Oncotarget* 6(28): 25917-25931, 2015.
14. Zhang Y, Tang L and Gonzalez V: Selected isothiocyanates rapidly induce growth inhibition of cancer cells. *Mol Cancer Ther* 2(10): 1045-1052, 2003.

15. Ni WY, Hsiao YP, Hsu SC, Hsueh SC, Chang CH, Ji BC, Yang JS, Lu HF and Chung JG: Oral administration of benzyl-isothiocyanate inhibits in vivo growth of subcutaneous xenograft tumors of human malignant melanoma A375.S2 cells. *In Vivo* 27(5): 623-626, 2013.
16. Hamsa TP, Thejass P and Kuttan G: Induction of apoptosis by sulforaphane in highly metastatic B16F-10 melanoma cells. *Drug Chem Toxicol.* 34(3): 332-340, 2011.
17. Do DP, Pai SB, Rizvi SAA and D'Souza MJ: Development of sulforaphane-encapsulated microspheres for cancer epigenetic therapy. *Int J Pharmaceutics* 386: 114-121, 2010.
18. Sharma AK, Sharma A, Desai D, Madhunapantula SV, Huh SJ, Robertson GP and Amin S: Synthesis and Anticancer Activity Comparison of Phenylalkyl Isoselenocyanates with Corresponding Naturally Occurring and Synthetic Isothiocyanates. *J Med Chem* 51(24): 7820-7826, 2008.
19. Thejass P and Kuttan G: Modulation of cell-mediated immune response in B16F-10 melanoma-induced metastatic tumor-bearing C57BL/6 mice by sulforaphane. *Immunopharmacol Immunotoxicol* 29(2): 173-186, 2007.
20. Rudolf K, Cervinka M, Rudolf E: Sulforaphane-induced apoptosis involves p53 and p38 in melanoma cells. *Apoptosis* 19(4): 734-747, 2014.
21. Huang SH, Hsu MH, Hsu SC, Yang JS, Huang WW, Huang AC, Hsiao YP, Yu CC and Chung JG: Phenethyl isothiocyanate triggers apoptosis in human malignant melanoma A375.S2 cells through reactive oxygen species and the mitochondria-dependent pathways. *Hum Exp Toxicol* 33(3): 270-283, 2014.
22. Huang SH, Wu LW, Huang AC, Yu CC, Lien JC, Huang YP, Yang JS, Yang JH, Hsiao YP, Wood WG, Yu CS and Chung JG: Benzyl isothiocyanate (BITC) induces G2/M phase arrest and apoptosis in human melanoma A375.S2 cells through reactive oxygen species (ROS) and both mitochondria-dependent and death receptor-mediated multiple signalling pathways. *J Agric Food Chem* 60(2): 665-675, 2012.
23. Thejass P and Kuttan G: Allyl isothiocyanate (AITC) and phenyl isothiocyanate (PITC) inhibit tumour-specific angiogenesis by down regulating nitric oxide (NO) and tumour necrosis factor- α (TNF- α) production. *Nitric Oxide* 16(2): 247-257, 2007.
24. Higdon JV, Delage B, Williams DE and Dashwood RH: Cruciferous vegetables and human cancer risk: epidemiologic evidence and mechanistic basis. *Pharmacol Res* 55(3): 224-236, 2007.
25. Bommareddy A, Hahm ER, Xiao D, Powolny AA, Fisher AL, Jiang Y and Singh VS: Atg5 Regulates Phenethyl Isothiocyanate-Induced Autophagic and Apoptotic Cell Death in Human Prostate Cancer Cells. *Cancer Res* 69(8): 3704-3712, 2009.
26. Herman-Antosiewicz A, Johnson DE and Singh SV: Sulforaphane Causes Autophagy to Inhibit Release of Cytochrome c and Apoptosis in Human Prostate Cancer Cells. *Cancer Res* 66(11): 5828-5835, 2006.

27. Xiao D, Bommareddy A, Kim SH, Sehrawat A, Hahm ER and Singh SV: Benzyl isothiocyanate causes FoxO1-mediated autophagic death in human breast cancer cells. *PLoS One* 7(3): e32597, 2012.
28. Xiao D, Powolny AA, Moura MB, Kelley EE, Bommareddy A, Kim SH, Hahm ER, Normolle D, Van Houten B and Singh SV: Phenethyl isothiocyanate inhibits oxidative phosphorylation to trigger reactive oxygen species-mediated death of human prostate cancer cells. *J Biol Chem* 285(34): 26558-26569, 2010.
29. Sakamaki K, Imai K, Tomii K and Miller DJ: Evolutionary analyses of caspase-8 and its paralogs. Deep origins of the apoptotic signalling pathways. *Bioessays* 37(7): 767-776, 2015.
30. Kim B, Srivastava SK and Kim SH: Caspase-9 as a therapeutic target for treating cancer. *Expert Opin Ther Targets* 19(1): 113-127, 2015.
31. Di Fazio P, Ocker M and Montalbano R: New drugs, old fashioned ways: ER stress induced cell death. *Curr Pharm Biotechnol* 13(11): 2228-2234, 2012.

Figure Legends

Figure 1. The effect of different concentrations of ITCs in A375 and HaCaT cells after 24 and 48h of treatment. Cells were subjected at a single bolus concentration ranging between 5-100 μ M of sulforaphane (**A**; A375 cells and **B**; HaCaT cells), phenethyl isothiocyanate (**C**; A375 cells and **D**; HaCaT cells) and benzyl isothiocyanate (**E**; A375 cells and **F**; HaCaT cells) for 24 and 48h. Data are representative of two independent experiments. An asterisk (*) or hashtag (#) indicates statistical significance when compared to their respective controls (DMSO) for 24 or 48h ($P < 0.05$).

Figure 2. The effect of different concentrations of ITCs on HaCaT, A431 and A375 cells after 48h of treatment. In general, cells were subjected to 0.1-5 μ M of sulforaphane (**A**), phenethyl isothiocyanate (**B**) and benzyl isothiocyanate (**C**) at 0.1-5 μ M for 48h with all treatments being refreshed after 24h. Data are representative from two independent experiments. An asterisk (*), hashtag (#) or diamond (\diamond) indicates statistical significance for HaCaT, A431 or

A375 treated cells, respectively when compared to their corresponding controls (DMSO) ($P<0.05$).

Figure 3. The effect of ITCs on enzymatic activity and protein expression levels of various caspases. A375 cells were subjected to 5 μ M of ITC (for 48h with all treatments being refreshed after 24h) and enzymatic activity levels were recorded for caspase 4 (**A**), caspase 8 (**B**) and caspase 9 (**C**). Additionally, under the same treatment conditions, protein expression levels were assessed for procaspase-3 (**D**), procaspase-6 (**E**) and procaspase-7 (**F**). Data are representative from two independent experiments. An asterisk (*) indicates statistical significance when compared to their respective controls ($P<0.05$)