

Northumbria Research Link

Citation: Goodall, Stuart (2018) The acclimatised spinal cord. *Journal of Physiology*, 596 (15). pp. 2949-2950. ISSN 0022-3751

Published by: Wiley-Blackwell

URL: <https://doi.org/10.1113/JP275552> <<https://doi.org/10.1113/JP275552>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/32773/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

1 **The acclimatised spinal cord**

2

3 Stuart Goodall

4

5 Department of Sport, Exercise, & Rehabilitation, Faculty of Health and Life Sciences, Northumbria
6 University, Newcastle, UK.

7

8

9 *A Perspective* written upon: **JP-RP-2017-274872R1** 'UBC-Nepal Expedition: Acclimatization to high-
10 altitude increases spinal motoneurone excitability during fatigue in humans' by Luca Ruggiero,
11 Alexandra F Yacyshyn, Jane Nettleton, and Chris J McNeil.

12

13

14 **Word count:** 894.

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32 **Address for correspondence:**

33 Stuart Goodall, PhD

34 Faculty of Health and Life Sciences

35 Northumbria University

36 Newcastle-upon-Tyne

37 NE1 8ST

38 UK

39

40 Tel: +44 191 227 4749

41 Fax: +44 191 227 4713

42 Email: stuart.goodall@northumbria.ac.uk

The acclimatised spinal cord

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

Fatigue is a universal and daily phenomenon that involves a myriad of complex mechanisms ultimately characterised as an exercise-induced decrease in the maximal force produced by a muscle (Gandevia *et al.*, 1996). The investigation of fatigue dates back to the early 19th century with the work from Alessandro Mosso and colleagues, however, today there is still debate among the scientific community about the appropriateness of defining and measuring ‘fatigue’ for different populations and exercise tasks (Enoka & Duchateau, 2016). A significant amount of the decline in maximal force occurs through processes which exist in the muscle, yet, changes in the performance of any, or all sites from the motor cortex to the muscle fibre, may contribute to fatigue. Thus, the impairment in muscle performance is not necessarily the limiting factor in force production from a fatigued muscle (Gandevia *et al.*, 1996). Over the last 70 years, stimulation techniques have been utilised to enable the measurement of fatigue and subsequently, sites of fatigue have been compartmentalised into those that exist within the muscle and the central nervous system (CNS). More specifically, over the last 20 years, the use of transcranial magnetic stimulation (TMS) has enabled greater investigation into the fatigued CNS and in line with the use of motor nerve stimulation, TMS enables excitability of the brain to muscle pathway to be quantified. As such, this technique has become a popular tool to monitor changes in CNS function in many paradigms within exercise science.

The sites implicated in the fatigue response are augmented when exercise is performed under conditions of environmental stress. The ability to perform exercise, whether it be an isolated limb activity or whole body manoeuvre, deteriorates in line with increasing altitude. At severe levels of altitude (arterial oxy-haemoglobin saturation <75%), the site of fatigue is predominantly located within the CNS and TMS has been used to demonstrate this in a number of investigations. Following a prolonged stay at altitude the level of CNS fatigue is attenuated, along with an associated increase in excitability of the brain to muscle pathway (Goodall *et al.*, 2014). However, one aspect that was not determined in the Goodall *et al.* (2014) investigation, and moreover a parameter that is infrequently studied during the investigation of fatigue, is how the change in spinal excitability contributes to the overall change in excitability of the corticospinal tract. Responsiveness of the motoneurone pool can be studied using cervicomedullary junction stimulation; the size of a resultant evoked potential (CMEP) provides information on excitability of the spinal tract. In this issue of *The Journal of Physiology*, a study has focused on the changes in motoneurone excitability and fatigue that are observed during exercise in acute hypoxia, and following a period of chronic hypoxia (Ruggiero *et al.*, 2017). Confirming previous findings, Ruggiero and colleagues (2017) found that the acute hypoxic environment elicited

1 the greatest amount of fatigue and specifically, this was due in part, to a sub optimal output from the
2 motor cortex. Supraspinal fatigue developed earlier in acute hypoxia (8 mins), compared to normoxia
3 and chronic hypoxia (16 mins). Furthermore, in acute hypoxia and normoxia, there was a fatigue
4 related reduction in motoneurone excitability, however, in chronic hypoxia responsiveness of the
5 motoneurone pool was not depressed and the CMEPs were larger than what was observed in the
6 acute condition (Ruggiero *et al.*, 2017).

7 The findings from Ruggiero *et al.* (2017) help to explain the results of Goodall *et al.* (2014) who
8 found a two fold increase in corticospinal excitability after a period of chronic hypoxia, however, those
9 authors concluded such a change was due to adaptations in spinal and/or supraspinal sites. It is now
10 clear that the heightened corticospinal excitability observed previously (Goodall *et al.*, 2014), was in
11 part, due to an acclimatised motoneurone pool (Ruggiero *et al.*, 2017). The timing of the
12 measurements made during chronic hypoxia in both of the aforementioned investigations (between
13 6 and 18 days), correspond with the window for maximal sympathetic norepinephrine concentration
14 following exposure to high altitude (Barnholt *et al.*, 2006). Ruggiero and colleagues (2017) explain that
15 norepinephrine acts as a potent neuromodulator, acting to increase motoneurone excitability. Such
16 an increase seems a logical mechanism for the manipulation of motoneurone responsiveness in
17 chronic hypoxia. Notably, given that output from the motoneurone pool is imperative for muscle
18 contraction, it follows that high altitude adaptation of intrinsic motoneurone properties can explain,
19 in part, the superior exercise performance commonly observed following a period of high altitude
20 acclimatisation.

21 Ruggiero and colleagues (2017) should be congratulated on the execution of this excellent
22 study which is the first to demonstrate a direct effect of acclimatisation within the spinal cord. From
23 a personal perspective, I know that venturing to altitude in order to undertake such experiment is not
24 without its challenges and complexities. Most investigations reporting the mechanisms of
25 acclimatisation tend to focus upon the cardiovascular and haematological adaptations which are of
26 course of great importance, but it is now clear that widespread changes are apparent throughout the
27 neuromuscular system. As a logical next step, it is worthwhile to investigate the change in intrinsic
28 motoneurone properties of locomotor muscles and understand the potential impact of this on the
29 common improvement in exercise performance observed following acclimatisation. However, most
30 notably for the field of high altitude physiology is that we are beginning to fully understand the
31 integrated adaptation which occurs throughout the human body during acclimatisation.

32

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15

References

Barnholt KE, Hoffman AR, Rock PB, Muza SR, Fulco CS, Braun B, Holloway L, Mazzeo RS, Cymerman A & Friedlander AL. (2006). Endocrine responses to acute and chronic high-altitude exposure (4,300 meters): modulating effects of caloric restriction. *American Journal of Physiology Endocrinology & Metabolism* **290**, E1078-1088.

Enoka RM & Duchateau J. (2016). Translating Fatigue to Human Performance. *Medicine & Science in Sports & Exercise* **48**, 2228-2238.

Gandevia SC, Allen GM, Butler JE & Taylor JL. (1996). Supraspinal factors in human muscle fatigue: evidence for suboptimal output from the motor cortex. *Journal of Physiology* **490**, 529-536.

Goodall S, Twomey R, Amann M, Ross EZ, Lovering AT, Romer LM, Subudhi AW & Roach RC. (2014). AltitudeOmics: exercise-induced supraspinal fatigue is attenuated in healthy humans after acclimatization to high altitude. *Acta Physiologica* **210**, 875-888.

Ruggiero L, Yacyshyn AF, Nettleton J & McNeil CJ. (2017). UBC-Nepal expedition: acclimatization to high-altitude increases spinal motoneurone excitability during fatigue in humans. *Journal of Physiology*.