

Northumbria Research Link

Citation: Nouioui, Imen, Carro, Lorena, Sangal, Vartul, Jando, Marlen, Igual, José Mariano, Goodfellow, Michael and Klenk, Hans-Peter (2018) Formal description of *Mycobacterium neglectum* sp. nov. and *Mycobacterium palauense* sp. nov., rapidly growing actinobacteria. *Antonie van Leeuwenhoek*, 111 (7). pp. 1209-1223. ISSN 0003-6072

Published by: Springer

URL: <https://doi.org/10.1007/s10482-018-1029-5> <<https://doi.org/10.1007/s10482-018-1029-5>>

This version was downloaded from Northumbria Research Link: <http://nrl.northumbria.ac.uk/33322/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



UniversityLibrary



Northumbria
University
NEWCASTLE

1 **Formal description of *Mycobacterium neglectum* sp. nov. and *Mycobacterium palauense***
2 **sp. nov., rapidly growing actinobacteria**

3

4 Imen Nouioui^{1*}, Lorena Carro¹, Vartul Sangal², Marlen Jando³, José Mariano Igual⁴, Michael
5 Goodfellow¹, Hans-Peter Klenk¹

6

7 ¹School of Natural and Environmental Sciences, Ridley Building, Newcastle University,
8 Newcastle upon Tyne, NE1 7RU, United Kingdom.

9 ²Faculty of Health and Life Sciences, Northumbria University, Newcastle upon Tyne NE1 8ST,
10 UK

11 ³Leibniz Institute DSMZ – German Collection of Microorganisms and Cell Cultures,
12 Inhoffenstraße 7B, 38124 Braunschweig, Germany

13 ⁴Instituto de Recursos Naturales y Agrobiología de Salamanca, Consejo Superior de
14 Investigaciones Científicas (IRNASA-CSIC), c/Cordel de Merinas 40-52, 37008 Salamanca,
15 Spain

16

17 To whom correspondence should be addressed: imen.nouioui@ncl.ac.uk

18

19 **Abstract**

20 The taxonomic positions of two fast growing mycobacteria (CECT 8778^T and CECT 8779^T)
21 were established using a polyphasic approach. The strains were shown to have
22 chemotaxonomic, cultural and morphological properties consistent with their classification in
23 the genus *Mycobacterium*. Multi-locus sequence analyses (MLSA) show that strain CECT
24 8778^T forms a well-supported clade together with the type strains of *Mycobacterium aurum*,
25 *Mycobacterium austroafricanum* and *Mycobacterium vanbaalenii* while strain CECT 8779^T
26 presents as a distinct branch that is well separated from its nearest phylogenetic neighbours;
27 it is also apparent from the MLSA genetic distances that these strains are most closely related
28 to the type strains of *Mycobacterium mageritense* and *M. vanbaalenii*, respectively. Digital
29 DNA:DNA hybridization and average nucleotide identity values between each of the strains
30 and its close phylogenetic neighbour are below the 70 and 96% threshold values for definition
31 of prokaryotic species; these results are underpinned by corresponding phenotypic data. Based
32 upon the consensus of the phenotypic and phylogenetic analyses, it can be concluded that the
33 two strains represent novel species within the genus *Mycobacterium* for which the following
34 names are proposed: *Mycobacterium neglectum* sp. nov., with the type strain CECT 8778^T (BN
35 3150^T = DSM 44756^T) and *Mycobacterium palauense* sp. nov., with the type strain CECT
36 8779^T (= DSM 44914^T).

37

38 **Keywords:** Actinobacteria, phenotyping, phylogeny, polyphasic taxonomy

39

40 **Introduction**

41 The genus *Mycobacterium* (Lehmann and Neumann 1896), the sole representative of the
42 family *Mycobacteriaceae* (Chester 1897) can be distinguished from all of the other genera
43 classified in the order *Corynebacteriales* by using a selection of genotypic and phenotypic
44 methods (Goodfellow and Jones 2012). The genus encompasses pathogenic and non-
45 tuberculous mycobacteria (Magee and Ward 2012; Forbes 2017; Gcebe et al. 2017) which can
46 be assigned to two groups based on growth rates. Slowly growing strains require 7 or more
47 days of incubation at optimal temperature to produce visible colonies from highly diluted
48 inocula whereas those of rapidly growing strains are evident in fewer than 7 days under
49 comparable conditions (Wayne and Kubica 1986). Polyphasic taxonomic procedures are now
50 used to detect novel mycobacterial species, as exemplified by the delineation of species
51 previously aggregated within the *Mycobacterium abscessus* and *Mycobacterium avium*

52 complexes (Ben Salah et al. 2009; Tortoli et al. 2016). Developments such as these are needed
53 to detect the causal agents of mycobacterial infections and to establish the primary reservoirs
54 of individual mycobacterial species within natural habitats (Tran and Dahl 2016; Shahraki et
55 al. 2017).

56

57 Environmental mycobacteria are common in aquatic and terrestrial ecosystems
58 (Nishiuchi et al. 2017; Roguet et al. 2016), including biofilms of water distribution systems
59 (September et al. 2004; Feazel et al. 2009; Gomez-Smith et al. 2015). This study was
60 undertaken to establish the taxonomic status of two rapidly growing mycobacteria: strain
61 CECT 8778^T (DSM 44756^T) was isolated from a biofilm of a water distribution system and
62 strain CECT 8779^T from marine sediment. A 16S rRNA gene sequence of strain DSM 44756
63 (then coded BN 3150) was deposited in GenBank (accession number AJ580802) under the
64 name “*Mycobacterium neglectum*”. This code and species epithet have been used quite
65 extensively in the literature (Thomas et al. 2008; Hussein et al. 2009; Jenkins et al. 2009; Salah
66 et al. 2009; Loret and Creub 2010; Pontiroli et al. 2013; Nishiuchi et al. 2017). However, at no
67 stage has a formal description been given for “*M. neglectum*” hence this name has no standing
68 in nomenclature (Rule 29 of International Code of Nomenclature of Prokaryotes [2008
69 revision]; Parker et al. 2015). In the present polyphasic study, we provide the first formal
70 description of *Mycobacterium neglectum* sp. nov., with the type strain CECT 8778^T, while a
71 second novel species represented by strain CECT 8779^T is named *Mycobacterium palauense*
72 sp. nov.

73

74 **Materials and methods**

75

76 Source, maintenance and cultivation of strains

77

78 Strain CECT 8778^T, isolated from a biofilm of an underground drinking water system in
79 Duisburg, Germany in 1999, and strain CECT 8779^T, isolated from a marine sediment collected
80 from the Republic of Palau in 2004, were obtained from the Spanish Type Culture Collection.
81 The strains, together with *Mycobacterium aurum* DSM 43999^T (Tsukamura 1966),
82 *Mycobacterium austroafricanum* DSM 44191^T (Tsukamura et al. 1983), *Mycobacterium*
83 *mageritense* DSM 44476^T (Domenech et al. 1997) and *Mycobacterium vanbaalenii* DSM 7152^T
84 (Khan et al. 2002) were maintained as suspensions in 35% (v/v) glycerol at -80°C. Biomass for

85 the chemotaxonomic and molecular systematic studies on the isolates was cultured in shake
86 flasks (200 revolutions per minute) of proteose peptone-meat extract-glycerol agar medium
87 (PMG; DSMZ medium 250); after incubation at 28°C for 5 days, cells were harvested and
88 washed three times in sodium chloride solution (0.9%, w/v). Cells for the chemotaxonomic
89 analyses were freeze dried and stored at room temperature; wet biomass for the fatty acid
90 analyses was prepared under the same conditions.

91

92 Phylogeny

93 Genomic DNA was extracted from strains CECT 8778^T and CECT 8779^T using the procedure
94 described by Amaro et al. (2008). The genomes of the strains were sequenced using an MiSeq
95 instrument (Illumina), as described by Sangal et al. (2015) and assembled into contigs using
96 SPAdes 3.9.0 with a kmer length of 127 (Bankevich et al. 2012). Annotation of the genomes
97 was achieved using the RAST pipeline available on the RAST server (Aziz et al. 2008, 2012).
98 Complete 16S rRNA gene sequences of strains CECT 8778^T and CECT 8779^T were extracted
99 from the draft genomes (accession numbers NVQE000000000 and NVQF000000000,
100 respectively) and deposited in GenBank under accession numbers MF769621 and MF769712.
101 Corresponding 16S rRNA gene sequences of the type strains of closely related *Mycobacterium*
102 spp. were retrieved from the EzBioCloud server (Yoon et al. 2017) and pairwise sequence
103 similarities calculated using the Genome-to-Genome Distance Calculator (GGDC) web server
104 (Meier-Kolthoff et al. 2013a, b). Phylogenies derived from the 16S rRNA gene sequences were
105 inferred using the GGDC web server adapted to single genes (Meier-Kolthoff et al. 2014).
106 Multiple sequence alignments were generated using MUSCLE software (Edgar 2004) and a
107 maximum-likelihood (ML) tree inferred from the alignment with RAxML (Stamatakis 2014)
108 using rapid bootstrapping together with the auto Maximal-Relative-Error (MRE) criterion
109 (Pattengale et al. 2010). Similarly, a maximum-parsimony (MP) tree was inferred from the
110 alignments with the ‘Tree analysis New Technology’ (TNT) program (Goloboff et al. 2008)
111 using 1000 bootstraps together with tree bisection and reconnection branch swapping and ten
112 random sequence replicates. The sequences were checked for computational bias using the X²
113 test implemented in PAUP* (Phylogenetic Analysis Using Parsimony) (Swofford 2002).

114 Partial sequences of three housekeeping genes, *hsp65* (heat shock protein), *rpoB* (RNA
115 polymerase beta subunit) and *recA* (recombination protein A) (McNabb et al. 2004;
116 Ramaprasad et al. 2016), were drawn from the draft genomes of strains CECT 8778^T and CECT
117 8779^T and deposited in GenBank under the accession numbers MF774022, MF774023,

118 MF774024 , MF77402 , MF774026, MF774027, respectively. A multilocus sequence
119 analysis (MLSA) tree was generated from 3203 nucleotides (nt) of concatenated sequences of
120 the three housekeeping genes and corresponding 16S rRNA gene sequences and ML and MP
121 trees inferred as described above. In addition, a neighbour-joining (NJ) tree (Saitou and Nei
122 1987) was generated from the MEGA 7 software package (Kumar et al. 2015). The alignment
123 of the concatenated sequences and the corresponding evolutionary distances were carried out
124 using CLUSTAL W software (Thompson et al. 1997) and the Kimura two parameter model
125 (Kimura 1980), respectively.

126 The average nucleotide identity (ANI) between strain CECT 8778^T and *M. aurum* DSM
127 43999^T (genome accession number is NZ_CVQQ000000000), *M. austroafricanum* DSM
128 44191^T genome accession number is (NZ_HG964469) and *M. vanbaalenii* PYR1^T (genome
129 accession number is CP000511) and between strain CECT 8779^T and *M. mageritense* DSM
130 44476^T (genome accession number is NZ_CCBF000000000), their respective near
131 phylogenetic neighbours, were calculated according to Rodriguez and Konstantinidis (2014).
132 Similarly, digital DNA-DNA hybridization (dDDH) similarities were determined between the
133 two strains and their close phylogenetic neighbours using the GGDC server (Meier-Kolthoff et
134 al. 2013a).

135

136 Chemotaxonomy

137 The chemotaxonomic profiles of strains CECT 8778^T and CECT 8779^T and their respective
138 close phylogenetic neighbours were determined using standard thin-layer chromatographic
139 procedures. To this end, the strains were examined for diaminopimelic acid isomers (A₂pm)
140 (Staneck and Roberts 1974), predominant menaquinones (Collins 1985), mycolic acids
141 (Mininkin et al. 1980), diagnostic sugars (Lechevalier and Lechevalier 1970) and polar lipids
142 (Mininkin et al. 1984). In addition, cellular fatty acids were extracted from freeze dried biomass
143 of the strains and fatty acid methyl esters (FAMES) prepared following saponification and
144 methylation using the procedure introduced by Miller (1982), as modified by Kuykendall et al.
145 (1988). The FAMES were analysed by gas chromatography (Agilent 6890 instrument) and the
146 resultant peaks automatically integrated. The identities of the fatty acids were determined using
147 the standard Microbial Identification (MIDI) System, version 4.5, and the Myco 6 database
148 (Sasser 1990).

149

150 Growth and cultural properties

151 Strains CECT 8778^T and CECT 8779^T were examined for their ability to grow and form
152 colonies and pigments on glucose-yeast extract-malt extract agar (GYM, DSMZ medium 65),
153 Löwenstein-Jensen (LJ) medium (Jensen 1932), Middlebrook 7H10 agar supplemented with
154 oleic acid, albumin dextrose and catalase (MB7H10, Lorian 1968), proteose peptone-meat
155 extract-glycerol agar (PMG; DSMZ medium 250) and tryptic soy agar (TSA, MacFaddin 1985)
156 for 14 days at 4, 10, 20, 25, 28 and 42°C under light and dark conditions. The strains were also
157 examined for acid-alcohol-fastness using the Ziehl-Neelsen method (Runyon et al. 1980) and
158 for their ability to grow on PMG agar under anaerobic conditions at 28°C using an anaerobic
159 bag system (Sigma-Aldrich 68061).

160 Phenotypic tests

161 The two strains and their close phylogenetic neighbours were examined for phenotypic tests
162 found to be useful in mycobacterial systematics (Magee and Ward 2012, Nouioui et al. 2017).
163 The strains were tested for their ability to use sole carbon and sole nitrogen sources, to grow in
164 the presence of several concentrations of sodium chloride, at a range of pH values and in the
165 presence of antibiotics using GENIII microplates and an Omnilog device (BIOLOG, Hayward,
166 CA). The tests were carried out in duplicate using freshly prepared inocula (OD₆₀₀-0.3-0.6)
167 harvested from the mid-logarithmic growth phase of PMG agar plates incubated at 28°C for 7
168 days. The resultant data were exported and analysed using the opm package version 1.3.36
169 (Vaas et al. 2012, 2013). The strains were also examined for their ability to produce
170 arylsulfatase after 3 and 14 days (Tomioka et al. 1990), catalase (Palomino et al. 2007) and
171 heat stable catalase (Sequeira de Latini and Barrera 2008) and for niacin accumulation (Kent
172 and Kubica 1985), resistance to potassium tellurite (Kent and Kubica 1985; Kilburn et al. 1969),
173 degradation of Tween 80 (Ribón 2012) and urea hydrolysis (Palomino et al. 2007) using the
174 media and incubation conditions described in these references. All of these tests were carried
175 out in duplicate using the standard inoculum.

176

177 **Results and discussion**

178 The chemotaxonomic, growth and staining properties of strains CECT 8778^T and CECT 8779^T
179 were shown to be consistent with their classification in the genus *Mycobacterium* (Magee and
180 Ward 2012). The organisms were found to be strictly aerobic, Gram-positive, acid-alcohol fast,
181 rapid growing, rod-shaped bacteria which contain *meso*-diaminopimelic acid, arabinose,
182 galactose, glucose, rhamnose and ribose in whole organism hydrolysates (wall chemotype IV
183 *sensu* Lechevalier and Lechevalier 1970); mixtures of saturated, unsaturated and 10-methyl

184 octadecanoic (tuberculostearic) fatty acids; mycolic acids; dihydrogenated menaquinones with
185 nine isoprene units (MK9(H₂)) as the predominant isoprenologue; and a polar lipid profile that
186 includes diphosphatidylglycerol, phosphatidylethanolamine (diagnostic phospholipid),
187 phosphatidylinositol, as well as a glycopospholipid and glycolipids (phospholipid type II;
188 Lechevalier et al. 1977). Both strains were found to produce unpigmented colonies under both
189 light and dark conditions on LJ, MB7H10, PMG and TSA plates after 5 days at 28°C; moderate
190 growth was observed at 20°C, 25°C and 37°C; optimal growth was detected at 28°C on GYM,
191 MB7H10 and PMG agar after 5 days. The strains were unable to grow on any of these media
192 at 4°C, 15°C, 42°C, or 45°C or under anaerobic conditions at 28°C on PMG agar. Strain CECT
193 8778^T and its nearest phylogenetic neighbours, *M. aurum* DSM 43999^T, *M. austroafricanum*
194 DSM 44191^T and *M. vanbaalenii* DSM 7152^T, share several features; they are all acid acid-
195 alcohol fast, rapid growing bacteria that grew on MB7H10 and PMG media at 28°C though
196 only the test strain formed non-pigmented colonies. Strain CECT 8779^T and *M. mageritense*
197 DSM 44476^T were found to have very similar cultural and morphological traits though only
198 the latter grew at 22, 30, 37 and 45°C on LJ media (Domenech, et al. 1997).

199 The pairwise 16S rRNA gene similarities between strain CECT 8778^T and *M. aurum*
200 NCTC 10437^T, *M. austroafricanum* DSM 44191^T, *Mycobacterium pyrenivorans* DSM 44605^T
201 (Derz et al. 2004), *Mycobacterium vaccae* ATCC 25954^T (Bönicke and Juhasz 1964) and *M.*
202 *vanbaalenii* PYR-1 were found to be 99.2%, 99.3%, 98.6%, 98.9%, 99.2%, respectively. It can
203 be seen from Figure 1 that the two strains were found to be well separated from the type strains
204 of the remaining fast growing *Mycobacterium* species. Strain CECT 8778^T was shown to form
205 a distinct branch at the periphery of a well-supported subclade that included all of the organisms
206 cited above. In turn, the pairwise 16S rRNA gene similarities between strain CECT 8779^T and
207 the type strains of *M. mageritense*, *Mycobacterium peregrinum* ATCC 14467^T (Kusunoki and
208 Ezaki 1992) and *Mycobacterium wollinskyi* ATCC 700010^T (Brown et al. 1999) were found to
209 be 98.8%, 98.8% and 98.9%, respectively; strain CECT 8779^T formed a well defined branch in
210 a subclade that also contained these organisms, albeit one that was supported by a high
211 bootstrap value only in the ML analysis. It can also be seen from Figure 1 that all of the rapidly
212 growing strains were sharply separated from the type strain of *Mycobacterium Tuberculosis*
213 (Zopf 1883; Lehmann and Neumann 1896), the type species of the genus.

214 The MLSA trees based on the sequences of the three housekeeping genes and the
215 corresponding 16S rRNA sequences are shown in Figure 2; this tree was inferred from 3257
216 nt, 604 of which were variable and 324 of which were parsimony-informative. The average
217 bootstrap supports for the ML and MP trees were found to be 90.0% and 92.3%, respectively.

218 It is evident from Figure 2 that strain CECT 8778^T forms a well supported subclade together
219 with the type strains of *M. aurum*, *M. austroafricanum* and *M. vanbaalenii* while strain CECT
220 8779^T forms a branch distinct from all of the other mycobacteria, including the type strains of
221 *M. mageritense*, *M. peregrinum* and *M. wollinskyi*. Moreover, it is clear from Table 1 that
222 strains CECT 8778^T and CECT 8779^T share close genetic distances, namely 0.02 and 0.03, with
223 the type strains of *M. vanbaalenii* and *M. mageritense*, respectively. However, in the
224 corresponding NJ tree, strain CECT 8778^T was shown to be more closely related to the *M.*
225 *austroafricanum* and *M. vanbaalenii* type strains than to the *M. aurum* strain though these
226 relationships were not supported by high bootstrap values (Fig. S1). The phylogenetic trees
227 based on the sequences of the individual housekeeping genes are shown in Figures S2-S4; the
228 relationships between strain CECT 8778^T and the type strains of *M. aurum*, *M. austroafricanum*
229 and *M. vanbaalenii* are evident in all of the trees though bootstrap values are low. In contrast,
230 strain CECT 8779^T was found to form a distinct branch in all of these trees.

231 The genome sizes of strains CECT 8778^T (NVQE000000000) and CECT 8779^T
232 (NVQF000000000) were both found to be ~6.2Mb with average *in silico* G+C contents of
233 65.3mol% and 69.4mol%, respectively. The draft genome of strain CECT 8778^T was generated
234 from 123 contigs with N lengths of 116368 and was found to contain 6159 predicted protein
235 coding sequences and 51 tRNA genes. Similarly, the draft genome of strain CECT 8779^T was
236 compiled from 210 contigs with N lengths of 61684 and was shown to have 5802 predicted
237 protein coding sequences and 71 tRNA genes. The coverage for strains CECT 8778^T and CECT
238 8779^T were 79X and 67X, respectively. Strains CECT 8778^T and CECT 8779^T contain, very
239 similar subsystem gene functions as exemplified in Table S1.

240 Strain CECT 8778^T and *M. aurum* NCTC 10437^T, *M. austroafricanum* DSM 44191^T
241 and *M. vanbaalenii* PYR-1^T, currently its closest phylogenetic neighbours, were found to share
242 dDDH similarities of 20.1%, 21.1% and 21.1%, respectively, values well below the 70% cut
243 off point recommended for the delineation of prokaryotic species (Wayne et al. 1987). The
244 corresponding ANI similarities between strain CECT 8778^T and the three strains mentioned
245 above were found to be 78.4%, 78.7% and 79.6%, values well below the 95-96% threshold
246 used to distinguish between closely related species of prokaryotes (Goris et al. 2007; Richter
247 and Rosselló-Móra 2009; Chun and Rainey 2014). Similarly, the dDDH and ANI values
248 between strain CECT 8779^T and *M. mageritense* DSM 44476^T, its current closest phylogenetic
249 neighbour, were 20.9% and 79.1%, respectively, values well below the species thresholds cited
250 above.

251 Identical results were recorded for all of the phenotypic tests that were carried out in
252 duplicate. It can be seen from Table 2 that strains CECT 8778^T and CECT 8779^T can be
253 distinguished from one another and from their respective reference strains using a combination
254 of phenotypic properties though it is apparent that all of these organisms share a common set
255 of features. Strain CECT 8778^T, unlike the type strains of *M. aurum*, *M. austroafricanum* and
256 *M. vanbaalenii*, was shown to grow in the presence of tetrazolium blue and tetrazolium violet
257 and also differed from these strains by its inability to utilise L-alanine, butyric acid, α -hydroxy-
258 butyric acid, α -keto-butyric acid, galactose, glycerol, L-lactic acid, pectin and sucrose or to
259 grow in the presence of lithium chloride, nalidixic acid and sodium chloride (up to 8%, w/v).
260 Similarly, strain CECT 8779^T, unlike *M. mageritense* DSM 44476^T, was shown to use D-
261 arabitol, D-glucose-phosphate, D-maltose, D-mannose, methyl pyruvate, pectin, D-sorbitol,
262 sucrose, D-trehalose and D-turanose as sole carbon sources and to grow in the presence of
263 lithium chloride and sodium chloride (8%, w/v). In turn, strain CECT 8778^T, unlike strain
264 CECT 8779^T, was seen to metabolise dextrin, *myo*-inositol, D- and L-malic acid, quinic acid,
265 D- saccharic acid, D- salicin, L-serine and bromo-succinic acid. Conversely, strain CECT
266 8779^T was found to use L-alanine, L-aspartic acid, butyric acid, citric acid, D-glucose-6-
267 phosphate, glycerol, α -keto-glutaric acid, L-histidine, L-lactic acid, D-maltose, N-acetyl- β -D-
268 mannosamine, methyl pyruvate, pectin, L-rhamnose and sucrose and to grow in the presence
269 of lithium chloride, nalidixic acid and sodium chloride.

270 The kind of mycolic acids synthesised by representatives of *Mycobacterium* species fall
271 into several well established patterns of taxonomic value (Mininkin et al. 1985; Magee and
272 Ward 2012). In the present study, the test strains were found to have different mycolic acid
273 profiles: strain CECT 8778^T was shown to contain α - and *decarboxy*- mycolates and strain
274 CECT 8779^T α -, *keto*- and *methoxy*- mycolates. This latter pattern serves to distinguish strain
275 CECT 8779^T from *M. mageritense* DSM 44476^T which is characterised by the presence of α -,
276 α' - and *epoxy*-mycolates (Domenech et al. 1997). Similarly, the mycolic acid profile
277 distinguishes strain CECT 8778^T from *M. aurum* DSM 43999^T and *M. austroafricanum* DSM
278 44191^T as these strains have α - and *keto*-mycolates and wax esters (Mininkin et al. 1985; Magee
279 and Ward 2012). Similarly, complex polar lipid pattern of strain CECT 8778^T serves to
280 distinguish it from both strain CECT 8779^T and from the type strains of *M. aurum*, *M.*
281 *austroafricanum* and *M. vanbaalenii*; all of these strains were found to contain
282 diposphatidylglycerol, phosphatidylethanolamine (diagnostic lipid), phosphatidylinositol and

283 glycophospholipids. Strain CECT 8779^T, unlike its near phylogenetic neighbour, *M.*
284 *mageritense* DSM 44476^T, was shown to have a lipid pattern that lacked phosphatidylglycerol.

285 All of the strains produced complex mixtures of straight-chain saturated, unsaturated
286 and 10-methyl-octadecanoic (tuberculostearic) fatty acids, a profile typical of members of the
287 genus *Mycobacterium* (Magee and Ward 2012). With few exceptions, strains CECT 8778^T, *M.*
288 *aurum* DSM 43999^T, *M. austroafricanum* DSM 44191^T and *M. vanbaalenii* DSM 7152^T were
289 found to have major proportions (>10% of total fatty acid) of C_{16:0} (13.3-72.6%), C_{18:1} ω9c
290 (7.6-19.7.0%) and summed features 2 (12.6-43.2%) and 3 (12.3-16.8%) though the
291 predominant component varied (Table 3). The fatty acid profiles of CECT 8778^T and the type
292 strain of *M. vanbaalenii* were distinct; the test strain, for instance, produced higher proportions
293 of summed feature 2 (43.2 against 27.6%). Even greater differences were found between the
294 fatty acid profiles of CECT 8778^T and *M. aurum* DSM 43999^T; the latter, for instance, was
295 especially rich in C_{16:0} (25.4% against 13.3.0%). Marked differences were found between the
296 fatty acid profiles of strain CECT 8779^T and *M. mageritense* DSM 44476^T as the former
297 contained moderate proportions of C_{16:1} ω9c and summed features 2 and a lower amount of
298 C_{16:0} (25.1 against 40.0%) (Table 3).

299 In summary, strain CECT 8778^T can be distinguished readily from *M. aurum* DSM
300 43999^T, *M. austroafricanum* DSM 44191^T and *M. vanbaalenii* DSM 7251^T, its close
301 phylogenetic neighbours, in the 16S rRNA and MLSA gene trees, by low ANI and dDDH
302 scores and by a range of chemotaxonomic and phenotypic markers. A similar wealth of
303 taxonomic data separate strain CECT 8779^T from the type strain of *M. mageritense*, its close
304 phylogenetic neighbour. These datasets clearly show that strains CECT 8778^T and CECT 8779^T
305 represent new centres of taxonomic variation within the genus *Mycobacterium*; the names
306 chosen for these species are *Mycobacterium neglectum* sp. nov. and *Mycobacterium palauense*
307 sp. nov., respectively. The Digital Protologue database TaxoNumbers for strains CECT 8778^T
308 and CECT 8779^T are TA00318 and TA000312, respectively.

309 **Description of *Mycobacterium neglectum* sp. nov.**

310 *Mycobacterium neglectum* (neg.lec'tum. L. adj. *neglectum*, neglected reflecting the history of
311 the strain)

312 Strict aerobic, Gram-stain positive, acid-alcohol fast, rapid growing organism which
313 forms unpigmented colonies after growth on Middelbrook 7H10, proteose peptone-meat
314 extract-glycerol and LJ media after incubation under the light and dark conditions after 5 days

315 at 28°C. Grows between 25°C and 28°C, optimally at 28°C and at pH 7. Produces arylsulfatase
316 after 3 and 14 days, catalase, nitrate reductase, urease and grows in the presence of potassium
317 tellurite. Additional phenotypic data are given in the text and in Table 1. Whole cell
318 hydrolysates are rich in *meso*-diaminopimelic acid, arabinose, galactose, glucose, ribose and
319 rhamnose; the polar lipid profile contains diphosphatidylglycerol, glycopospholipids (GPL1-
320 2), phosphatidylethanolamine, phosphatidylinositol and a glycolipid; MK9 (H₂) is the
321 predominant menaquinone and the major fatty acid is C_{17:1} ω7c/18 alcohol. Contains α- and
322 *decarboxy*-mycolic acids. The DNA G+C content determined from the draft genome of the
323 type strain is 65.3 mol%.

324 The type strain, CECT 8778^T (= BN3150^T = DSM 44756^T) was isolated from a biofilm
325 of an underground drinking water system in Germany. The Genbank accession number of the
326 draft genome sequence of strain CECT 8778^T is NVQE000000000.

327 **Description of *Mycobacterium palauense* sp. nov.**

328 *Mycobacterium palauense* (pa.lau.en'se N.L. neut. adj. *palauense* referring to the Republic
329 Palau, the source of the strain)

330 Strict aerobic, Gram-stain positive, acid-alcohol fast, rapid growing organism which
331 forms unpigmented colonies after growth on Middelbrook 7H10, proteose peptone-meat
332 extract-glycerol and LJ media after incubation under the light and dark conditions for 5 days
333 at 28°C. Grows between 25°C and 28°C, optimally at 28°C and at pH 7 and in the presence of
334 up to 8% w/v NaCl. Produces arylsulfatase after 3 and 14 days, catalase, accumulates niacin,
335 degrades Tween 80 and grows in presence of potassium tellurite. Additional phenotypic data
336 are given in the text and in Table 1. Whole cell hydrolysates are rich in *meso*-diaminopimelic
337 acid, arabinose, galactose, glucose, ribose and rhamnose; the polar lipid profile contains
338 diphosphatidylglycerol, phosphatidylethanolamine, phosphatidylinositol, as well as
339 glycopospholipids (GPL₁₋₂) and a glycolipid; MK9 (H₂) is the predominant menaquinone and
340 the major fatty acids are C_{16:0} and C_{18:1} ω9c. Contains α, *keto-methoxy* mycolic acids. The DNA
341 G+C content determined from the draft genome of the type strain is 69.4 mol %.

342 The type strain, CECT 8779^T (= DSM 44914^T) was isolated from marine sediment from
343 the Republic of Palau. The Genbank accession number of the draft genome sequence of strain
344 CECT 8779^T is NVQF000000000.

345

346 **Acknowledgements** This project was supported by the School of Natural and Environmental
347 Sciences (Newcastle University). IN. and LC. are grateful to Newcastle University for
348 postdoctoral fellowships.

349 **Compliance with ethical standards**

350 **Conflict of interest** The authors declare that they have no conflicts of interest.

351 **Ethical statement** This article does not contain any studies inoculating human participants or
352 animals.

353

354 **References**

355

356 Amaro A, Duarte E, Amado A, Ferronha H, Botelho A (2008) Comparison of three DNA
357 extraction methods for *Mycobacterium bovis*, *Mycobacterium tuberculosis* and
358 *Mycobacterium avium* subsp. *avium*. Lett Appl Microbiol 47:8-11

359 Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K et al (2008) The
360 RAST Server: rapid annotations using subsystems technology. BMC Genomics 9:75

361 Aziz RK, Devoid S, Disz T, Edwards RA, Henry CS, Olsen GJ, Olson R et al (2012) SEED
362 servers: high-performance access to the SEED genomes, annotations, and metabolic
363 models. PLoS One 7:e48053

364 Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM et al
365 (2012) SPAdes: a new genome assembly algorithm and its applications to single-cell
366 sequencing. J Comput Biol 19:455-477

367 Ben Salah I, Cayrou C, Raoult D, Drancourt M. (2009) *Mycobacterium marseillense* sp. nov.,
368 *Mycobacterium timonense* sp. nov. and *Mycobacterium bouchedurhonense* sp. nov.,
369 members of the *Mycobacterium avium* complex. Int J Syst Evol Microbiol 59:2803-
370 2808

371 Bönicke R, Juhasz SE (1964) Beschreibung der neuen Species *Mycobacterium vaccae* n. sp.
372 Zentralb Bakteriol, Parasitenkd Infektionskr Hyg, Abt I Orig 192:133-135

373 Brown BA, Springer B, Steingrube VA, Wilson RW, Pfyffer GE, Garcia MJ, Menendez MC,
374 et al (1999) *Mycobacterium wolinskyi* sp. nov. and *Mycobacterium goodii* sp. nov., two
375 new rapidly growing species related to *Mycobacterium smegmatis* and associated with
376 human wound infections: A cooperative study from the International Working Group
377 on Mycobacterial Taxonomy. Int J Syst Bacteriol 49:1493-1511

378 Chester FD (1897) Report of the mycologist: bacteriological work. Del Agric Exp Sta Bull
379 9:38-145

380 Chun J, Rainey FA (2014) Integrating genomics into the taxonomy and systematics of the
381 *Bacteria* and *Archaea*. Int J Syst Evol Microbiol 64:316–324.

382 Collins MD (1985) Analysis of isoprenoid quinones. Meth Microbiol 18:329-366

383 Derz K, Klinner U, Schuphan I, Stackebrandt E, Kroppenstedt RM (2004) *Mycobacterium*
384 *pyrenivorans* sp. nov., a novel polycyclic-aromatic-hydrocarbon-degrading species. Int
385 J Syst Evol Microbiol 54:2313-2317

386 Domenech P, Jimenez MS, Menendez MC, Bull TJ, Samper S, Manrique A, Garcia MJ (1997)
387 *Mycobacterium mageritense* sp. nov. Int J Syst Bacteriol 47:535-540

388 Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high
389 throughput. Nucleic Acids Res 32:1792-1797

390 Feazel LM, Baumgartner LK, Peterson KL, Frank DN, Harris JK, Pace NR (2009)
391 Opportunistic pathogens enriched in showerhead biofilms. PNAS 106: 16393-16399.

392 Forbes BA (2017) Mycobacterial taxonomy. J Clin Microbiol 55: 380-383.

393 Gcebe N, Rutten V, van Pittius NG, Naicker B, Michel A (2017) *Mycobacterium*
394 *malmesburyense* sp. nov., a non-tuberculous species of the genus *Mycobacterium*
395 revealed by multiple gene sequence characterization. Int J Syst Evol Microbiol 67: 832-
396 838

397 Goodfellow M, Jones A (2012) Order V. *Corynebacteriales* ord. nov. In Goodfellow M,
398 Kämpfer P, Busse HJ, Trujillo ME, Suzuki KI, Ludwig W, Whitman WB (eds) Bergey's
399 Manual of Systematic Bacteriology, 2nd edn, vol 5, The *Actinobacteria*, Part A. Springer,
400 New York, pp 232-243

401 Goloboff PA, Farris JS, Nixon KC (2008) TNT, a free program for phylogenetic analysis.
402 Cladistics 24:774-786

403 Gomez-Smith CK, . LaPara TM, Hozalski RM (2015) Sulfate reducing bacteria and
404 mycobacteria dominate the biofilm communities in a chloraminated drinking water
405 distribution system. Environ Sci Technol. 49: 8432-8440.

406 Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM (2007)
407 DNA-DNA hybridization values and their relationship to whole-genome sequence
408 similarities. Int J Syst Evol Microbiol 57:81-91.

409 Hussein Z, Landt O, Wirths B, Wellinghausen N (2009) Detection of non-tuberculous
410 mycobacteria in hospital water by culture and molecular methods. Int J Med Microbiol
411 299: 281-290.

412 Jensen KA (1932) Reinzüchtung und Typenbestimmung von Tuberkelbazillenstämmen.
413 Zentralbl Bakteriologie I. Orig 125:222-239

414 Jenkins SN, Waite IS, Blackburn A, Husband R, Rushton SP, Manning DC, O'Donnell AG
415 (2009) Actinobacterial community dynamics in long term managed grasslands. Antonie
416 van Leeuwenhoek 95:319-334.

417 Kent PT, Kubica GPW (1985) Public Health Mycobacteriology A Guide for the Level III
418 Laboratory. U.S. Department. of Health and Human Services, Public Health Service,
419 Centers for Disease Control, Atlanta

420 Khan AA, Kim SJ, Paine DD, Cerniglia CE (2002) Classification of a polycyclic aromatic
421 hydrocarbon-metabolizing bacterium, *Mycobacterium* sp. strain PYR-1, as
422 *Mycobacterium vanbaalenii* sp. nov. Int J Syst Evol Microbiol 52:1997-2002

423 Kilburn JO, Silcox VA, Kubica GP (1969) Differential identification of mycobacteria. V. The
424 tellurite reduction test. Am Rev Resp Dis 99:94-100

425 Kimura M (1980) A simple method for estimating evolutionary rates of base substitutions
426 through comparative studies of nucleotide sequences. J Mol Evol 16:111-120

427 Kumar S, Stecher G, Tamura K (2015) MEGA 7: Molecular Evolutionary Genetics Analysis
428 version 7. 0 for bigger datasets. Mol Biol Evol 3:1870-1874

429 Kusunoki S, Ezaki T (1992) Proposal of *Mycobacterium peregrinum* sp. nov., nom. rev., and
430 elevation of *Mycobacterium chelonae* subsp. *abscessus* (Kubica et al.) to species status:
431 *Mycobacterium abscessus* comb. nov. Int J Syst Bacteriol 42:240-245

432 Kuykendall LD, Roy MA, O'Neill JJ, Devine TT (1988) Fatty acids, antibiotic resistance, and
433 deoxyribonucleic acid homology groups of *Bradyrhizobium japonicum*. Int J Syst Evol
434 Microbiol 38:358-361

435 Lechevalier MP, Lechevalier HA (1970) Composition of whole-cell hydrolysates as a criterion
436 in the classification of aerobic actinomycetes. In: Prauser H (ed) *The Actinomycetales*,
437 Gustav Fischer Verlag, Jena, pp 311-316

438 Lechevalier MP, De Bièvre C, Lechevalier H (1977) Chemotaxonomy of aerobic
439 actinomycetes: phospholipid composition. Biochem Syst Evol 5:249-260

440 Lehmann KB, Neumann R (1896) Atlas und Grundriss der Bakteriologie und Lehrbuch der
441 speziellen bakteriologischen Diagnostik. J. F. Lehmann, München

442 Loret JF, Greub G (2010) Free-living amoebae: Biological by-passes in water treatment. Int J
443 Hyg Environ Heal 213: 167–175.

444 Lorian V (1968) Differentiation of *Mycobacterium tuberculosis* and Runyon Group 3 "V"
445 strains on direct cord-reading agar. Am Rev Resp Dis 97:1133-1135

446 MacFaddin JF (1985) Media for Isolation - Cultivation - Identification - Maintenance of
447 Medical Bacteria. vol 1, Williams & Wilkins, Baltimore

448 Magee JG, Ward AC (2012) Genus I. *Mycobacterium* Lehmann and Neumann 1896, 363^{AL}, In:
449 Goodfellow M, Kämpfer P, Busse HJ, Trujillo ME, Suzuki KI, Ludwig W, Whitman

450 WB (eds) Bergey's Manual of Systematic Bacteriology, 2nd edn, vol 5. The
451 *Actinobacteria*, Part A. Springer, New York, pp 312-375

452 McNabb A, Eisler D, Adie K, Amos M, Rodrigues M, Stephens G, Black WA et al (2004)
453 Assessment of partial sequencing of the 65-kilodalton heat shock protein gene (*hsp65*)
454 for routine identification of *Mycobacterium* species isolated from clinical sources. J
455 Clin Microbiol 42:3000-3011

456 Meier-Kolthoff JP, Auch AF, Klenk HP, Göker M (2013a) Genome sequence-based species
457 delimitation with confidence intervals and improved distance functions. BMC
458 Bioinformatics 14:60

459 Meier-Kolthoff JP, Göker M, Spröer C, Klenk HP (2013b) When should a DDH experiment
460 be mandatory in microbial taxonomy? Arch Microbiol 195(6):413-418

461 Meier-Kolthoff JP, Hahnke RL, Petersen J, Scheuner C, Michael V, Fiebig A, Rohde C et al
462 (2014) Complete genome sequence of DSM 30083(T), the type strain (U5/41(T)) of
463 *Escherichia coli*, and a proposal for delineating subspecies in microbial taxonomy.
464 Stand Genomic Sci 9:2

465 Miller LT (1982) Single derivatization method for routine analysis of bacterial whole-cell fatty
466 acid methyl esters, including hydroxy acids. J Clin Microbiol 16(3):584-586

467 Minnikin DE, Hutchinson IG, Caldicott AB, Goodfellow M (1980) Thin-layer chromatography
468 of methanolysates of mycolic acid-containing bacteria. J ChromatogrA. 188: 221-233

469 Minnikin DE, O'Donnell AG, Goodfellow M, Alderson G, Athalye M, Schaal A, Parlett JH
470 (1984) An integrated procedure for the extraction of bacterial isoprenoid quinones and
471 polar lipids. J Microbiol Meth 2(5):233-241

472 Minnikin DE, Minnikin SM, Parlett JH, Goodfellow M (1985) Mycolic acid patterns of some
473 rapidly-growing species of *Mycobacterium*. Zentralbl Bakteriol Mikrobiol Hyg A.
474 259:446-60

475 Nishiuchi Y, Iwamoto T, Maruyama F (2017) Infection sources of a common non-tuberculous
476 mycobacterial pathogen, *Mycobacterium avium* complex. Front Med. 4: 27.

477 Nouioui I, Carro L, Teramoto K, Igual JM, Jando M, Montero-Calasanz MC, Sutcliffe I et al
478 (2017) *Mycobacterium eburneum* sp. nov., a non-chromogenic, fast-growing strain
479 isolated from sputum. Int J Syst Evol Microbiol 67: 3174-3181.

480 Palomino JC, Leão SC, Ritacco V (2007) Tuberculosis 2007- From basic science to patient
481 care. Institute of Tropical Medicine. Belgium, Brazil, Argentina

482 Parker CT, Tindall BJ, Garrity GM (2015) International Code of Nomenclature of Prokaryotes
483 (2008 Revision). Int J Syst Microbiol doi:10.1099/ijsem.0.000778

484 Pattengale ND, Alipour M, Bininda-Emonds OR, Moret BM, Stamatakis A (2010) How many
485 bootstrap replicates are necessary? *J Comput Biol* 17(3):337-354

486 Pontiroli A, Khera TT, Oakley BB, Mason S, Dowd SE, Travis ER, Erenso G, Aseffa A,
487 Courtenay O, Wellington EMH (2013) Prospecting environmental mycobacteria:
488 combined molecular approaches reveal unprecedented diversity. *PLOS One* 8:e68648.

489 Ramaprasad EV, Rizvi A, Banerjee S, Sasikala C, Ramana CV (2016) *Mycobacterium oryzae*
490 sp. nov., a scotochromogenic, rapidly growing species is able to infect human
491 macrophage cell line. *Int J Syst Evol Microbiol* 66:4530-4536

492 Ribón W (2012) Biochemical isolation and identification of mycobacteria, In: Jimenez-Lopez
493 JC (ed) *Biochemical Testing*. InTech. doi:10.5772/34309.
494 [https://www.intechopen.com/books/biochemical-testing/biochemical-isolation-and-](https://www.intechopen.com/books/biochemical-testing/biochemical-isolation-and-identification-of-mycobacteria)
495 [identification-of-mycobacteria](https://www.intechopen.com/books/biochemical-testing/biochemical-isolation-and-identification-of-mycobacteria)

496 Richter M, Rosselló-Móra R (2009) Shifting the genomic gold standard for the prokaryotic
497 species definition. *Proc Natl Acad Sci U S A* 106(45):19126-19131

498 Rodriguez RLM, Konstantinidis KT (2016) The enveomics collection: a toolbox for
499 specialized analyses of microbial genomes and metagenomes. *PeerJ Preprints*
500 4:e1900v1

501 Roguet A, Therial C, Saad M, Boudahmane L, Moulin L, Lucas F. S (2016) High
502 mycobacterial diversity in recreational lakes. *Antonie van Leeuwenhoek*. 109: 619-631.

503 Runyon EH, Karlson AG, Kubica GP, Wayne LG (1980) *Mycobacterium*. American Society
504 for Microbiology. Washington DC.

505 Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing
506 phylogenetic trees. *Mol Biol Evol* 4:406-425

507 Salah IB, Ghigo E, Drancourt M (2009) Free-living amoebae, a training field for macrophage
508 resistance of mycobacteria. *Clin Microbiol Infect* 15:894-905.

509 Sangal V, Jones AL, Goodfellow M, Hoskisson PA, Kämpfer P, Sutcliffe IC (2015) Genomic
510 analyses confirm close relatedness between *Rhodococcus defluvi* and *Rhodococcus*
511 *equi* (*Rhodococcus hoagii*). *Arch Microbiol* 197:113-116

512 Sasser MJ (1990) Identification of bacteria by gas chromatography of cellular fatty acids.
513 Technical Note 101, Microbial ID USA: Inc, Newark, Del

514 September SM, Brözel VS, Venter SN (2004) Diversity of nontuberculoïd *Mycobacterium*
515 species in biofilms of urban and semi urban drinking water distribution systems. *Appl*
516 *Environ Microbiol* 70: 7571–7573.

517

518 Sequeira de Latini MD, Barrera L (2008) Manual para el Diagnóstico Bacteriológico de la
519 Tuberculosis: Normas y Guía Técnica, Parte I Baciloscopía. Organización
520 Panamericana de la Salud

521 Shahraki AH, Trovato A, Mirsaeidi M, Borroni E, Heidarieh P, Hashemzadeh M, et al (2017)
522 *Mycobacterium persicum* sp. nov., a novel species closely related to *Mycobacterium*
523 *kansasii* and *Mycobacterium gastri*. Int J Syst Evol Microbiol 67:1766–1770

524 Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of
525 large phylogenies. Bioinformatics 30:1312-1313

526 Staneck JL, Roberts GD (1974) Simplified approach to identification of aerobic actinomycetes
527 by thin layer chromatography. Appl Microbiol 28:226-231

528 Swofford DL (2002) PAUP*: Phylogenetic Analysis Using Parsimony (*and Other Methods),
529 Version 4.0. Sinauer Associates, Sunderland

530 Thomas V, Loret JF, Jousset M, Greub G (2008) Biodiversity of amoebae and amoebae-
531 resisting bacteria in a drinking water treatment plant. Environ Microbiol 10: 2728–2745.

532 Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997) The Clustal X
533 windows interface: flexible strategies for multiple sequence alignment aided by quality
534 analysis tools. Nucleic Acids Res 25:876-4882

535 Tomioka H, Saito H, Sato K, Dawson DJ (1990) Arylsulfatase activity for differentiating
536 *Mycobacterium avium* and *Mycobacterium intracellulare*. J Clin Microbiol 28:2104-
537 2106

538 Tortoli E, Kohl TA, Brown-Elliott BA, Trovato A, Leão SC, Garcia MJ et al (2016) Emended
539 description of *Mycobacterium abscessus*, *Mycobacterium abscessus* subsp. *abscessus*
540 and *Mycobacterium abscessus* subsp. *bolletii* and designation of *Mycobacterium*
541 *abscessus* subsp. *massiliense* comb. nov. Int J Syst Evol Microbiol 66:4471-4479

542 Tran PM, Dahl JL (2016) *Mycobacterium sarraceniae* sp. nov. and *Mycobacterium helvum* sp.
543 nov., isolated from the pitcher plant *Sarracenia purpurea*. Int J Syst Evol Microbiol
544 66:4480-4485

545 Tsukamura M (1966) Adansonian classification of mycobacteria. J Gen Microbiol 45:253-273

546 Tsukamura M, van der Meulen HJ, Grabow WOK (1983) Numerical taxonomy of rapidly
547 growing, scotochromogenic mycobacteria of the *Mycobacterium parafortuitum*
548 complex: *Mycobacterium austroafricanum* sp. nov. and *Mycobacterium diernhoferi* sp.
549 nov., nom. rev. Int J Syst Bacteriol 33:460-469

- 550 Vaas LAI, Sikorski J, Michael V, Göker M, Klenk HP (2012) Visualization and curve-
551 parameter estimation strategies for efficient exploration of phenotype microarray
552 kinetics. PLoS ONE 7: e34846
- 553 Vaas LAI, Sikorski J, Hofner B, Fiebig A, Buddruhs N, Klenk HP, Göker M (2013) Opm: an
554 R package for analysing OmniLog (R) phenotype microarray data. Bioinformatics
555 29:1823–1824
- 556 Wayne LG, Kubica GP (1986) The mycobacteria, In: Sneath PHA, Mair S, Sharpe ME, Holt
557 JG (eds) Bergey's Manual of Systematic Bacteriology, vol 2, Williams & Wilkins,
558 Baltimore, pp 1435-1457
- 559 Wayne L, Brenner DJ, Colwell RR, Grimont PAD, Kandler O, Krichevsky MI, Moore LH
560 et al (1987) Report of the *ad hoc* committee on reconciliation of approaches to bacterial
561 systematics. Int J Syst Evol Bacteriol 37:463-464
- 562 Yoon SH, Ha SM, Kwon S, Lim J, Kim Y, Seo H, Chun J (2017) Introducing EzBioCloud: A
563 taxonomically united database of 16S rRNA and whole genome assemblies. Int J Syst
564 Evol Microbiol 67:1613-1617
- 565 Zopf W (1883) Die Spaltpilze. Edward Trewendt, Breslau, pp 1-100

566

567

568 **Table 1.** Genetic distances between strains CECT 8778^T and CECT 8779^T and between them and their close phylogenetic neighbours

		MLSA (Kimura 2-parameter) distance													
		1	2	3	4	5	6	7	8	9	10	11	12	13	14
1	Strain CECT 8779 ^T														
2	Strain CECT 8778 ^T	0,051													
3	<i>Mycobacterium peregrinum</i>	0,031	0,039												
4	<i>Mycobacterium mageritense</i>	0,030	0,039	0,009											
5	<i>Mycobacterium wolinskyi</i>	0,032	0,036	0,012	0,012										
6	<i>Mycobacterium chlorophenicum</i>	0,036	0,034	0,031	0,034	0,034									
7	<i>Mycobacterium chubuense</i>	0,037	0,036	0,033	0,036	0,037	0,006								
8	<i>Mycobacterium psychrotolerans</i>	0,037	0,039	0,035	0,036	0,037	0,015	0,016							
9	<i>Mycobacterium austroafricanum</i>	0,048	0,028	0,046	0,048	0,049	0,029	0,032	0,036						
10	<i>Mycobacterium vanbaalenii</i>	0,047	0,026	0,044	0,046	0,046	0,026	0,029	0,035	0,003					
11	<i>Mycobacterium aurum</i>	0,050	0,029	0,038	0,041	0,041	0,032	0,030	0,037	0,028	0,026				
12	<i>Mycobacterium rufum</i>	0,040	0,036	0,034	0,037	0,036	0,009	0,012	0,018	0,035	0,034	0,036			
13	<i>Mycobacterium arcueilense</i>	0,054	0,044	0,030	0,035	0,036	0,049	0,051	0,046	0,054	0,052	0,046	0,051		
14	<i>Mycobacterium alvei</i>	0,056	0,043	0,032	0,036	0,039	0,049	0,052	0,047	0,059	0,057	0,046	0,052	0,008	
15	<i>Tsukamurella paurometabola</i>	0,071	0,077	0,074	0,076	0,073	0,073	0,073	0,073	0,083	0,081	0,087	0,077	0,084	0,086

569

570

571 **Table 2.** Phenotypic features that distinguish strains CECT 8778^T and CECT 8779^T from one another and from their near phylogenetic
572 neighbours. All data are from the present study.
573

	strain CECT 8778 ^T	<i>M. aurum</i> DSM 43999 ^T	<i>M. austroafricanum</i> DSM 44191 ^T	<i>M. vanbaalenii</i> DSM 7152 ^T	strain CECT 8779 ^T	<i>M. mageritense</i> DSM 44476 ^T
Biochemical tests:						
Arylsulfatase 3 days	+	-	+	+	+	+
Heat stable catalase 68°C	+	-	-	-	-	-
Niacin, Tween 80	+	+	-	-	+	+
Urea hydrolysis	+	-	-	+	-	-
GEN III Biolog microplate tests						
Utilisation of sugars:						
D-Arabitol,	+	-	+	-	+	-
Dextrin	+	-	-	+	-	+
D-Galactose	-	+	+	-	-	-
<i>N</i> -acetyl-D-Glucosamine	+	-	+	-	+	+
3- <i>o</i> -methyl-D-Glucose	-	-	+	-	-	-
D-Glucose-6-phosphate	-	-	+	-	+	-
Glycerol	-	+	+	+	+	+
<i>myo</i> -Inositol	+	+	+	+	-	-
D-Maltose	-	-	+	-	+	-
<i>N</i> -acetyl-β-D-Mannosamine	-	-	+	-	+	+
D-Mannose, D-sorbitol, D-trehalose	+	+	+	+	+	-
L-Rhamnose	-	-	+	-	+	+
D-Salicin	+	-	+	-	-	-

Sucrose	-	+	+	+	+	-
D-Turanose	+	+	-	+	+	-
Utilisation of organic acids:						
Butyric acid	-	+	+	+	+	+
β -amino- <i>n</i> -Butyric acid	+	+	-	-	+	+
α -hydroxy-Butyric acid, α -keto-Butyric acid	-	+	+	+	-	+
Citric acid	-	-	-	-	+	+
D-Galacturonic acid	-	+	-	-	-	-
α -keto-Glutamic acid	-	+	-	+	+	+
L-Lactic acid	-	+	+	+	+	+
D-and L-Malic acid	+	+	+	+	-	+
Methyl pyruvate	-	+	+	+	+	-
Quinic acid	+	-	+	-	-	-
D-Saccharic acid	+	+	+	+	-	-
Bromo-Succinic acid	+	+	+	+	-	+
Utilisation of amino acids:						
L-Alanine	-	+	+	-	+	+
L-Aspartic acid	-	-	+	-	+	+
Glycyl- L-proline	+	-	+	+	+	+
L-Histidine	-	-	-	-	+	+
D-Serine #2, L- Pyroglutamic acid	-	+	-	-	-	+
L-Serine	+	-	+	-	-	+
Resistance to:						
Lincomycin	-	-	-	-	-	+
Lithium chloride	-	+	+	+	+	-
Nalidixic acid, sodium chloride (1% w/v)	-	+	+	+	+	+

Rifamycin SV, sodium bromate	+	+	-	+	+	+
Sodium chloride (4% w/v)	-	+	-	-	+	+
Sodium chloride (8% w/v)	-	+	-	-	+	-
Sodium formate	+	+	+	+	-	+
Tetrazolium violet	+	-	-	-	+	+
Tetrazolium blue	+	-	-	-	-	-
Troleandomycin	-	+	-	-	-	-
Vancomycin	-	+	-	+	-	-
Growth in presence of:						
Gelatin	-	+	-	-	-	+
Pectin	-	+	+	+	+	-
Tween 40	+	-	+	+	+	+
Chemotaxonomic traits						
Polar lipids	DPG, PE, PI, GL GPL1-2	DPG, PE, PI, GL GPL1-2	DPG, PE, PI, GL GPL1-2	DPG, PE, PI, GL GPL1-2	DPG, PE, PI, GL GPL1-2	DPG, PE, PI, PG, GL GPL1-2
Fatty acids (>20 %)	Summed feature 2	C _{16:0}	C _{16:0}	Summed feature 2	C _{16:0} , C _{18:1} ω9c	C _{16:0} , C _{18:1} ω9c
Mycolic acids	<i>α</i> - and <i>decarboxy</i> -mycolates	<i>α</i> - <i>keto</i> -mycolates and wax esters*	<i>α</i> - <i>keto</i> -mycolates and wax esters*	<i>α</i> - <i>keto</i> -mycolates and wax esters*	<i>α</i> , <i>keto</i> - <i>methoxy</i> mycolic acids	<i>α</i> -, <i>α</i> '- and <i>epoxy</i> -mycolates**
DNA G+C content (%)	65.3	67.5	67.4	67.8	69.4	66.6

574 + Positive reaction; - negative reaction.

575 Positive results recorded for all of the strains: arylsulfatase (14 days) and catalase (biochemical tests); utilisation of acetic acid, acetoacetic acid, L-arginine, β-*hydroxy*-butyric acid, D-glucose, D-fructose, D-gluconic acid, L-glutamic acid, D-mannitol and propionic acid; growth at pH 6 and in presence of aztreonam, potassium tellurite and 1% sodium lactate. Negative results detected for all of the strains: utilisation of D-aspartic acid, D-

578 cellobiose, D-fructose-6-phosphate, D-and L-fucose, L-galactonic acid- γ -lactone, *N*-acetyl-D-galactosamine, β -gentiobiose, β -methyl-D-glucoside,
579 glucuronamide, D-glucuronic acid, D-lactic acid methyl ester, α -D-lactose, D-melibiose, mucic acid, *N*-acetyl-neuraminic acid, *p*-hydroxy-
580 phenylacetic acid, D-raffinose and stachyose; growth at pH 5 and in the presence of fusidic acid, guanidine hydrochloride, inosine, minocycline
581 and niaproof. Abbreviation: DPG: diphosphatidylglycerol; GPL1-2: glycopospholipids; GL: glycolipid; PE: phosphatidylethanolamine; PI:
582 phosphatidylinositol; PG: glycopospholipid; summed features 2, C_{17:1} ω 7c /18 alcohol; *data taken from Mininkin et al. 1985, Magee and Ward 2012;
583 ** data taken from Domenech et al. 1997.

584

585

586

587

588

589

590

591

592

593

594 **Table 3.** Fatty acid profiles (%) of strain CECT 8778^T (1), *M. aurum* DSM 43999^T (2), *M.*
 595 *austroafricanum* DSM 44191^T (3), *M. vanbaalenii* DSM 7152^T (4), strain CECT 8779^T (5) and
 596 *M. mageritense* DSM 44476^T (6). All data are from the present study.

	1	2	3	4	5	6
C _{14:0}	3.6	3.4	2.6	2.4	5.2	7.3
C _{16:1} ω9c	1.0	0.8	0.9	1.5	3.6	-
C _{16:1} ω6c	4.2	6.9	10.7	6.8	6.5	10.3
C _{16:1} ω7c	0.5	-	-	-	1.6	4.1
C _{16:0}	13.3	25.4	27.6	17.0	25.4	40.0
Summed feature 2	43.2	12.6	15.7	27.6	7.6	-
C _{18:1} ω9c	7.6	19.7	8.9	13.8	26.2	28.3
C _{18:0}	-	1.6	2.7	1.6	-	-
10 Me-C _{18:0}	11.0	12.6	14.2	8.6	7.8	8.7
Summed feature 3	12.3	15.2	15.0	16.8	-	-
C _{20:0}	-	-	0.3	0.3	-	-

597 Summed features 2, C_{17:1} ω7c /18 alcohol and summed feature 3, 20:0 ALC 18.838/ECL20:0
 598 ALC.

599
 600
 601
 602


603 **Figure legends**

604 Fig. 1. Maximum-likelihood phylogenetic tree based on almost complete 16S rRNA gene
605 sequences generated using the GTR+GAMMA model and midpoint-rooting showing
606 relationships between strains CECT 8778^T and CECT 8779^T and between them and their close
607 phylogenetic neighbours. The numbers above the branches are bootstrap support values greater
608 than 60% for ML (left) and MP (right). The scale bar indicates 0.007 substitutions per site.

609 Fig. 2. Maximum-likelihood phylogenetic tree based on concatenated sequences of 16S rRNA,
610 *hsp65*, *rpoB* and *recA* gene sequences (3257 nt) showing relationships between strains CECT
611 8778^T and CECT 8779^T and between them and their close phylogenetic neighbours. The tree
612 was inferred using the GTR+GAMMA model. The branches are scaled in terms of the expected
613 number of substitutions per site. The numbers above the branches are bootstrap support values
614 when larger than 60% from ML (left) and MP (right). The scale bar indicates 0.02 substitutions
615 per site. The accession numbers of the MLSA gene sequences are listed in Table S2.


616

617




618
619
620
621
622

0.005



623

0.02



[Click here to download Figure Fig. S1..pdf](#)

Fig. S1. Neighbour-joining MLSA phylogenetic tree based on concatenated sequences of 16S rRNA, *hsp65*, *rpoB* and *recA* genes showing relationships between strains CECT 8778^T and CECT 8779^T and between them and their close phylogenetic neighbours. The numbers above the branches are bootstrap support values greater than 60%. The scale bar indicates 0.005 substitutions per site.

Figure
0.06




Fig. S2. Maximum-likelihood phylogenetic tree based on *recA* partial gene sequences generated using the GTR+GAMMA model and midpoint-rooting showing relationships between strains CECT 8778^T and CECT 8779^T and between them and their close phylogenetic neighbours. The numbers above the branches are bootstrap support values greater than 60% for ML (left) and MP (right). The scale bar indicates 0.06 substitutions per site.




Fig. S3. Maximum-likelihood phylogenetic tree based on *rpoB* partial gene sequences generated using the GTR+GAMMA model and midpoint-rooting showing relationships between strains CECT 8778^T and CECT 8779^T and between them and their close phylogenetic neighbours. The numbers above the branches are bootstrap support values greater than 60% for ML (left) and MP (right). The scale bar indicates 0.02 substitutions per site.

Figure
0.02

⊗ LFN KH UH WR GRZO ORDC




Fig. S4. Maximum-likelihood phylogenetic tree based on *hsp65* partial gene sequences generated using the GTR+GAMMA model and midpoint-rooting showing relationships between strains CECT 8778^T and CECT 8779^T and between them and their close phylogenetic neighbours. The numbers above the branches are bootstrap support values greater than 60% for ML (left) and MP (right). The scale bar indicates 0.02 substitutions per site.