

Northumbria Research Link

Citation: Fitzpatrick, John, Hicks, Kirsty and Hayes, Phil (2018) Dose-Response Relationship between Training Load and Changes in Aerobic Fitness in Professional Youth Soccer Players. *International Journal of Sports Physiology and Performance*, 13 (10). pp. 1365-1370. ISSN 1555-0265

Published by: Human Kinetics

URL: <http://dx.doi.org/10.1123/ijsp.2017-0843> <<http://dx.doi.org/10.1123/ijsp.2017-0843>>

This version was downloaded from Northumbria Research Link: <http://nrl.northumbria.ac.uk/34469/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



Northumbria
University
NEWCASTLE



UniversityLibrary

Title: Dose-Response Relationship between Training Load and Changes in Aerobic Fitness in Professional Youth Soccer Players

Submission Type: Original Investigation

Authors: John F. Fitzpatrick^{1,2}, Kirsty M. Hicks¹, Philip R. Hayes¹

Author Affiliations:

¹Department Sport, Exercise and Rehabilitation, Northumbria University, Newcastle-upon-Tyne, UK.

²Sports Science and Medical Department, Newcastle United Football Club, Newcastle-upon-Tyne, UK.

Corresponding Author:

Mr John F. Fitzpatrick,
Sports Science and Medical Department,
Newcastle United Football Club,
Greenlee Drive, Little Benton,
Newcastle upon Tyne,
NE7 7SF;
Tel: 0191 2153200;
Email: john.fitzpatrick@nufc.co.uk

Preferred Running Head: Dose-Response Relationship in Soccer

Abstract Word Count: 246

Text Only Word Count: 3195

Number of Figures: 3

Number of Tables: 2

Abstract

Purpose: The aim of this study was to compare the dose-response relationship between, traditional arbitrary speed thresholds versus an individualised approach, with changes in aerobic fitness in professional youth soccer players. **Methods:** Fourteen youth soccer players, completed a 1500 metre time trial to estimate maximal aerobic speed (km.h⁻¹, (MAS)) at the start and the end of a six week period. Training load was monitored on a daily basis during this study. External load measures were; total distance covered (TD), total acceleration and deceleration distance > 2m.s⁻² (A/D Load). Arbitrary high speed running measures were; metres covered and time spent > 17 km.h⁻¹ (m>HSD, t>HSD) and 21 km.h⁻¹ (m>VHSD, t>VHSD). Individualised high speed running measures were; metres covered and time spent > MAS km.h⁻¹ (m>MAS, t>MAS) and 30% anaerobic speed reserve (m>30ASR, t>30ASR). In addition, internal load measures were also collected; heart rate exertion (HRE) and rating of perceived exertion (RPE). Linear regression analysis was used to establish the dose-response relationship between mean weekly training load and changes in aerobic fitness. **Results:** Substantial very large associations were found between t>MAS and changes in aerobic fitness ($R^2 = 0.59$). Substantial large associations were found for t>30ASR ($R^2 = 0.38$) and m>MAS ($R^2 = 0.25$). Unsubstantial associations were found for all other variables. **Conclusion:** An individualised approach to monitoring training load, in particular t>MAS, may be a more appropriate method than using traditional arbitrary speed thresholds when monitoring the dose-response relationship between training load and changes in aerobic fitness.

Introduction

The physiological response to a given training load is commonly called the dose-response relationship and is considered a fundamental component of training.¹ It has been suggested that a valid measure of training load should show a strong dose-response relationship with a particular training outcome, such as, fitness level, fatigue status or injury risk.² Training load measures that demonstrate a strong dose-response relationship will provide practitioners with a greater understanding of how their athletes may respond to a given training stimulus.³ Giving them an ability to prescribe training with confidence, and that it will produce a predictable outcome within a defined period.¹ Improving a practitioner's understanding of the dose-response relationship will allow them to optimally plan training to maximise fitness, whilst minimising fatigue and injury risk.

There is a wealth of information within the scientific literature about soccer players training and match loads.⁴ Equally, there are numerous studies documenting specific responses to training, whether it be fitness,⁵ fatigue⁶ or injury risk.⁷ However, there is still very limited evidence about the dose-response relationship between training load and specific training outcomes in soccer players. It is well established that the internal response to a given external load is what drives training adaptation.⁸ A comparison of methods used to establish internal load was investigated by Akubat et al⁹, with an individualised training impulse (iTRIMP) showing the strongest relationship with changes in fitness. However, given the time associated with conducting the assessments needed to use iTRIMP, this method may not be the most practically applicable for a squad of soccer players. Two other examples of dose-response studies within the literature have investigated the relationship between training load and fatigue. Thorpe et al¹⁰ found a large relationship between high intensity running (>14.4 km.h-1) and next day subjective fatigue. Scott and Lovell¹¹ built upon this work and compared the dose-response relationship between arbitrary high intensity running (>17.8 km.h-1) and various

individualised methods of assessing high speed running, with subsequent fatigue. Although, relationships were classified as small, there was no difference between arbitrary and individualised methods of quantifying training load. However, the literature on the relationship between external training load and fitness is sparse, particularly with regards to individualising external load.

Abt and Lovell¹² first proposed that high intensity speed thresholds, used during soccer match-play, should be individualised based on the second ventilatory threshold in a similar manner to the methods proposed by Lucia et al¹³. They found the median velocity at the second ventilatory threshold to be 15 km.h⁻¹. This resulted in a three-fold increase in the high intensity distance covered when compared to the arbitrary distance covered above 19.8 km.h⁻¹, meaning practitioners are substantially underestimating the amount of high intensity running their players are undertaking during training and match play. Although these findings show the need to individualise training loads based on physiological characteristics, the methods used by Abt and Lovell¹² require extensive laboratory testing which may not be applicable in an applied environment. Other methods have been employed to assess players training load with reference to individual fitness characteristics gained via field based assessments. Percentage of maximal sprint speed (MSS)¹⁴ has been used previously, however, it has been highlighted that using one single fitness characteristic may not reflect the complete locomotor profile of a player.^{15,16} Alternatively, Mendez-Villanueva et al¹⁷ used a technique which encapsulated field based testing data to estimate a players' aerobic (maximal aerobic speed [MAS]) and anaerobic capabilities (MSS). This technique allows for an estimation of a players' anaerobic speed reserve (ASR) and has been used to establish a players' transition (>30ASR) into sprint work.^{17,18} Furthermore, Hunter et al¹⁵ stated that a method using field based measures of MAS, MSS and ASR poses a more ecologically valid, economical and practical technique for individualising thresholds.

It is well established that aerobic capacity is an important physiological factor during soccer performance.^{19,20} The assessment of MAS is therefore warranted within soccer as a performance indicator. Given MAS has been used as a measure of change in physical fitness in previous studies²¹ and its usefulness in an applied setting when being used to prescribe training loads²², MAS was selected as a dependent variable, alongside MSS, in the present study.

To our knowledge there has yet to be a study comparing the dose-response relationship between arbitrary and individualised methods for assessing training loads and a specific training outcome, such as changes in aerobic fitness. It was therefore the aim of this study to compare the dose-response relationship between traditional arbitrary speed thresholds versus an individualised approach utilising MAS and MSS with changes in aerobic fitness in professional youth soccer players.

Methods

Subjects

Fourteen male youth soccer players (age: 17.1 ± 0.5 years, height: 178.3 ± 4.6 cm, body mass: 70.9 ± 5.8 kg) (defenders = 5, midfielders = 6, forwards = 3), competing in the English Under-18 Premier League agreed to participate in the study. Data were collected over a 6-week period, during an in-season competition phase (August-September).

Design

For this observational research, players took part in normal team training throughout the 6-week period as prescribed by club coaching staff. This included 6 competitive matches, 23 training sessions, and 13 rest days. No structured conditioning was conducted throughout this study. Physical assessments were completed at the start and the end of the 6-week period, with training load monitored throughout. Before inclusion in the study, players were examined

by the club medical staff and were deemed to be free from illness and injury. The study was granted institutional ethics approval prior to commencement and conformed to the declaration of Helsinki. Informed consent was provided by all players and by their parents for players under 18 years of age.

Methodology

Prior to the start of the 6-week in-season period, players completed a testing battery to estimate MSS and MAS. Following a standardised warm up players completed two maximal 40 metre sprints, with three minutes recovery between efforts. Split times were recorded at 30 and 40 metres (Brower Timing Systems, Draper, UT), with the time taken to complete this 10 metre split used as MSS ($\text{km}\cdot\text{h}^{-1}$).¹⁷ The best MSS over the two sprints was used for the purpose of this study (minimum detectable change [MDC] 1.7%, unpublished observations). Following 10 minutes of rest players then completed a 1500-metre time trial (TT) on an outdoor artificial pitch. The time taken to complete this TT was recorded and the average speed calculated to estimate MAS ($\text{km}\cdot\text{h}^{-1}$) (MDC 1.3%, unpublished observations). This method of assessing aerobic fitness has previously been validated.^{23,24} Additionally, ASR was calculated from this data in accordance with previous literature¹⁷ (MDC 1.2%, unpublished observations).

Training load was calculated for every training session and match played during the 6-week period, using a number of different methods; global positioning system (GPS), heart rate telemetry (HR) and session rating of perceived exertion (sRPE). External load was measured using GPS units (MinimaxX S4, Catapult Sports, Melbourne, Australia) sampling at a frequency of 10 Hz. GPS devices were switched on at least 15 minutes prior to each training session and match to ensure a full satellite signal (number of satellites: 14.4 ± 0.5 ; horizontal dilution of precision: 0.81 ± 0.10). Players were fitted with the same device for each session.

The GPS devices were worn between the scapular in a tight-fitting vest to reduce movement

artefact. Following each training session and match, data were downloaded using the manufacturer's software (Catapult Sprint, Version 5.1.7, Catapult Sports, Melbourne, Australia). This GPS system has previously been shown to provide valid and reliable estimates of instantaneous velocity during acceleration, deceleration, and constant-velocity movements during linear, multidirectional, and soccer-specific activities.^{25,26}

The external load measures used for analysis were; total distance (TD), acceleration and deceleration distance $>2\text{m}\cdot\text{s}^{-2}$ (AD Load). Arbitrary high speed running measures were; metres covered above $17\text{ km}\cdot\text{h}^{-1}$ (m>HSD) and $21\text{ km}\cdot\text{h}^{-1}$ (m>VHSD) and time spent above $17\text{ km}\cdot\text{h}^{-1}$ (t>HSD) and $21\text{ km}\cdot\text{h}^{-1}$ (t>VHSD). These arbitrary speed thresholds were selected to match the group mean thresholds for MAS and 30% ASR. Individualised high speed running measures were; metres covered above MAS (m>MAS) and time spent above MAS (t>MAS). As suggested by previous research,^{15,17} to define a player's transition into anaerobic work, metres covered above 30% ASR (m>30ASR) and time spent above 30% ASR (t>30ASR) were also calculated.

Measurements of HR were collected using a short-range telemetry HR transmitter strap recording at 5 s intervals (Polar T34, Polar Electro, OY, Finland). Data were downloaded and analysed using specific software (Catapult Sprint, Version 5.1.7, Catapult Sports, Melbourne, Australia). A heart rate exertion (HRE) score was calculated based on Edwards²⁷ training impulse, using the time spent in five HR zones and multiplied by a zone specific weighting factor.

Approximately 30 minutes after each training session and match, players reported their RPE using the method of Foster et al²⁸ Each player was asked verbally, in private, how hard they found each session, reporting their subjective perception of effort using the Borg 10-point category-ratio scale. sRPE was subsequently calculated by multiplying the RPE by the number

of training or match minutes played. Players were familiarised with the use of the RPE scale prior to the start of the six-week study period.

Statistical Analysis

Descriptive statistics are presented as means \pm standard deviations. Pre-and post-measures of MAS and MSS were compared via standardised changes in the mean (effect size; ES) using a custom spreadsheet.²⁹ The following criteria were adopted to interpret the magnitude of change; $>0.2-0.6$, small; $>0.6-1.2$, moderate; $>1.2-2$, large; >2 , very large.³⁰ The magnitude of change was classified as a substantial increase or substantial decrease when there was a 75% or greater likelihood of the change being equal to or greater than the ES \pm 0.2 (small).³⁰ To understand the strength and direction of the dose-response relationship between the mean weekly training load and changes in fitness, Pearson's product moment correlation coefficients (r) were calculated. Where the 90% confidence interval overlaps both the positive and negative threshold by $\geq 5\%$ the relationship was deemed unclear.³⁰ The following criteria were adopted to interpret the magnitude of the relationship; 0.0-0.1 trivial; $>0.1-0.3$ small; $>0.3-0.5$ moderate; $>0.5-0.7$ large; $>0.7-0.9$ very large; $>0.9-0.99$ nearly perfect; 1.00 perfect.²⁵

Linear regression analysis was conducted following visual inspection of all relationships to identify a linear or curvilinear relationship. To determine the level of variance in the dependent variable explained by training load the coefficient of determination (R^2) was calculated via linear regression analysis. Additionally, to understand the error associated with each dose-response relationship the standard error of prediction (SEP) was calculated.³¹

Results

A total of 387 training and match files were analysed for the 14 players during the 6-week in-season training period. Mean \pm SD weekly training loads are displayed in Table 1. Mean \pm SD weekly and daily $t >$ MAS during the training period are displayed in Figure 1.

The mean change in MAS over the training period was $0.11 \pm 0.12 \text{ km}\cdot\text{h}^{-1}$ (ES: 0.15, possibly trivial, 31/69/0) and the mean change for MSS was $0.27 \pm 0.20 \text{ km}\cdot\text{h}^{-1}$ (ES: 0.16, possibly trivial, 26/74/0).

A *very large* linear relationship was found between $t > \text{MAS}$ and changes in MAS ($r = 0.77$ [90% CI 0.48 to 0.91], $R^2 = 0.59$) (Figure 2). Also, *large* relationships were found between $t > 30\text{ASR}$ ($r = 0.62$ [90% CI 0.22 to 0.84], $R^2 = 0.38$), $m > \text{MAS}$ ($r = 0.50$ [90% CI 0.06 to 0.78], $R^2 = 0.25$) and changes in MAS. Relationships between all other mean weekly arbitrary and individualised training load measures and changes in fitness parameters were found to be *unclear* (Tables 2 & 3).

Other external load measures, TD ($r = 0.26$ [90% CI -0.23 to 0.64]) and AD Load ($r = 0.20$ [90% CI -0.29 to 0.60]) displayed *unclear* relationships with changes in MAS. Similarly, internal load measures, HRE ($r = -0.21$ [90% CI -0.61 to 0.28]) and sRPE ($r = 0.22$ [90% CI -0.26 to 0.62]) were also *unclear*. In contrast, relationships with changes in MSS were identified for TD ($r = 0.46$ [90% CI 0.00 to 0.76] *possibly moderate*, $R^2 = 0.21$), AD Load ($r = 0.57$ [90% CI 0.15 to 0.81] *possibly large*, $R^2 = 0.32$) and HRE ($r = 0.40$ [90% CI -0.07 to 0.73] *possibly moderate*, $R^2 = 0.16$). However, sRPE ($r = 0.37$ [90% CI -0.11 to 0.71]) displayed an *unclear* relationship with changes in MSS.

Discussion

The aim of the present study was to examine the dose-response relationship between a range of measures quantifying training load and changes in aerobic fitness using a field based test of MAS. The training load was quantified using both arbitrary speed thresholds and an individualised approach utilising MAS and MSS. The key finding from the present study is that the use of individualised thresholds, specifically $t > \text{MAS}$, demonstrated a stronger dose-response relationship with changes in aerobic fitness than commonly used arbitrary thresholds.

This is the first study to assess the relationships between various measures of external training load and changes in aerobic fitness in youth soccer players. It has previously been shown that using individualised thresholds may better represent the true physiological demands of soccer training and match play placed on the individual.^{12,15} However, linking these individualised measures to specific training outcomes, such as improvements in aerobic fitness, had not previously been researched. The present study found that using an individualised approach, in which players training and match load was assessed based on the time spent above their MAS, has the strongest relationship with changes in aerobic fitness. This provides practitioners with important information for planning training loads, allowing them to more clearly understand the physical outcome from a given training dose.

It has been shown that high intensity distance covered ($>19.8 \text{ km}\cdot\text{h}^{-1}$) is one of the most commonly used measures of training load in elite soccer.³² This study has demonstrated however, that arbitrary thresholds of $17.0 \text{ km}\cdot\text{h}^{-1}$ and $21.0 \text{ km}\cdot\text{h}^{-1}$ presented *unclear* correlations with changes in MAS and MSS (Table 2). This should call into question the usefulness of these measures for assessing training load. Furthermore, as the dose-response relationship between these measures is unclear, it is very difficult for practitioners to make informed decisions about the desired training outcome based on arbitrary speed thresholds, which could lead to over/under-training, injury or illness. The authors acknowledge that the use of arbitrary thresholds is required to make comparisons between players possible, from a performance and talent identification standpoint. It could therefore be suggested that a joint approach, utilising both arbitrary and individualised thresholds is most advantageous.

Another novel aspect of the present study was the inclusion of time spent above both arbitrary and individualised speed thresholds. When analysing external training loads the most commonly used measure is distance covered above a specific threshold, this is evident in both research³³ and practice.³² However, the present study would suggest that time spent above a

high speed threshold may be a more robust measure of training load, given the stronger associations between time variables compared to distance variables. This would seem to be in line with research on endurance athletes where time spent around $\dot{V}\text{O}_2$ max is an important consideration when looking to improve aerobic fitness.³⁴ The differences between time spent and distance covered above a threshold may be a consequence of players running at higher speeds, this will lead to a large number of metres covered in a small amount of time. Therefore if $t > \text{MAS}$ is the parameter practitioners are looking to target, even paced running at a speed just above MAS (100-110%) may be a key factor when looking to improve aerobic fitness.

Depicted in Figure 1 is the mean weekly and daily training load for $t > \text{MAS}$ across this 6-week study period. Although this study was conducted over a relatively short period, the data may suggest that $t > \text{MAS}$ follows a similar weekly variation to that of other external load variables previously described.³⁵ It is worth noting the limited amount of time players spend above this threshold during match play (2.3 min). This was lower than during a match day -4 training session (3.0 min). Future research should look to assess $t > \text{MAS}$ longitudinally to confirm weekly training and match loads. It could be suggested that given the mean weekly $t > \text{MAS}$ of 7.4 min and the trivial mean change in fitness, that greater $t > \text{MAS}$ is needed throughout the training week, if players are to improve their fitness throughout the in-season. Furthermore, given the low $t > \text{MAS}$ acquired during soccer match play, specific running based interventions could be used to increase mean weekly $t > \text{MAS}$ and therefore improve aerobic fitness.

A limitation of the present study is the small sample size ($n=14$), although this is common in studies of players at a professional level. Moreover, this study was conducted over a relatively small in-season period; future work should look to replicate these results over longer training periods. Additionally, intervention studies looking to specifically target $t > \text{MAS}$ in order to improve aerobic fitness in soccer players would be warranted.

Practical Applications

There are a number of practical applications which practitioners may take from the present study. The results have shown a *very large* dose-response relationship between weekly $t > \text{MAS}$ and changes in aerobic fitness. This may allow the implementation of specific training programs from which practitioners can understand the expected outcome. By using the regression analysis data displayed in Table 3 and Figure 2 an estimated percentage change in aerobic fitness can be obtained from a given mean weekly $t > \text{MAS}$. For example, 6.5 minutes per week over a 6-week period would estimate a 0% change in fitness, which may be the target of an in-season maintenance phase. However, it is important to appreciate the error associated with that estimate, for $t > \text{MAS}$ this is 1% (Table 3), meaning 6.5 minutes per week may result in a -1% to +1% change in aerobic fitness. Therefore, if the desired outcome is to maintain fitness a target that is greater than the MDC may be more appropriate. 8.3 minutes per week would predict an improvement of $1.3\% \pm 1.0\%$, giving a range of 0.3% to 2.3% change in aerobic fitness (Table 3.). It should be noted that this study was conducted on a small sample, and the specific results are cohort specific, however, practitioners may still use this information to help inform their training programs and to ensure their athletes are going to achieve the desired outcome.

Conclusion

This is the first study to examine the dose-response relationship between a range of measures quantifying training load and changes in aerobic fitness, using a field based test of maximal aerobic speed in professional youth soccer players. Results show a *very large* relationship between time above maximal aerobic speed and changes in aerobic fitness compared to a unclear relationships with arbitrary thresholds. The practical applications provided may help practitioners to effectively plan their training programs based on the desired

training outcome. Detailed practical recommendations have been given on how linear regression analysis and standard error of the estimate can be used to help improve the training process.

Acknowledgments

Funding for this research was provided by Newcastle United Football Club and Northumbria University. The authors would like to thank Craig Musham for his help and expertise throughout the testing procedures and the players and staff at Newcastle United Football Club for their hard work and cooperation throughout the study.

Reference

1. Akubat I. Training load monitoring in soccer. In: Van Winckel J, ed. *Fitness in soccer: The science and practical application*. Klein-Gelmen: Moveo Ergo Sum; 2014:167-184.
2. Banister EW. Modeling elite athletic performance. In: MacDougall JD, Wenger HA, Green HJ, eds. *Physiological testing of elite athletes*. Champaign, Illinois: Human Kinetics; 1991:403-424.
3. Manzi V, Iellamo F, Impellizzeri F, D'ottavio S, Castagna C. Relation between individualized training impulses and performance in distance runners. *Med Sci Sports Exerc*. 2009;41(11):2090-2096.
4. Malone JJ, Di Michele R, Morgans R, Burgess D, Morton JP, Drust B. Seasonal training-load quantification in elite english premier league soccer players. *Int J Sports Physiol Perform*. 2015;10(4):489-497.
5. Owen AL, Wong DP, Paul D, Dellal A. Effects of a periodized small-sided game training intervention on physical performance in elite professional soccer. *J Strength Cond Res*. 2012;26(10):2748-2754.
6. Nédélec M, McCall A, Carling C, Legall F, Berthoin S, Dupont G. Recovery in soccer: Part 1 — post-match fatigue and time course of recovery. *Sports Med*. 2012;42(12):997-1015.
7. Gabbett TJ. The training-injury prevention paradox: Should athletes be training smarter and harder? *Br J Sports Med*. 2016:bjsports-2015-095788.
8. Impellizzeri FM, Rampinini E, Marcora SM. Physiological assessment of aerobic training in soccer. *J Sports Sci*. 2005;23(6):583-592.
9. Akubat I, Patel E, Barrett S, Abt G. Methods of monitoring the training and match load and their relationship to changes in fitness in professional youth soccer players. *J Sports Sci*. 2012;30(14):1473-1480.
10. Thorpe RT, Strudwick AJ, Buchheit M, Atkinson G, Drust B, Gregson W. Monitoring fatigue during the in-season competitive phase in elite soccer players. *Int J Sports Physiol Perform*. 2015;10(8):958-964.
11. Scott D, Lovell R. Individualisation of speed thresholds does not enhance the dose-response determination in football training. *J Sports Sci*. 2017:1-10.
12. Abt G, Lovell R. The use of individualized speed and intensity thresholds for determining the distance run at high-intensity in professional soccer. *J Sports Sci*. 2009;27(9):893-898.
13. Lucia A, Hoyos J, Santalla A, Earnest C, Chicharro JL. Tour de france versus vuelta a espana: Which is harder? *Med Sci Sports Exerc*. 2003;35(5):872-878.

14. Harley JA, Barnes CA, Portas M, Lovell R, Barrett S, Paul D, Weston M. Motion analysis of match-play in elite u12 to u16 age-group soccer players. *J Sports Sci.* 2010;28(13):1391-1397.
15. Hunter F, Bray J, Towlson C, Smith M, Barrett S, Madden J, Abt G, Lovell R. Individualisation of time-motion analysis: A method comparison and case report series. *Int J Sports Med.* 2015;36(1):41-48.
16. Weston M. Difficulties in determining the dose-response nature of competitive soccer matches. *J Athl Enhanc.* 2013;2(1):1-2.
17. Mendez-Villanueva A, Buchheit M, Simpson B, Bourdon P. Match play intensity distribution in youth soccer. *Int J Sports Med.* 2013;34(2):101-110.
18. Bundle MW, Hoyt RW, Weyand PG. High-speed running performance: A new approach to assessment and prediction. *J Appl Physiol.* 2003;95(5):1955-1962.
19. Bangsbo J. Energy demands in competitive soccer. *J Sports Sci.* 1994;12:5-12.
20. Stølen T, Chamari K, Castagna C, Wisløff U. Physiology of soccer. *Sports Med.* 2005;35(6):501-536.
21. Buchheit M, Simpson M, Al Haddad H, Bourdon P, Mendez-Villanueva A. Monitoring changes in physical performance with heart rate measures in young soccer players. *Eur J Appl Physiol.* 2012;112(2):711-723.
22. Dupont G, Akakpo K, Berthoin S. The effect of in-season, high-intensity interval training in soccer players. *J Strength Cond Res.* 2004;18(3):584-589.
23. Lorenzen C, Williams MD, Turk PS, Meehan DL, Kolsky DJC. Relationship between velocity reached at vo2max and time-trial performances in elite Australian rules footballers. *Int J Sports Physiol Perform.* 2009;4(3):408-411.
24. Bellenger CR, Fuller JT, Nelson MJ, Hartland M, Buckley JD, DeBenedictis TA. Predicting maximal aerobic speed through set distance time-trials. *Eur J Appl Physiol.* 2015;115(12):2593-2598.
25. Akenhead R, French D, Thompson KG, Hayes PR. The acceleration dependent validity and reliability of 10 hz gps. *J Sci Med Sport.* 2014;17(5):562-566.
26. Varley MC, Fairweather IH, Aughey RJ. Validity and reliability of gps for measuring instantaneous velocity during acceleration, deceleration, and constant motion. *J Sports Sci.* 2012;30(2):121-127.
27. Edwards S. *The heart rate monitor book.* New York: Polar Electro Oy; 1993.
28. Foster C, Daines E, Hector L, Snyder AC, Welsh R. Athletic performance in relation to training load. *Wis Med J.* 1996;95(6):370-374.
29. Hopkins WG. Spreadsheets for analysis of controlled trials, crossovers and time series. 2017; (sports.org/2017/wghxls.htm).

30. Hopkins WG, Marshall S, Batterham A, Hanin J. Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exerc.* 2009;41(1):3.
31. Weir JP. Quantifying test-retest reliability using the intraclass correlation coefficient and the sem. *J Strength Cond Res.* 2005;19(1):231-240.
32. Akenhead R, Nassis GP. Training load and player monitoring in high-level football: Current practice and perceptions. *Int J Sports Physiol Perform.* 2016;11(5):587-593.
33. Jones CM, Griffiths PC, Mellalieu SD. Training load and fatigue marker associations with injury and illness: A systematic review of longitudinal studies. *Sports Med.* 2017;47(5):943-974.
34. Billat VL, Flechet B, Petit B, Muriaux G, Koralsztein J-p. Interval training at vo2max: Effects on aerobic performance and overtraining markers. *Med Sci Sports Exerc.* 1999;31(1):156-163.
35. Akenhead R, Harley JA, Tweddle SP. Examining the external training load of an english premier league football team with special reference to acceleration. *J Strength Cond Res.* 2016;30(9):2424-2432.




Figure 1. Weekly (A) and daily (B) time spent above maximal aerobic speed ($t > \text{MAS}$) during the 6-week in-season training period, mean \pm SD.




Figure 2. Linear relationship between mean weekly time spent above maximal aerobic speed ($t > \text{MAS}$) and % change in MAS during the 6-week in-season training period. Intercept = -4.41 (%), Slope = 0.68 (min), SEP = 1.02 (%).

Table 1: Mean \pm SD weekly training loads throughout the 6-week training period.

	Weekly Mean \pm SD
TD (m)	29,324 \pm 4037
AD Load (m)	1477 \pm 254
m>HSD (m)	2613 \pm 576
t>HSD (min)	7.70 \pm 1.66
m>MAS (m)	2512 \pm 507
t>MAS (min)	7.41 \pm 1.72
m>VHSD (m)	940 \pm 242
t>VHSD (min)	2.35 \pm 0.58
m>30ASR (m)	770 \pm 176
t>30ASR (min)	1.95 \pm 0.48
HRE (au)	957 \pm 107
sRPE (au)	2091 \pm 380

Table 2: Relationship between mean weekly arbitrary and individualised training load measures and % changes in fitness. Pearson’s product moment correlation coefficients (r) with 90% confidence intervals (CI) and magnitude based inference of the relationship.

		m>HSD (m)	t>HSD (min)	m>VHSD (m)	t>VHSD (min)	m>MAS (m)	t>MAS (min)	m>30ASR (m)	t>30ASR (min)
MAS (km.h ⁻¹)	r	0.22	0.37	-0.07	0.05	0.50 *	0.77 *	0.20	0.62 *
	90% CI	-0.27, 0.62	-0.10, 0.71	-0.51, 0.40	-0.42, 0.50	0.06, 0.78	0.48, 0.91	-0.28, 0.61	0.22, 0.84
	MBI	Unclear	Unclear	Unclear	Unclear	Possibly Large	Possibly Very Large	Unclear	Possibly Large
MSS (km.h ⁻¹)	r	0.32	0.34	0.25	0.27	0.30	0.21	-0.09	-0.15
	90% CI	-0.17, 0.68	-0.15, 0.69	-0.24, 0.64	-0.22, 0.65	-0.18, 0.67	-0.28, 0.61	-0.53, 0.39	-0.57, 0.33
	MBI	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear

Abbreviations; m>HSD; distance > 17 km.h⁻¹, t>HSD; time > 17 km.h⁻¹, m>VHSD; distance > 21 km.h⁻¹, t>VHSD; time > 21 km.h⁻¹, m>MAS; distance > MAS km.h⁻¹, t>MAS; time > MAS km.h⁻¹, m>30ASR; distance > 30% ASR km.h⁻¹, t>30ASR; time > 30% ASR km.h⁻¹.

* Substantial relationship (90% CI does not overlap both the positive and negative thresholds by ≥5% [Hopkins et al. 2009])

Table 3: Relationship between mean weekly arbitrary and individualised training load measures and % change in fitness. Linear regression coefficient of determination (R^2), slope, intercept and standard error of prediction (SEP). Also displayed is the minimum training load (TL) required to elicit the minimum detectable change (MDC) in fitness.

		m>HSD (m)	t>HSD (min)	m>VHSD (m)	t>VHSD (min)	m>MAS (m)	t>MAS (min)	m>30ASR (m)	t>30ASR (min)
MAS (km.h ⁻¹)	R ²	0.05	0.14	0.00	0.00	0.25	0.59	0.04	0.38
	Slope (%)	0.58	0.34	-0.44	0.13	1.52	0.68	1.77	1.96
	Intercept (%)	-0.86	-1.99	1.07	0.35	-3.17	-4.41	-0.71	-3.16
	SEP (%)	1.55	1.48	1.59	1.59	1.37	1.02	1.56	1.25
	TL to elicit MDC	3672	9.48	-449	7.16	2914	8.30	1116	2.26
MSS (km.h ⁻¹)	R ²	0.10	0.11	0.06	0.07	0.09	0.04	0.01	0.02
	Slope (%)	0.74	0.27	1.40	0.63	0.80	0.16	-0.67	-0.43
	Intercept (%)	-1.13	-1.29	-0.51	-0.68	-1.21	-0.41	1.32	1.64
	SEP (%)	1.33	1.32	1.36	1.35	1.34	1.37	1.40	1.39
	TL to elicit MDC	3820	10.97	1580	3.77	3626	12.86	-565	-0.14

Abbreviations; m>HSD; distance > 17 km.h⁻¹, t>HSD; time > 17 km.h⁻¹, m>VHSD; distance > 21 km.h⁻¹, t>VHSD; time > 21 km.h⁻¹, m>MAS; distance > MAS km.h⁻¹, t>MAS; time > MAS km.h⁻¹, m>ASR; distance > 30% ASR km.h⁻¹, t>ASR; time > 30% ASR km.h⁻¹.

Note: Slope for distance measures is unit increase per 1000 metres covered