

Northumbria Research Link

Citation: Scott, Jason (2015) Delivering clinical case management for frequent callers to ambulance services. *Emergency Medicine Journal*, 32 (5). p. 342. ISSN 1472-0205

Published by: BMJ Publishing Group

URL: <https://doi.org/10.1136/emered-2014-204354> <<https://doi.org/10.1136/emered-2014-204354>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/37502/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



**Northumbria
University**
NEWCASTLE



UniversityLibrary

Delivering clinical case management for frequent callers to ambulance services

Jason Scott¹

¹Institute of Health and Society, Newcastle University

Author Details:

Dr Jason Scott
Institute of Health and Society
Newcastle University
Baddiley-Clark Building,
Richardson Road,
Newcastle upon Tyne,
NE2 4AX.

Email: jason.scott@ncl.ac.uk

Telephone: 0191 208 8848

Word Count: 588

Ambulance services in England are playing an increasing role in the provision of care. Their role has changed from one of picking up patients in the community and dropping them off at an emergency department (ED), to more advanced care models. These now include the provision of clinical advice over the telephone (hear and treat), the treatment of patients on-scene (see and treat), and the more traditional conveyance to the ED.[1] These changes are occurring during a prolonged period of increased demand on all areas of emergency care, including a 47% rise in ED attendance over 15 years.[2] Further, it is known that frequent users of healthcare use a disproportionate amount of resources.[3]

Clinical Case Management for Frequent Callers

The paper by Edwards et al. [include paper reference here] provides one of the first examples of delivering interventions for frequent callers to an English ambulance service, and is one of few publications on the topic internationally.[4] There are three notable findings that will be discussed in this commentary:

- (1) Frequent callers were found to be heterogeneous with 95 (86%) falling within more than category;
- (2) Ten different types of interventions were used, although 90 (82%) patients received five or fewer;
- (3) The interventions as a whole showed a reduction in median calls from five per month pre-intervention to zero per month post-intervention.

These findings are of importance for both research and practice. Firstly, the heterogeneous profile of patients within this study suggests that, similar to frequent users of the ED,[5] there are potential subgroups of frequent callers. However there is still much research to be done before they can be fully identified due to the paucity of research within ambulance services.[4]

Secondly, that there were ten intervention types demonstrates the complexity of care associated with this group of patients who may require the support of various health and social care services. Whilst this potentially serves as a note of caution, complexity should be sought to be understood rather than avoided. Indeed this could serve as an opportunity for ambulance services, which have

been recognised as being able to coordinate and lead the integration of health and social care services,[6] which coincides with perspectives on transforming the health care system for the delivery of more integrated care.[7]

Thirdly the reduction in calls is a promising indication that, from an ambulance service perspective, providing interventions can reduce service usage and possibly free up resources that are needed elsewhere. Nevertheless, caution should be noted when interpreting the impact of clinical case management. As Edwards et al.[include paper reference here] have acknowledged, the lack of a control group means that reductions in usage cannot be fully attributed to the interventions. Roland and Abel[8] have recently cautioned against ignoring regression to the mean; given that patients were registered for up to 24.5 months at the maximum end of the range, it is entirely possible that this contributed to the result. Further research would provide an insight into the extent that regression to the mean has impacted on the study outcome.

What does quality mean to frequent callers?

There is a risk that frequent callers, once identified by ambulance services, are simply passed to another service[9] so that budgetary rather than patient needs are met. As such it must be remembered that many patients will be frequently calling ambulance services because they are not receiving adequate care, as opposed to calling for vexatious reasons.[5] Qualitative research is needed to identify outcomes that are of importance to frequent callers and to understand the mechanisms by which interventions for frequent callers work.

References

1. Department of Health. *Taking healthcare to the patient: Transforming NHS Ambulance Services*. London: Department of Health, 2005.
2. National Audit Office. *Emergency admissions to hospital: managing the demand*. London: The Stationery Office, 2013.
3. Pines JM, Asplin BR, Kaji AH, et al. Frequent Users of Emergency Department Services: Gaps in Knowledge and a Proposed Research Agenda. *Academic Emergency Medicine* 2011;**18**(6):e64-9 doi: 10.1111/j.1553-2712.2011.01086.x.
4. Scott J, Strickland AP, Warner K, et al. Frequent callers to and users of emergency medical systems: A systematic review. *Emergency Medicine Journal* 2013 doi: 10.1136/emermed-2013-202545.
5. LaCalle E, Rabin E. Frequent Users of Emergency Departments: The Myths, the Data, and the Policy Implications. *Annals of Emergency Medicine* 2010;**56**(1):42-48
6. House of Commons Health Committee. *Urgent and Emergency Services: Second Report of Session 2013–14*. London: The Stationery Office, 2013.
7. Naylor C, Imison C, Addicott R, et al. *Transforming our health care system*. London: The King's Fund, 2013.
8. Roland M, Abel G. Reducing emergency admissions: are we on the right track? *BMJ* 2012;**345**:e6017 doi: 10.1136/bmj.e6017.
9. Mason SM. Frequent attendance at the emergency department is a symptom but not a disease. *Emergency Medicine Journal* 2014 doi: 10.1136/emermed-2014-203674.