

Northumbria Research Link

Citation: Menting, Stein Gerrit Paul, Elferink-Gemser, Marije Titia, Huijgen, Catharina and Hettinga, Florentina (2019) Pacing in lane-based head-to-head competitions: A systematic review on swimming. *Journal of Sports Sciences*, 37 (20). pp. 2287-2299. ISSN 0264-0414

Published by: Taylor & Francis

URL: <https://doi.org/10.1080/02640414.2019.1627989>
<<https://doi.org/10.1080/02640414.2019.1627989>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/37917/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

1 **Pacing in lane-based head-to-head competitions: A systematic review**
2 **on swimming**

3 Stein Gerrit Paul Menting^{1,2,3}, Marije Titia Elferink-Gemser¹, Barbara
4 Catharina Huijgen¹, Florentina Johanna Hettinga^{2,3*}

5 *1. Center for Human Movement Sciences, University Medical Center Groningen,*
6 *University of Groningen, Groningen, the Netherlands.*

7 *2. School of Sport, Rehabilitation and Exercise Sciences, University of Essex,*
8 *Colchester, the United Kingdom.*

9 *3. Department of Sport, Exercise & Rehabilitation, Faculty of Health and Life*
10 *Sciences, Northumbria University, Newcastle, United Kingdom*

11

12 * Correspondence:

13 Department of Sport, Exercise & Rehabilitation

14 Faculty of Health and Life Sciences

15 Northumbria University

16 Room 238, Northumberland Building

17 Newcastle Upon Tyne

18 NE1 8ST

19 United Kingdom

20 E-mail: florentina.hettinga@northumbria.ac.uk;

21 Tel: +44 (0)1912273989

22

23

24

25

26 Pacing in lane-based head-to-head competitions: A systematic review **27 on swimming**

28 Athletes' energy distribution over a race (e.g. pacing behaviour) varies across different sports.
29 Swimming is a head-to-head sport with unique characteristics, such as propulsion through
30 water, a multitude of swimming stroke types and lane-based racing. The aim of this paper was
31 to review the existing literature on pacing behaviour in swimming. According to PRISMA
32 guidelines, 279 articles were extracted using the PubMed and Web of Science databases. After
33 the exclusion process was conducted, 16 studies remained. The findings of these studies
34 indicate that pacing behaviour is influenced by the race distance and stroke type. Pacing
35 behaviours in swimming and time-trial sports share numerous common characteristics. This
36 commonality can most likely be attributed to the lane-based racing set-up. The low efficiency
37 of swimming resulting from propulsion through the water induces a rapid accumulation of
38 blood lactate, prompting a change in swimmers' biomechanical characteristics, with the goal
39 of minimising changes in velocity throughout the race. Although the literature on youth
40 swimmers is scarce, youth swimmers demonstrate more variable pacing profiles and have
41 more difficulty in selecting the most beneficial energy distribution.

42

43 Keywords: pacing behaviour; swimming; athletic performance; psychology; adolescent; talent
44 development.

45 **Introduction**

46 Pacing behaviour can be defined as the outcome of an individual's continuous, goal-
47 directed, decision-making process regarding the distribution of energy resources over
48 time (Edwards & Polman, 2013; Smits, Pepping, & Hettinga, 2014). In head-to-head
49 and time-trial sports environments, the goal of the pacing process is to achieve optimal
50 performance, which requires that athletes deplete all possible energy stores prior to
51 finishing the race, but not so fast that a meaningful slowdown occurs before the end of
52 the race (Foster et al., 2003; Ulmer, 1996). The application of a broad range of
53 theoretical models and the findings of experimental studies have shown that pacing
54 behaviour is primarily influenced by the duration of the competitive event (Foster et
55 al., 2003; Ulmer, 1996; van Ingen Schenau & Cavanagh, 1990). In addition, recent
56 studies have shown that different competitive environments influence pacing
57 behaviour (Hettinga, Konings, & Pepping, 2017; Konings & Hettinga, 2017). Lastly,
58 in the finals of elite competitions in multiple sports, the pacing behaviour of more
59 successful performers seems to differ from of less successful performers (Konings,
60 Noorbergen, Parry, & Hettinga, 2016; Muehlbauer, Schindler, & Panzer, 2010).

61 The representation of an athletes' pacing behaviour over a race is termed
62 'pacing profile'. Although general pacing profiles have been distinguished (Abbiss &
63 Laursen, 2008), it is assumed that pacing behaviour is associated with the different
64 biomechanical and physiological limitations of the athlete (Stoter et al., 2016) as well
65 as with the different competitive environments (De Koning et al., 2011; Konings &
66 Hettinga, 2017) the athlete competes in.

67 Although the cognitive skills necessary for an inherent pacing ability are
68 apparently present in young children (Micklewright et al., 2012), the brain areas
69 associated with pacing behaviour continue to develop throughout adolescence

70 (Edwards & Polman, 2013; Elferink-Gemser & Hettinga, 2017; Giedd et al., 1999;
71 Rubia et al., 2001). Studies have shown that pacing behaviour develops during
72 adolescence in elite athletes (Menting, Konings, Elferink-Gemser, & Hettinga, 2018;
73 Wiersma, Stoter, Visscher, Hettinga, & Elferink-Gemser, 2017). Moreover, adolescent
74 athletes whose pacing profiles resemble profiles of adult elite performers earlier on in
75 their development, seem to achieve a higher performance level in their later career
76 compared to their peers (Wiersma et al., 2017). Therefore, an exploration of how youth
77 athletes pace their races and develop their pacing behaviour throughout adolescence is
78 particularly salient.

79

80 Swimming is a head-to-head sport entailing a unique combination of characteristics.
81 Firstly, swimmers propel themselves through water, which requires more energy than
82 overcoming air resistance during running or cycling races (Toussaint, 1990; Toussaint
83 et al., 1988). Because of the extensive energy loss to the environment, it is essential
84 for swimmers to reduce drag and to optimise propulsion (Barbosa et al., 2010; Holmér,
85 1974). Increased propulsion can be achieved by increasing the number of strokes for a
86 given distance, defined as the stroke rate (SR), or by increasing the distance covered
87 per stroke, namely the stroke length (SL). Due to the propulsion through water, an
88 increase in SR will induce an increase in drag and, therefore, an increase in the amount
89 of energy lost to the environment (Barbosa et al., 2010). Hence, elite swimmers mostly
90 increase SL and reduce drag compared with non-elite swimmers (Barbosa et al., 2010).
91 It has been posited that to ensure an optimally paced race, a swimmer should minimise
92 fluctuations in velocity throughout the race, thereby minimising energy loss to the
93 environment in the form of drag (Barbosa et al., 2010; De Koning et al., 2011).
94 Moreover, a key phase of the race is the underwater phase that follows the start and

95 turns. During this phase, the highest race velocity is achieved due to the increased
96 impulse following the dive or push off from the wall and the decrease in drag as a
97 result of the adoption of a streamlined body position (Hochstein & Blickhan, 2014;
98 Vantorre, Chollet, & Seifert, 2014).

99 Swimming entails several different stroke types and various race lengths, each
100 associated with a specific technical skillset and energetic demand (Barbosa et al., 2006;
101 Capelli, Pendergast, & Termin, 1998; Zamparo et al., 2005). The race distance in a
102 pool ranges from 50 m to 200 m for the breaststroke, backstroke and butterfly events
103 and up to 1,500 m for freestyle races (FINA, 2017). In open water, races can range
104 from 5 to 25 km (Swimming World Magazine, 2017). Moreover, pool swimming
105 competitions are generally organised as a qualifying structure comprising heats, semi-
106 finals and finals.

107 A final characteristic is that during pool swimming events, the competitors are
108 separated by lanes. Consequently, competitors do not have to compete to be positioned
109 in the ideal line, as is common in other head-to-head competitions such as (track-)
110 cycling, running, short-track speed skating or Boat Race rowing.

111

112 Because of the unique combination of characteristic relevant to the sport of swimming,
113 the pacing behaviour in swimming could deviate from those of other sports. The
114 present review is aimed at offering insights into sport specific pacing behaviour in
115 swimming. The primary aim is to provide an overview of studies on this subject. As
116 there is a wide range of distances covered in swimming events, each of which entails
117 particular energetics and techniques, it was decided to focus on 100–800 m pool races.
118 The durations of these events (the world records for the 100 m and 800 m freestyle
119 races are 46.91 s and 452.12 s, respectively (FINA, 2017)) best match those of other

120 sports, such as track cycling as well as short- and long-track speed skating, as described
121 in the literature (Hettinga, De Koning, & Foster, 2009; Konings et al., 2016;
122 Muehlbauer, Schindler, & Panzer, 2010; Stoter et al., 2016; van Ingen Schenau, De
123 Koning, & De Groot, 1992). In addition to providing an overview of the literature,
124 potential factors that influence the pacing behaviour of swimmers were identified and
125 discussed. As adolescence is a crucial phase of pacing behaviour development, a
126 particular focus of the review is on studies that explore the pacing strategies of youth
127 swimmers, namely juniors (aged 12–16 years) as well as adolescent swimmers (aged
128 16–21 years).

129

130 **Methods**

131 Following PRISMA guidelines, the PubMed and Web of Science databases were
132 searched for studies about pacing behaviour in swimming up to until April 2017 using
133 the following combination of terms:

134 (1) Pacing (OR performance strategy* OR energy distribution* OR pacing
135 behaviour* OR velocity profile)

136 AND

137 (2) Swim*

138 NOT

139 (3) Triathlon* OR Animal* OR Fish* OR Pacemaker* OR Bacter*

140

141 The inclusion terms focused on articles written in English and published in peer
142 reviewed journals, covering pacing behaviour in swimming in relation to performance.
143 Therefore, all included articles described pacing profiles with outcome variables such
144 as lap times or (normalised) velocity distribution over the race. Additionally, the

145 variability of pacing profiles over multiple races, expressed as the coefficient of
146 variation (CV), was analysed in several studies. To provide an extensive overview of
147 the literature, included were articles featuring participants of all age groups and
148 performance levels. The initial search yielded 279 articles. After duplicate studies had
149 been discarded, a total of 244 articles remained. The titles and abstracts of the
150 remaining articles were read and papers lacking relevant links to pacing behaviour in
151 swimming were excluded, resulting in 22 potential papers. After reading the bodies of
152 these remaining articles, six were excluded because the articles did not meet the
153 inclusion criteria. Therefore, a total of 16 studies were reviewed (Figure 1). Quality
154 assessment of the articles was performed following guidelines provided by Letts et al.
155 (2007). Articles with a score above seven were considered of good methodological
156 quality.

157

158 *** Please insert Figure 1 near here***

159

160 Pacing profiles have been described in previous studies using velocity expressed as a
161 percentage of the mean velocity in the race (e.g., ‘normalised velocity’). This method
162 provides a way of comparing the profiles of participants whose performance levels,
163 sex and age differ. To avoid any misinterpretation in the description of pacing profiles,
164 the definitions of general pacing profiles provided by Abbiss and Laursen (2008) and
165 adapted for swimming by Mauger, Neuloh, & Castle (2012) were used in the current
166 review. In a negative pacing profile, the velocity increases throughout the race. By
167 contrast, velocity decreases in a positive pacing profile. In an even pacing profile, the
168 velocity remains constant throughout the race. In a parabolic shaped pacing profile,
169 the velocity decreases after the initial phase of the race and subsequently increases in

170 the final phase. Finally, the fast-start-even pacing profile is characterised by a high
171 velocity in the initial phase, followed by a lower, constant velocity during the
172 remainder of the race. To the authors' knowledge, there are no specific percentages
173 determined in the literature whereby these pacing profiles can be quantified.

174 As the qualification of participant performance level varied throughout the
175 different included articles, there was a need for a standard qualification system to
176 properly compare the outcomes reported in the included articles. Therefore,
177 performance levels were categorised based on the world record in the year of
178 publication of the article. Participants were divided into three groups: elite, sub-elite
179 and competitive. Elite swimmers were defined as those with performances within
180 110% of the world record (Vantorre, Chollet, & Seifert, 2014). Sub-elite swimmers
181 were defined when those whose total race time was 110–120% of the world record.
182 Finally, competitive swimmers were defined as swimmers who performed in a
183 competitive environment but whose total race time exceeded 120% of the world
184 record.

185

186 **Results**

187 *General pacing profiles*

188 All of the reviewed articles ($n = 16$) were of good methodological quality (with total
189 scores ≥ 7 ; Table 1). Therefore, studies were not distinguished based on qualitative
190 weight. Table 2 presents a summary of the characteristics and outcomes reported in
191 the reviewed articles. In the majority of studies, the participants were elite ($n = 9$),
192 followed by sub-elite ($n = 4$) and competitive swimmers ($n = 3$). Most of the studies
193 analysed freestyle swimming ($n = 13$), followed by breaststroke ($n = 5$), backstroke (n
194 $= 3$) and butterfly ($n = 3$). The pool lengths were 25 m ($n = 3$), 50 m ($n = 11$) and not

195 specified in two studies. The pacing profiles identified in the studies were positive (n
196 = 11), negative (n = 2), even (n = 3), parabolic (n = 8) and fast-start-even (n = 8). In a
197 majority of the articles (n = 11), pacing profiles were analysed using data collected
198 during actual swimming competitions (e.g. 'real competition'). The articles which
199 collected data in real competition analysed races from either a combination of the
200 heats, semi-finals and finals (n = 6), only the semi-finals and finals (n = 3) or
201 exclusively the finals (n = 2). Additionally, several studies were conducted in more
202 controlled settings (e.g., 'simulated competition') in which participants were tasked
203 with swimming a time-trial without an opponent (n = 6). One article explored data
204 collected during real and simulated competition scenarios entailing one or two
205 opponents. No significant difference was found between pacing profiles in simulated
206 competitions (with an opponent) and in real competitions ($P > 0.22$). However, in real
207 competitions ($P < 0.001$), absolute velocity was higher during all sections of the race
208 (Skorski, Faude, Rausch, & Meyer, 2013). Three of the studies conducted in a
209 simulated competition examined the effect of an imposed manipulation of swimmers'
210 pacing behaviours on their performance outcomes. Only one study of swimmers in real
211 competition related observed pacing profiles to total race time.

212 A total of 12 of the 16 reviewed studies showed a higher velocity in the starting
213 phase of the race. This phenomenon was observed for all four distances (100 m: n = 3,
214 200 m: n = 7, 400 m: n = 8, 800 m: n = 3) and for all stroke types. Pacing profiles for
215 100 m and 200 m races showed a high velocity during the first 50 m (Dormehl &
216 Osborough, 2015; Nikolaidis & Knechtle, 2017; Robertson, Pyne, Hopkins, & Anson,
217 2009; Skorski, Faude, Caviezel, & Meyer, 2014; Veiga & Roig, 2016). For the 400 m
218 profile, a high velocity either occurred during the first 50 m (Mytton et al., 2015;
219 Skorski et al, 2014b) or 100 m (Robertson, et al., 2009). In the 800 m freestyle races,

220 a high velocity was reported for the initial 100 m freestyle (Nikolaidis & Knechtle,
221 2017; Skorski, Faude, Rausch, & Meyer, 2013). The high velocity during the starting
222 phase was reported in several studies that excluded the first 15 m of the race in the
223 velocity measurements (Dormehl & Osborough, 2015; Mauger et al., 2012).
224 Correspondingly, it was reported in studies in which swimmers were instructed to start
225 from the water (Figueiredo, Zamparo, Sousa, Vilas-Boas, & Fernandes, 2011;
226 Schnitzler, Seifert, & Chollet, 2009). Thus, it can be stated that the high velocity during
227 the starting phase occurs independently of the dive start.

228

229 ***Please insert Table 1 near here***

230

231 ***Race distance***

232 Different features in the pacing profile were observed depending on the race distance.
233 In all three studies of 100 m races, it was observed that the pacing profiles of
234 swimmers, both elite and competitive, were positive (Dormehl & Osborough, 2015;
235 Nikolaidis & Knechtle, 2017; Robertson et al., 2009). The pacing profiles of swimmers
236 in races over 200 m were analysed in 10 studies. In studies that focused on 200 m
237 competitions, both real and simulated, in 25 m and 50 m pools, elite swimmers showed
238 a high velocity start followed by a large decrease in velocity during the second lap, a
239 small decrease in velocity during the third lap and a constant velocity up to the end of
240 the race (Figueiredo et al., 2011; Skorski et al., 2014b; Veiga & Roig, 2016). In one
241 study that investigated competitive swimmers performing 200 m freestyle, the velocity
242 increased in the final lap (2.1%) (Nikolaidis & Knechtle, 2017). The pacing profiles
243 of swimmers in 400 m races were examined in a total of 10 studies. Elite swimmers
244 performing freestyle and medley in real competitions displayed parabolic pacing

245 profiles, with significantly higher velocities during the first and last sections than in
246 other sections of the race ($P < 0.001$) (Mytton et al., 2015; Robertson et al., 2009;
247 Saavedra, Escalante, Garcia-Hermoso, Arellano, & Navarro, 2012; Skorski et al.,
248 2014b). Swimmers with parabolic pacing profiles performed significantly better than
249 the swimmers who displayed one of the other pacing profiles. This finding applied to
250 both males ($P < 0.001$) and females ($P < 0.001$) (Taylor, Santi, & Mellalieu, 2016).
251 Two studies found that during 400m races in real competitions, the most common
252 pacing profiles among elite swimmers performing freestyle were the fast-start-even
253 and parabolic profiles (Mauger et al., 2012; Taylor et al., 2016). Three studies
254 examined the pacing profiles of swimmers performing in 800 m races. A first section
255 at a fast pace, followed by a gradual decrease in the normalised velocity during the
256 200–700 m and increased normalised velocity during the final 100 m was observed
257 among adolescent sub-elite swimmers (Skorski et al., 2013). A study of in elite female
258 swimmers performing freestyle during real competition confirmed these
259 characteristics, reporting a gradual decrease in velocity throughout the race, with the
260 slowest lap time for eleventh lap (500–550 m) (Lipinska, Allen, & Hopkins, 2016).
261 Among competitive freestyle swimmers, split times increased during the 100–200 m
262 (8.8%) and 200–600m sections (0.2% – 1.0%) and decreased during the 600–700m
263 (0.3%) and 700–800m sections (3.4%) ($P < 0.001$) (Nikolaidis & Knechtle, 2017).

264

265 *Stroke types*

266 In addition to the duration of the race, certain deviations in pacing behaviour were
267 caused by the different stroke types. Elite swimmers in 200 m freestyle, butterfly and
268 backstroke races tended to display a fast-even profile, with a fast first 50 m section for
269 all three stroke types ($P < .001$ for all other sections). Additionally, in the freestyle and

backstroke events, the second 50 m lap was also faster than the third and fourth laps ($P < 0.001$) (Skorski et al., 2014b). The pacing profile of swimmers performing breaststroke was characterised by a high velocity during the first 50 m lap and a gradual decrease in normalised velocity with every 50 m lap ($P < 0.001$) (Skorski et al., 2014b; Thompson, MacLaren, Lees, & Atkinson, 2003, 2004). Furthermore, more variability was observed in the pacing profiles of swimmers performing breaststroke during the entire race (Skorski et al., 2014b). Individual medley events were examined in two studies. Elite swimmers participating in 200 m and 400 m individual medley real competitions demonstrated a parabolic pacing profile in which they performed butterfly strokes for the smallest percentage of the total race time, followed by freestyle, backstroke and breaststroke (Robertson et al., 2009; Saavedra et al., 2012).

281

282 ***Biomechanics and metabolic systems***

The metabolic systems used in swimming competition were described in three studies (Figueiredo et al., 2011; Thompson et al., 2003; Thompson et al., 2004). One study that investigated a 200 m freestyle race reported that the percentage of the total metabolic power output covered by the aerobic energy system increased over the duration of the race, from 45% during the first lap to 83% in the third lap, with a drop to 66% during the final lap. Conversely, the coverage of the anaerobic system decreased over the duration of the race, from 55% during the first lap to 17% during the third lap, with a small increase to 24% during the final lap (Figueiredo et al., 2011). It was reported that as the race time increased, the blood lactate peak value correspondingly increased, which was linked to increasing fatigue throughout the race (Figueiredo et al., 2011; Thompson et al., 2003; Thompson et al., 2004). In all four studies that measured biomechanical characteristics in adult swimmers, the SL

295 decreased throughout the race. One study on sub-elite swimmers performing freestyle
296 in a 400 m race showed a drop in SL after the first 50 m and during the last 100 m,
297 whereas the SR remained unchanged during the race (Schnitzler, Seifert, & Chollet,
298 2009). However, the other three studies reported a decrease in SL accompanied by an
299 increase in SR (Figueiredo et al., 2011; Thompson et al., 2003; Thompson et al., 2004).
300 Additionally, swimming performance was subdivided into surface and underwater
301 swimming in one study, reporting that although the velocity in surface swimming
302 decreased by 6–8% over a 200 m freestyle race, the underwater velocity remained
303 constant (Veiga & Roig, 2016).

304

305 *Medallists vs non-medallists*

306 The pacing behaviour of swimmers in the finals of elite competitions were compared
307 in three studies. Two studies found that the pacing behaviour expressed in lap times,
308 and therefore representing absolute velocity, was similar for swimmers ranked in first
309 to sixteenth (Robertson et al., 2009) or first to eighth (Mytton et al., 2015) place.
310 However, a comparison of the normalised velocity showed that medallists had a
311 relatively lower normalised velocity in both the first 100 m ($102.2 \pm 1.2\%$ vs $103.1 \pm$
312 1.1% , $P = 0.03$) and the second 100 m ($97.7 \pm 0.8\%$ vs $98.2 \pm 0.6\%$, $P < 0.001$)
313 compared to swimmers ranked fourth to eighth place. In the third 100 m, there was no
314 difference between medallists and non-medallists ($98.5 \pm 1.0\%$ vs $98.4 \pm 0.6\%$, $p =$
315 0.63). In the final 100 m, medallists had a higher normalised velocity compared to non-
316 medallists ($101.8 \pm 1.7\%$ vs $100.5 \pm 1.2\%$, $P \leq 0.01$) (Mytton et al., 2015). Among
317 elite swimmers performing a 200 m medley, it was observed that medallists had a
318 higher absolute velocity than non-medallists (fourth to sixteenth place) during
319 throughout the race. However, medallists invested more time in butterfly and freestyle

320 strokes ($P < 0.001$) and less in backstroke ($P < 0.001$) and breaststroke ($P < 0.021$)
321 than swimmers ranked in ninth to sixteenth place (Saavedra et al., 2012). In the 400 m
322 medley, medallists invested more time in butterfly strokes ($P < 0.001$) and less in
323 backstroke ($P = 0.018$) and breaststroke ($P = 0.024$) compared with swimmers ranked
324 in ninth to sixteenth place (Saavedra et al., 2012).

325

326 *Pacing in youth swimmers*

327 Three studies focused on the pacing behaviours of youth swimmers. One study found
328 no difference in the pacing profiles of young and adolescent competitive swimmers
329 (group 1: aged 14.4 ± 0.7 years; group 2: aged $17.0 \pm .8$ years) performing in 200 m
330 freestyle real competitions (Dormehl & Osborough, 2015). The pacing profile
331 observed in this study corresponds to the profile displayed by elite swimmers
332 competing in the same event (Skorski et al., 2014b; Veiga & Roig, 2016). A
333 comparison of adolescent sub-elite swimmers (aged 16.9 ± 2.1 years) with elite
334 swimmers (aged 22.8 ± 2.9 years) participating in 200 m and 400 m freestyle races
335 revealed that the variability of the pacing profiles of both elite and adolescent
336 swimmers was low throughout the race. However, in the last quarter of the race, the
337 variability was higher among adolescent swimmers than among elite swimmers
338 (Skorski et al., 2014b; Skorski et al., 2013). Furthermore, the findings of a study of
339 sub-elite youth swimmers (males: 19.2 ± 2.0 years, females: 16.2 ± 1.8 years)
340 participating in a 400 m freestyle race, revealed better performances of seven out of
341 15 swimmers in a trial with an imposed manipulated pacing profile compared with
342 performances in trials entailing a self-regulated pace (Skorski et al., 2014a).

343

344 *** Please insert Table 2 near here***

345

346 **Discussion**

347 Pacing behaviour in swimming is characterised by a high velocity start and is
348 influenced by racing distance. The study findings indicate that when the racing
349 distance increases, the swimmers' pacing profiles change from being positive (100 m
350 races) to being more parabolic (400 m and 800 m races). In elite finals, the best
351 performing swimmers demonstrated a higher absolute velocity throughout the race
352 (Mytton et al., 2015; Robertson et al., 2009; Saavedra et al., 2012). However, the
353 pacing profiles of the top three performers differed from those of the other finalists
354 (Mytton et al., 2015; Saavedra et al., 2012). Namely, medallists showed a lower
355 normalised velocity during the first half of the race and a higher normalised velocity
356 during the last portion of the race (Mytton et al., 2015). Notably, all of these
357 characteristics are similar to those reported in studies of time-trial competitions (Foster
358 et al., 2003; Foster et al., 2004; Hettinga et al., 2009; Muehlbauer Schindler, & Panzer,
359 2010; Stoter et al., 2016; Ulmer, 1996; Wiersma, Stoter, Visscher, Hettinga, &
360 Elferink-Gemser, 2017). Swimming is a head-to-head competition, in which the
361 winner is the athlete who covers the given race distance first, regardless of the time
362 taken. Nevertheless, distinct differences between athletes' pacing behaviours were
363 observed in 400 m swimming and 1,500 m running competitions, although both sports
364 entail head-to-head competition of similar duration (Mytton et al., 2015). Additionally,
365 the characteristics of the pacing profile in swimming are similar to those of athletes in
366 time-trial sports. This similarity is most likely caused by the separation of competitors
367 through the use of lanes, thereby preventing tactical behaviour as seen in classic head-
368 to-head sports (e.g., drafting behind an opponent), which enables a swimmer to be
369 more independent of other competitors. This explanation is supported by the fact that

370 studies on rowing, another head-to-head sport in which competitors are separated by
371 lanes, have reported pacing behaviour which resembles pacing profiles of time-trial
372 sports (Garland, 2005; Muehlbauer, Schindler, & Widmer, 2010).

373

374 The different strokes types are a distinctive feature of pool swimming. There appear
375 to be marked differences in swimmers' pacing profiles associated with different stroke
376 types (Skorski et al., 2014b; Veiga & Roig, 2016), indicating that stroke type affects
377 pacing behaviour. Most notably, the pacing profiles of swimmers performing
378 breaststroke, in contrast to other strokes, were characteristically positive (Skorski et
379 al., 2014b; Thompson et al., 2003; Thompson et al., 2004). In addition, in races, the
380 variability of pacing profiles was higher for swimmers performing the breaststroke
381 than that for swimmers performing other strokes (Skorski et al., 2014b). A possible
382 explanation could be found in the finding that the breaststroke technique features a
383 large intracyclic variation of swimming velocity (Barbosa et al., 2006). Higher
384 intracyclic variations in velocity prompt more mechanical work by swimmers and
385 consequently induce greater energy expenditure (Barbosa et al., 2006). This increased
386 energy expenditure could be the reason for the decrease in swimming velocity in the
387 last lap as well as the increased variation throughout the race.

388

389 A comparison of contribution of energy systems in the course of a swimming race to
390 a track cycling task of a similar duration (141.30 ± 4.47 s for swimming vs 133.8 ± 6.6
391 s for cycling) reveals a clear difference between the two sports (Figueiredo et al., 2011;
392 Foster et al., 2004). The contribution of the anaerobic system during swimming is
393 around 56% after the first 50 m, thereafter decreasing with a corresponding increase
394 in the aerobic contribution during the race, which reaches a high point of 83% during

395 the third lap (Figueiredo et al., 2011). In track cycling, the contribution of the anaerobic
396 energetic system is around 75% during the first 30 seconds (Foster et al., 2004), which
397 is comparable to the first 50 m in swimming. The aerobic system only takes over as
398 the predominant energy system at the 100 s mark (Foster et al., 2004). This difference
399 in the contributions of the two energetic systems could be attributed to low efficiency
400 in swimming caused by the increased energy loss to the environment. This low
401 efficiency could place a greater demand on the anaerobic system to maintain velocity.
402 Consequently, the accumulation of blood lactate, and in association symptoms of
403 fatigue, occurs earlier during a swimming event than in a track cycling event of the
404 same duration. This relatively fast onset of blood lactate accumulation is also reflected
405 in biomechanical characteristics. As blood lactate level increases over the duration of
406 the race, SL tends to decrease (Schnitzler et al., 2009, Figueiredo et al., 2011;
407 Thompson et al., 2003, 2004). However, as noted in previous studies, it is essential to
408 minimise large variations in velocity throughout the race (Barbosa et al., 2010; De
409 Koning et al., 2011). Therefore, to maintain velocity, swimmers must increase SR
410 during the race. Notably, a high SR is associated with a higher level of drag than a
411 high SL and a low SR.

412 Additionally, it appears that whereas elite swimmers maintain underwater
413 velocity during the race, surface velocity decreases (Veiga & Roig, 2016). This finding
414 accords with the previously mentioned goals of minimising drag and maintaining
415 velocity throughout the race. As for the underwater phase of the lap, drag is minimised
416 through the streamlined body position. Consequently, the highest velocity is achieved
417 during this phase of the race. A recent study that examined behavioural differences in
418 pacing between and within the laps of 32 elite swimmers confirmed the occurrence of
419 changes in biomechanical characteristics resulting from increasing fatigue as well as

420 the maintenance of constant underwater velocity throughout the race (Simbaña-
421 Escobar, Hellard, Pyne, & Seifert, 2017). This study concluded that swimmers' pacing
422 profiles within the first lap evidenced a decreasing velocity because of the loss of
423 velocity following the dive. The dive is the fastest part of the race because of the initial
424 acceleration as well as the airborne locomotion, compared with the rest of the race in
425 which locomotion occurs in water (Vantorre et al., 2014). Additionally, the swimmers'
426 pacing behaviour within the second and third laps is characterised by a decrease in
427 velocity at the end of the lap as they prepare to turn and by an increase of velocity
428 during the underwater phase attributed to decreased drag.

429

430 Because of the scarce literature on youth swimmers' pacing behaviour ($n = 3$), it is
431 difficult to provide a detailed description of the pacing behaviour of junior and
432 adolescent swimmers. No direct differences in the pacing profiles of youth and adult
433 swimmers were found. However, pacing profiles of youth swimmers were evidently
434 more variable, and these swimmers demonstrated difficulty in self-selecting the most
435 beneficial pacing profile. This could indicate that youth swimmers struggle to regulate
436 their energy distribution in the most efficient manner. This inability to pace efficiently
437 was also found in a study of junior swimmers (15 ± 1.5 years) performing a swimming
438 incremental step test (Scruton et al., 2015). Youth swimmers' incompetence in
439 stabilising their pacing behaviour may be related to the finding that pacing skills are
440 contingent on prior experience and the level of (meta-) cognitive functioning, requiring
441 time to fully develop (Elferink-Gemser & Hettinga, 2017; Foster et al., 2009; Ulmer,
442 1996, Micklewright et al., 2012).

443 A recently proposed model for developing athletes' pacing skills emphasises
444 the importance of both the experiential and self-regulatory aspects of skill learning

445 (Elferink-Gemser & Hettinga, 2017). Self-regulation has proven essential for an
446 efficient training regime (Toering, Elferink-Gemser, Jordet, & Visscher, 2009). By
447 supporting the multiple cyclical facets of self-regulation learning (reflection, planning,
448 performance and evaluation), coaches can facilitate the development of young
449 athletes' pacing behaviour. The importance of this development was recently
450 highlighted in a longitudinal study of adolescent speed skaters (Wiersma et al., 2017).
451 The findings indicated that youth athletes whose pacing profiles resemble those of elite
452 performers in an earlier stage of their development went on to achieve higher
453 performance levels in their later careers, compared to their peers at youth level
454 (Wiersma et al., 2017). As swimmers' pacing behaviours resemble those of athletes in
455 time-trial sports like speed skating, it is plausible that swimmers also demonstrate a
456 similar relation between the development of their pacing behaviour and their
457 performance in later stages of their careers. Further research on the development of
458 pacing behaviour in swimming is required to address this question.

459

460 **Conclusion**

461 The present study is the first systematic investigation of the body of literature on
462 pacing behaviour in pool swimming. Although swimming is a head-to-head sport, the
463 pacing behaviour of swimmers in this type of competition is similar to that of athletes
464 in time-trial sports. A positive profile is evident in shorter races (100 m), whereas a
465 more parabolic profile is prevalent in the longer races (400 and 800 m). Additionally,
466 elite medallists demonstrate more conservative pacing behaviour, characterised by a
467 lower normalised velocity in the initial phase of the race and a higher normalised
468 velocity in the final phase. Given the unique characteristics of the breaststroke event,
469 the swimmers' pacing profile markedly deviates from those of other strokes, being

470 more positive. Blood lactate accumulates throughout the race, prompting a decrease in
471 SL and a consequent increase in SR during the course of the race to minimise variations
472 in velocity. The pacing profiles of youth swimmers are more variable than those of
473 elite swimmers and young swimmers tend to have difficulty effectively regulating their
474 energy distribution to achieve the highest performance outcome. The relationship
475 between pacing behaviour and performance development in swimmers needs to be
476 further explored in future studies.

477

478 **Declaration of interests**

479 The authors report no potential conflicts of interest that are related to the content of
480 this review.

481

482 **References**

- 483 Abbiss, C. R., & Laursen, P. B. (2008). Describing and understanding pacing strategies during
484 athletic competition. *Sports Medicine*, 38(3), 239–252.
- 485 Barbosa, T. M., Bragada, J. A., Reis, V. M., Marinho, D. A., Carvalho, C., & Silva, A. J.
486 (2010). Energetics and biomechanics as determining factors of swimming performance:
487 updating the state of the art. *Journal of Science and Medicine in Sport*, 13(2), 262–269.
- 488 Barbosa, T. M., Fernandes, R., Keskinen, K. L., Colaço, P., Cardoso, C., Silva, J., & Vilas-
489 Boas, J. P. (2006). Evaluation of the energy expenditure in competitive swimming
490 strokes. *International Journal of Sports Medicine*, 27(11), 894–899.
- 491 Brick, N. E., MacIntyre, T. E., & Campbell, M. J. (2016). Thinking and action: a cognitive
492 perspective on self-regulation during endurance performance. *Frontiers in Physiology*,
493 7.
- 494 Capelli, C., Pendergast, D. R., & Termin, B. (1998). Energetics of swimming at maximal
495 speeds in humans. *European Journal of Applied Physiology and Occupational*
496 *Physiology*, 78(5), 385–393.
- 497 De Koning, J. J., Foster, C., Lucia, A., Bobbert, M. F., Hettinga, F. J., & Porcari, J. P. (2011).
498 Using modeling to understand how athletes in different disciplines solve the same
499 problem: swimming versus running versus speed skating. *International Journal of Sports*
500 *Physiology and Performance*, 6(2), 276–280.

501 Dormehl, S. J., & Osborough, C. D. (2015). Effect of Age, Sex, and Race Distance on Front
 502 Crawl Stroke Parameters in Subelite Adolescent Swimmers During Competition.
 503 *Pediatric Exercise Science*, 27(3), 334–344.

504 Edwards, A. M., & Polman, R. C. J. (2013). Pacing and awareness: brain regulation of physical
 505 activity. *Sports Medicine*, 43(11), 1057–1064.

506 Elferink-Gemser, M. T., & Hettinga, F. J. (2017). Pacing and Self-Regulation: Important Skills
 507 for Talent Development in Endurance Sports. *International Journal of Sports Physiology*
 508 *and Performance*, 1–17.

509 Figueiredo, P., Zamparo, P., Sousa, A., Vilas-Boas, J. P., & Fernandes, R. J. (2011). An energy
 510 balance of the 200 m front crawl race. *European Journal of Applied Physiology*, 111(5),
 511 767–777.

512 FINA. (2017). World records. Retrieved from
 513 http://www.fina.org/sites/default/files/wr_50m_oct_3_2018.pdf

514 Foster, C., De Koning, J. J., Hettinga, F., Lampen, J., La Clair, K. L., Dodge, C., ... Porcari,
 515 J. P. (2003). Pattern of energy expenditure during simulated competition. *Medicine and*
 516 *Science in Sports and Exercise*, 35(5), 826–831.

517 Foster, C., Hendrickson, K. J., Peyer, K., Reiner, B., Lucia, A., Battista, R. A., ... Wright, G.
 518 (2009). Pattern of developing the performance template. *British Journal of Sports*
 519 *Medicine*, 43(10), 765–769.

520 Foster, C., Hettinga, F., Lampen, J., Dodge, C., Bobbert, M., & Porcari, J. P. (2004). Effect of
 521 competitive distance on energy expenditure during simulated competition. *International*
 522 *Journal of Sports Medicine*, 25(03), 198–204.

523 Garland, S. W. (2005). An analysis of the pacing strategy adopted by elite competitors in 2000
 524 m rowing. *British Journal of Sports Medicine*, 39(1), 39–42.

525 Giedd, J. N., Blumenthal, J., Jeffries, N. O., Castellanos, F. X., Liu, H., Zijdenbos, A., ...
 526 Rapoport, J. L. (1999). Brain development during childhood and adolescence: a
 527 longitudinal MRI study. *Nature Neuroscience*, 2(10), 861–863.

528 Hettinga, F. J., De Koning, J. J., & Foster, C. (2009). VO₂ response in supramaximal cycling
 529 time trial exercise of 750 to 4000 m. *Medicine & Science in Sports & Exercise*, 41(1),
 530 230–236.

531 Hettinga, F. J., Konings, M. J., & Pepping, G.-J. (2017). The science of racing against
 532 opponents: Affordance competition and the regulation of exercise intensity in head-to-
 533 head competition. *Frontiers in Physiology*, 8.

534 Hochstein, S., & Blickhan, R. (2014). Body movement distribution with respect to swimmer's
 535 glide position in human underwater undulatory swimming. *Human Movement Science*,
 536 38, 305–318.

537 Holmér, I. (1974). Propulsive efficiency of breaststroke and freestyle swimming. *European*

538 *Journal of Applied Physiology and Occupational Physiology*, 33(2), 95–103.

539 Konings, M. J., & Hettinga, F. J. (2017). The impact of different competitive environments on
540 pacing and performance. *International journal of sports physiology and performance*,
541 13(6):701-708

542 Konings, M. J., Noorbergen, O. S., Parry, D., & Hettinga, F. J. (2016). Pacing Behavior and
543 Tactical Positioning in 1500-m Short-Track Speed Skating. *International Journal of*
544 *Sports Physiology and Performance*, 11(1), 122–129.

545 Letts, L., Wilkins, S., Law, M., Stewart, D., Bosch, J., & Westmorland, M. (2007). Guidelines
546 for critical review form: Qualitative studies (Version 2.0). *McMaster University*
547 *Occupational Therapy Evidence-Based Practice Research Group*.

548 Lipinska, P., Allen, S. V., & Hopkins, W. G. (2016). Modeling parameters that characterize
549 pacing of elite female 800-m freestyle swimmers. *European Journal of Sport Science*,
550 16(3), 287–292.

551 Mauger, A. R., Neuloh, J., & Castle, P. C. (2012). Analysis of pacing strategy selection in elite
552 400-m freestyle swimming. *Medicine & Science in Sports & Exercise*, 44(11), 2205–
553 2212.

554 Menting, S. G. P., Konings, M. J., Elferink-Gemser, M. T., & Hettinga, F. J. (2018). Pacing
555 Behaviour of Elite Youth Athletes: Analysing 1500-m Short-Track Speed Skating.
556 *International Journal of Sports Physiology and Performance*, 1–22.

557 Micklewright, D., Angus, C., Suddaby, J., St Clair, G. A., Sandercock, G., & Chinnasamy, C.
558 (2012). Pacing strategy in schoolchildren differs with age and cognitive development.
559 *Medicine and Science in Sports and Exercise*, 44(2), 362–369.

560 Micklewright, D., Kegerreis, S., Raglin, J., & Hettinga, F. (2016). Will the Conscious–
561 Subconscious Pacing Quagmire Help Elucidate the Mechanisms of Self-Paced Exercise?
562 New Opportunities in Dual Process Theory and Process Tracing Methods. *Sports*
563 *Medicine*, 1–9.

564 Muehlbauer, T., Schindler, C., & Panzer, S. (2010). Pacing and performance in competitive
565 middle-distance speed skating. *Research Quarterly for Exercise and Sport*, 81(1), 1–6.

566 Muehlbauer, T., Schindler, C., & Widmer, A. (2010). Pacing pattern and performance during
567 the 2008 Olympic rowing regatta. *European Journal of Sport Science*, 10(5), 291–296.

568 Mytton, G. J., Archer, D. T., Turner, L., Skorski, S., Renfree, A., Thompson, K. G., ... Gibson,
569 A. S. (2015). Increased variability of lap speeds: differentiating medalists and
570 nonmedalists in middle-distance running and swimming events. *International Journal of*
571 *Sports Physiology and Performance*, 10(3), 369–373.

572 Nikolaidis, P. T., & Knechtle, B. (2017). Pacing in age-group freestyle swimmers at The XV
573 FINA World Masters Championships in Montreal 2014. *Journal of Sports Sciences*,
574 35(12), 1165–1172.

575 Noorbergen, O. S., Konings, M. J., Micklewright, D., Elferink-Gemser, M. T., & Hettinga, F.
576 J. (2016). Pacing Behavior and Tactical Positioning in 500-and 1000-m Short-Track
577 Speed Skating. *International Journal of Sports Physiology and Performance*, 11(6), 742–
578 748.

579 Swimming World Magazine (2017). Sport Publications, Inc. Retrieved from
580 <http://www.swimmingworldmagazine.com/results/open-water>

581 Robertson, E., Pyne, D., Hopkins, W., & Anson, J. (2009). Analysis of lap times in
582 international swimming competitions. *Journal of Sports Sciences*, 27(4), 387–395.

583 Rubia, K., Russell, T., Overmeyer, S., Brammer, M. J., Bullmore, E. T., Sharma, T., ...
584 Andrew, C. M. (2001). Mapping motor inhibition: conjunctive brain activations across
585 different versions of go/no-go and stop tasks. *Neuroimage*, 13(2), 250–261.

586 Saavedra, J. M., Escalante, Y., Garcia-Hermoso, A., Arellano, R., & Navarro, F. (2012). A 12-
587 year analysis of pacing strategies in 200- and 400-m individual medley in international
588 swimming competitions. *Journal of Strength and Conditioning Research*, 26(12), 3289–
589 3296.

590 Schnitzler, C., Seifert, L., & Chollet, D. (2009). Variability of coordination parameters at 400-
591 m front crawl swimming pace. *Journal of Sports Science and Medicine*, 8(2), 203–210.

592 Scruton, A., Baker, J., Roberts, J., Basevitch, I., Merzbach, V., & Gordon, D. (2015). Pacing
593 accuracy during an incremental step test in adolescent swimmers. *Open Access Journal*
594 *of Sports Medicine*, 6, 249–257.

595 Simbaña Escobar, D., Hellard, P., Pyne, D. B., & Seifert, L. (2018). Functional role of
596 movement and performance variability: Adaptation of front crawl swimmers to
597 competitive swimming constraints. *Journal of Applied Biomechanics*, 34(1), 53–64.

598 Skorski, S., Faude, O., Abbiss, C. R., Caviezel, S., Wengert, N., & Meyer, T. (2014). Influence
599 of Pacing Manipulation on Performance of Juniors in Simulated 400-m Swim
600 Competition. *International Journal of Sports Physiology and Performance*, 9(5), 817–
601 824.

602 Skorski, S., Faude, O., Caviezel, S., & Meyer, T. (2014). Reproducibility of Pacing Profiles in
603 Elite Swimmers. *International Journal of Sports Physiology and Performance*, 9(2),
604 217–225.

605 Skorski, S., Faude, O., Rausch, K., & Meyer, T. (2013). Reproducibility of Pacing Profiles in
606 Competitive Swimmers. *International Journal of Sports Medicine*, 34(2), 152–157.

607 Smits, B. L. M., Pepping, G.-J., & Hettinga, F. J. (2014). Pacing and decision making in sport
608 and exercise: the roles of perception and action in the regulation of exercise intensity.
609 *Sports Medicine*, 44(6), 763–775.

610 Stoter, I. K., MacIntosh, B. R., Fletcher, J. R., Pootz, S., Zijdewind, I., & Hettinga, F. J. (2016).
611 Pacing Strategy, Muscle Fatigue, and Technique in 1500-m Speed-Skating and Cycling

612 Time Trials. *International Journal of Sports Physiology and Performance*, 11(3), 337–
613 343.

614 Taylor, J. B., Santi, G., & Mellalieu, S. D. (2016). Freestyle race pacing strategies (400m) of
615 elite able-bodied swimmers and swimmers with disability at major international
616 championships. *Journal of Sports Sciences*, 34(20), 1913–1920.

617 Thompson, K. G., MacLaren, D. P., Lees, A., & Atkinson, G. (2003). The effect of even,
618 positive and negative pacing on metabolic, kinematic and temporal variables during
619 breaststroke swimming. *European Journal of Applied Physiology*, 88(4–5), 438–443.

620 Thompson, K. G., MacLaren, D. P. M., Lees, A., & Atkinson, G. (2004). The effects of
621 changing pace on metabolism and stroke characteristics during high-speed breaststroke
622 swimming. *Journal of Sports Sciences*, 22(2), 149–157.

623 Toering, T. T., Elferink-Gemser, M. T., Jordet, G., & Visscher, C. (2009). Self-regulation and
624 performance level of elite and non-elite youth soccer players. *Journal of Sports Sciences*,
625 27(14), 1509–1517.

626 Toussaint, H. M. (1990). Differences in propelling efficiency between competitive and
627 triathlon swimmers. *Medicine and Science in Sports and Exercise*, 22(3), 409–415.

628 Toussaint, H. M., Beelen, A., Rodenburg, A., Sargeant, A. J., de Groot, G., Hollander, A. P.,
629 & van Ingen Schenau, G. J. (1988). Propelling efficiency of front-crawl swimming.
630 *Journal of Applied Physiology*, 65(6), 2506–2512.

631 Ulmer, H.-V. (1996). Concept of an extracellular regulation of muscular metabolic rate during
632 heavy exercise in humans by psychophysiological feedback. *Cellular and Molecular Life
633 Sciences*, 52(5), 416–420.

634 van Ingen Schenau, G. J., & Cavanagh, P. R. (1990). Power equations in endurance sports.
635 *Journal of Biomechanics*, 23(9), 865–881.

636 van Ingen Schenau, G. J., De Koning, J. J., & De Groot, G. (1992). The distribution of
637 anaerobic energy in 1000 and 4000 metre cycling bouts. *International Journal of Sports
638 Medicine*, 13(06), 447–451.

639 Vantorre, J., Chollet, D., & Seifert, L. (2014). Biomechanical analysis of the swim-start: a
640 review. *Journal of Sports Science and Medicine*, 13(2), 223–231.

641 Veiga, S., & Roig, A. (2016). Underwater and surface strategies of 200 m world level
642 swimmers. *Journal of Sports Sciences*, 34(8), 766–771.

643 Wiersma, R., Stoter, I. K., Visscher, C., Hettinga, F. J., & Elferink-Gemser, M. T. (2017).
644 Development of 1500m Pacing Behavior in Junior Speed Skaters: A Longitudinal Study.
645 *International Journal of Sports Physiology and Performance*, 1–20.


646 Zamparo, P., Bonifazi, M., Faina, M., Milan, A., Sardella, F., Schena, F., & Capelli, C. (2005).
647 Energy cost of swimming of elite long-distance swimmers. *European Journal of Applied
648 Physiology*, 94(5–6), 697–704.

649

650 Insertions and captions

651 Figure 1. Flow diagram of the literature selection process, including the number of
652 articles excluded at each stage.

653



654

655 *Figure 1. Flow diagram of the literature selection process, including the number of articles excluded at each stage.*

656

657

658 Table 1. A quality assessment of the included articles in alphabetical order applying
659 the guidelines developed by Letts et al. (2007).

660 Table 1. A quality assessment of the included articles in alphabetical order applying
661 the guidelines developed by Letts et al. (2007).

Assessment questions →	1	2	3	4	5	6	7	8	9	10	11	12	13	14	Total
1. Dormehl and Osborough (2015)	1	1	1	0	0	1	0	0	1	1	1	1	0	0	8
2. Figueiredo, Zamparo, Sousa, Vilas-Boas, & Fernandes (2011)	1	1	1	0	0	1	0	0	1	1	1	1	0	0	8

3.	Lipinska, Allen, and Hopkins (2016)	1	1	1	0	0	1 ^a	0	0	1	1	0	1	1	0	8
4.	Mauger, Neuloh, & Castle (2012)	1	1	1	0	0	1 ^b	0	0	1	1	1	1	1	1	10
5.	Mytton et al. (2015)	1	1	1	0	1	1 ^a	0	0	1	1	1	1	0	1	10
6.	Nikolaidis and Knechtle (2017)	1	1	1	0	0	1 ^a	0	0	1	1	1	1	1	1	10
7.	Robertson, Pyne, Hopkins, & Anson (2009)	1	1	1	0	0	1 ^a	0	0	0	1	1	1	1	1	9
8.	Saavedra, Escalante, Garcia-Hermoso, Arellano, & Navarro (2012)	1	1	1	0	0	1 ^a	1	1	1	1	1	1	0	0	10
9.	Schnitzler, Seifert, & Chollet (2009)	1	1	1	0	0	1	0	0	1	1	1	1	1	0	9
10.	Skorski, Faude, Abbiss, et al. (2014)	1	1	1	0	0	1	1	1	1	1	1	1	1	1	12
11.	Skorski, Faude, Caviezel, & Meyer (2014)	1	1	1	0	0	1 ^a	1	1	1	1	1	1	1	1	12
12.	Skorski, Faude, Rausch, & Meyer et al. (2013)	1	1	1	0	0	1	0	0	1	1	1	1	1	1	10
13.	Taylor, Santi, & Mellalieu (2016)	1	1	1	0	0	1 ^b	0	0	1	1	1	1	0	1	9
14.	Thompson, MacLaren, Lees, & Atkinson (2003)	1	1	1	0	0	1	0	0	1	1	1	1	0	0	8
15.	Thompson, MacLaren, Lees, & Atkinson (2004)	1	1	1	0	0	1	0	0	1	1	1	1	0	0	8
16.	Veiga and Roig (2016)	1	1	1	0	0	1 ^b	0	0	1	1	1	1	0	1	9

662

a = records were in the public domain. b = no informed consent but there was ethical approval.

663

Included questions (scored either 0 or 1): 1.Was the aim of the study and purpose stated clearly? 2.Was relevant background literature reviewed? 3.Was the study design appropriate for the research question? 4.Were the participants relevant to the research question and was their selection well-reasoned? 5.Was the sample size justified? 6.Was informed consent obtained? 7.Were the outcome measures reliable? 8.Were the outcome measures valid? 9.Were results reported in terms of statistical significance? 10.Were the data collection methods appropriate for the research design? 11.Did a meaningful picture of the phenomenon under study emerge? 12.Were conclusions appropriate given the study findings? 13.Are there any implications for future research given the results of the study? 14.Were limitations of the study acknowledged and described by the authors?

670

671

672 Table 2. An overview of the reviewed studies on pacing behaviour in pool swimming
673 ordered by race distance (n =16).

674

675 Table 2. An overview of the reviewed studies on pacing behaviour in pool swimming
676 ordered by race distance (n =16).

Study	Race distance	Gender and number of participants	Age (years)	Performance level	Stroke type	Competition type (stage of competition)	Methods	Statistical analyses	Pacing profile	Main results
-------	---------------	-----------------------------------	-------------	-------------------	-------------	---	---------	----------------------	----------------	--------------

Dormehl & Osborough (2015)	100m ¹ , 200m ¹	Male (n=56), Female (n=56)	Group 1: 14.4±0.7 Group2: 17.0±0.9	Competitive	Freestyle	Real competition (heats, semi- finals and finals)	Races collected at international schools swimming championships ¹ . Race split up in quarters. For 200m: laps 1, 3 and 5 for quarter 1, 2 and 3. Quarter 4 is combination of laps 7 and 8 Measurements: -Race time -Velocity per quarter. (Velocity of first quarter was measured between 15m and 20m to account for dive. Remainders over a 10m midsection of the pool)	-Repeated measurements ANOVA's -Post-Hoc (Bonferroni)	-Positive	-No difference in pacing profile between groups 100m: -Velocity decreased for each quarter. 200m: -Velocity decreased for each quarter except for the last quarter in which it did not differ from the third quarter.
Robertson, Pyne, Hopkins, & Anson (2009)	100m ² , 200m ² , 400m ²	Male (n=1530), Female (n=1527)	n/a	Elite	100m, 200m: Freestyle, breaststroke, butterfly, backstroke. 400m: Freestyle, medley	Real competition (semi-finals and finals)	Races collected during OG, WC, EC and CG over a 7 year period. Measurements: -Total race time -Split times (50m or 100m) -Placing for top 16 finishers	Lap times of finalists and semi-finalists, the mean lap times for all the swimmers (placed 1–16) were plotted between- athletes and within-athletes.	100m: Positive 200m: Fast- start- even. 400m: Parabolic	-Winners maintained a lead through each of the intermediate laps. -Pacing profile: faster first lap (~1-3s), followed by evenly paced middle laps and an evenly paced or slightly faster (~1s) final lap. -The most successful(top 3 of 16) swimmers were faster in all the laps.
Nikolaidis & Knechtle (2017)	100m ² , 200m ² , 400m ² , 800m ²	Males (n=2260), Females (n=2221)	25-94	Competitive	Freestyle	Real competition (heats, semi- finals and finals)	Races were collected during the Masters championships 2014. Measurements -50m split times (200m, 400m) -100m split times (800m) -Total race time	-Mixed-design factorial ANOVA -Post-Hoc (Bonferroni) test. Effect size eta squared (η^2): small ($0.010 < \eta^2 \leq 0.059$), moderate ($0.059 < \eta^2 \leq 0.138$) and large ($\eta^2 > 0.138$).	- Parabolic -Positive	100m: Velocity 1 st lap > 2 th lap (+11.6%) 200m: Velocity: 1 st lap > 2 th lap (+11.6%) > 3 th lap (+3.8%) < 4 th lap (-2.1%) ($P < 0.001$, $\eta^2 = 0.847$), - Larger changes in older age groups than in the younger groups, both in women ($P < 0.001$, $\eta^2 = 0.195$)and men ($P < 0.001$, $\eta^2 = 0.200$). 400m: -Swimming time: 50–100 m (+11.1%), 101–150 m (+2.9%), 151–200 m (+1.2%), 201–250m (unchanged), 251– 300 m (+0.5%), 301– 350 m (-0.6%), 351– 400 m (-4.5%) ($P < 0.001$, $\eta^2 = 0.856$). -Larger changes in older age groups than in the younger groups, both in women ($P < 0.001$, $\eta^2 = 0.176$) and men ($P < 0.001$, $\eta^2 = 0.131$). 800m: -Swimming time: 100–200 m

										(+8.8%), 201–300 m (+1.0%), 301–400 m (+0.5%), 401–500 m (+0.2%), 501–600 m (+0.2%), 601–700 m (–0.3%) 701–800 m (–3.4%) ($P < 0.001$, $\eta^2 = 0.842$). - Larger changes in older age groups than in the younger groups in men ($P < 0.001$, $\eta^2 = 0.105$).
Thompson, MacLaren, Lees & Atkinson (2003)	200m. 175m	Male (n=9)	21.2±2.6	Sub-Elite	Breaststroke	Simulated competition (without opponent)	200m test trial 3 paced 175m trials. Measurements: -50m split times -HR -RPE -La (post-trial)	-Dependent t-tests -one-way ANOVA -Factorial ANOVA -Post hoc (Tukey's HSD)	-Even -Positive - Negative	-Difference in split times ($p < 0.01$) -No difference in finishing times ($p < 0.05$). -The 200 test trial: positively paced (split time: 72.2±8.6s; finish time: 148.0±13.2s) -RPE: even < positive, 200m trial ($p < 0.05$). -HR: negative trial < others ($p < 0.05$)
Thompson, MacLaren, Lees, & Atkinson (2004)	200m	Male (n=9)	22.5±4.5	Competitive	Breaststroke	Simulated competition (without opponent)	200m test trial 3 paced 200m trials: -98% of 200m time -100% of 200m time -102% of 200m time Measurements: -50m split times -HR -Respiratory exchange ratio (RER) (post-trial) -La (post-trial) 50m, 100m, 150m and 200m trial at 200m velocity. No dive, no underwater phase. Measurements: -50m split times. -Velocity during every 50m. -VO ₂ -La (post-trial) -Aerobic, anaerobic (lactate and alactic) -Total energy expenditure	-Dependent t-tests a -One-way ANOVA -Factorial ANOVA -Post-hoc (Tukey's HSD)	-Positive	-Finishing times: different ($F=28.37$, $p < 0.01$). 102% > 100% (0.8%, $p < 0.05$). - 102% was positively paced ($t=4.88$, $p < 0.006$). -RER and blood lactate 102% > 100% & 98% ($p < 0.05$) -PRE: 102% > 98% ($p < 0.05$) -HR: at 100m 102% > 98% ($F=4.00$, $p < 0.03$). No difference at 200m.
Figueiredo, Zamparo, Sousa, Vilas-Boas, & Fernandes (2011)	200m ¹	Male (n=10)	21.6±2.4	Elite	Freestyle	Simulated competition (without opponent)	50m, 100m, 150m and 200m trial at 200m velocity. No dive, no underwater phase. Measurements: -50m split times. -Velocity during every 50m. -VO ₂ -La (post-trial) -Aerobic, anaerobic (lactate and alactic) -Total energy expenditure	-One-way repeated measures ANOVA -Post-Hoc (Bonferroni) -Cohen's f; small ($0 \leq f \leq 0.10$), medium ($0.10 < f \leq 0.25$); and large effect size ($f > 0.25$).	Fast-start-even.	-Velocity: first lap > other laps ($F_{3,27}=24.72$, $p < 0.01$, $f=1.04$) -Split time: first lap < other laps ($F_{3,27}=30.753$, $p < 0.001$, $f=1.23$) -Aerobic contribution: stable the last 3 laps, lower in 1 st lap ($F_{3,27}=110.515$, $p < 0.001$, $f=5.69$). -Anaerobic anlactic contribution: 1 st lap > other laps ($F_{3,27}=925.91$, < 0.01 , $f=5.69$) -Anaerobic lactate contribution: 1 st lap > other laps ($F_{3,27}=66.131$, $p < 0.001$, $f=1.73$) -Total energy expenditure: 1 st & 4 th lap > other laps ($F_{3,27}=19.578$, $p < 0.001$, $f=0.59$) -Total energy expenditure: 2 th lap > 3 th lap ($F_{3,27}=29.137$, $p < 0.001$, $f=0.80$).

Veiga & Roig (2016)	200m ²	Males (n=64), Females (n=64)	n/a.	Elite	Butterfly, backstroke, breaststroke, freestyle.	Real competition (semi-finals and finals)	Races collected during the FINA WC 2013. Measurements: -average underwater velocity -average free swimming velocity -average lap velocity	-Repeated-measurement ANOVA - Univariate analyses using Wilks' methods.	-Positive pacing	-Free swimming velocity: 1 st lap > 2 th lap (-0.08 m·s ⁻¹ , -0.07 to -0.09 m·s ⁻¹ , P = 0.001) 2 th lap > 3 th lap > 4 th lap (both -0.02 m·s ⁻¹ , -0.01 to -0.03 m·s ⁻¹ , P = 0.001). -Underwater velocity : 1 st turn > 2 th turn (-0.03 m·s ⁻¹ , -0.01 to -0. m·s ⁻¹ , P = 0.005), 2 th = 3 th turn (0.01 m·s ⁻¹ , -0.01 to 0.03 m·s ⁻¹ , P = 0.55). -Average velocity: 1 st lap > 2 th lap (-0.15 m·s ⁻¹ , -0.15 to -0.16 m·s ⁻¹ , P = 0.001) 2 th lap > 3 th lap (-0.03 m·s ⁻¹ , -0.02 to -0.03 m·s ⁻¹ , P = 0.001). 3 th lap = 4 th lap (-0.01 m·s ⁻¹ , -0.02 to 0.00 m·s ⁻¹ , P = 0.29).
Saavedra, Escalante, Garcia-Hermoso, Arellano, & Navarro (2012)	200m ² , 400m ²	Male (n=821), Female (n=822)	n/a	Elite	Medley	Real competition (semi-finals and finals)	Races were collected during OG. WC, EC, CG, PPC, U.S. Olympic team trials, Australian Olympic team trial in 2000-2011. Measurements: -Total race time -50m split times -percentage of total time spend in a lap.	-A two-way ANOVA sex*classification -Post-Hoc (Bonferroni)	- Parabolic	200m: -The percentage of time spend per stroke: Butterfly men (22.59±0.42), women (22.65±0.42) < freestyle men (23.20±0.42), women (22.90±0.46) < backstroke men (25.62±0.53), women (25.52±0.52) < breaststroke men (28.59±0.60), women (28.93±0.65) -The best swimmers: greater percentage in butterfly and freestyle (p<0.001) and less in backstroke (p<0.001) and breaststroke (p<0.021) compared to the lowest classified swimmers. 400m: -The best swimmers: greater percentage in butterfly and backstroke (p<0.001) and less in backstroke (p<0.018) and breaststroke (p<0.024) compared to the lowest classified swimmers.
Skorski, Faude, Caviezel, & Meyer (2014)	200m ² , 400m ²	Male (n=158)	22.8±2.9	Elite	200m: freestyle, butterfly, backstroke, breaststroke. 400m: freestyle	Real competition (heats, semi-finals and finals)	Races of top 50 swimmers collected during 22 national and international events as well as the races of the finals (1 th -16 th place) of the PPC and EC. Measurements:	-Repeated measures ANOVA (factor 1, competition; factor 2, section of the race) -Post-Hoc (Scheffé).	-Fast-start-even -Positive - Parabolic	-Average performance improvement from heat to final was 1.2% (CL 0.6-2.2%). -Pacing pattern: Fast-start-even pattern in 200m freestyle, butterfly and backstroke. Velocity in 1 th lap > others. (P<.001) and

							<ul style="list-style-type: none"> -Overall race times -50m split times -Normalized velocity 			<p>2th lap > 3th and 4th in freestyle and backstroke (P<.001). Positive profile in breaststroke (P<.001). Parabolic profile in 400m freestyle. Velocity in 1st lap > others (P<.001). Last lap > others (P<.001). -Heat paced similar to finals (interaction: all P>.06). 50m split times were faster in finals (P<.02). -Normalized pacing pattern was not significantly different between competitions 1 and 2 (P>.18). -CV's for intra-individual differences in split times between heats and finals were small for all 200m races (<2.2%; CL 0.6-3.2%). In 400m freestyle, values increased in the course of the race up to 2.9% (CL 2.2-4.5%) in the last section.</p>
Skorski, Faude, Rauch, & Meyer. (2013)	200m ² , 400m ² , 800m ²	Male (n=9), Female (n=7)	16.9±2.1	Sub-Elite	Freestyle	<p>Simulated competition (with opponents) & Real competition (heats, semi-finals and finals)</p>	<ul style="list-style-type: none"> -Six simulated competitions (SC: 2x 200m, 2x 400m and 2x 800m. -Real competition races (RC) Measurements: -50m splits times (200m) and 100m splits (400m, 800m). -Peak blood lactate values (post-trial) -HR (post trial) 	<ul style="list-style-type: none"> -2-way repeated measures ANOVA test*section of test -Cohen's d -Within-subject-variation by means of the SEM and log-transformed CV. 	<ul style="list-style-type: none"> -Fast-start-even 	<p>-Fast-start profile during SC (p<0.002) and RC (p<0.001). -CV for test-retest small for first 3 sections (CV < 2.0%, for first 6 sections of 800m) and increased towards the end. -Pacing pattern SC = RC (p>0.22). -Pacing pattern for absolute velocities SC = RC (p>0.10), all section times faster during RC (p<0.001). -SEM in split times between SC and RC were small in the middle of the race during 800m (200m-600m) and 400m (200m-300m) (SEM <1.6s). The first section higher SEM in both distances (>1.8s). The last section of the during the 400m (300m-400m) and the 2 last sections during the 800m (600m-800m) showed higher SEM (>1.8s).</p>
Mauger, Neuloh, & Castle. (2012)	400m ²	Male (n=147), Female (n=117)	n/a	Elite	Freestyle	Real competition (finals)	<p>Races collected at the EC, WC, British and Australian national championship and International Invitational Meets, CG and OG in period 2003-2010.</p>	<p>Data were analyzed using a three-way ANOVA (pacing strategy _ sex _ swimming suit) in an unrelated design (p=0.05).</p>	<ul style="list-style-type: none"> - Parabolic -Fast-start-even -Positive - Negative -Even 	<p>-Fast-start-even and parabolic pacing profiles used the most, with parabolic profiles preferred by men. -Fast-start-even pacing profile performed at 96.08±2.12% of the (228.4±4.66s).</p>

							Measurements: -Total race time -50m split times -Mean velocity of every lap (excluding first 10m after the start and first and last 5m of every lane). Pacing profiles were determined by an algorithm based on normalized velocity.			-Parabolic pacing profile performed at 96.04±2.2% of the WR (228.7±4.84s) -Positive pacing profile performed at 95.4±2.19% of the (230.15±4.82s) -1.7s performance difference between fast-start-even and positive pacing ($F_{2,228} = 1.00, P > 0.05$).
Taylor, Santi, & Mellalieu (2016)	400m ²	Male (n=489), Female (n=312)	n/a	Elite	Freestyle	Real competition (heats, semi-finals and finals)	Races collected at the WC, EC, OG between 2006 and 2012. Measurements: -50m split times -Normalized 50m split time	-k-means cluster analysis -One-way ANOVA -Cohen's d.	-Fast-start -Positive - Parabolic	-In males mean race time in parabolic pacing (mean race time = 230.57 s, 95% CL = 229.51–231.63) < fast-start-even pacing (mean race time = 235.91 s, 95% CL = 234.81–237.01), and positive (mean race time = 252.66 s, 95% CL = 249.26–256.06) -In females mean race time in parabolic pacing profile (mean race time = 249.59 s, 95% CL = 248.47–250.71) < fast-start-even (mean race time = 253.94 s, 95% CL = 252.87–255.01), positive (mean race time = 262.76 s, 95% CL = 260.05–265.47) -Fast-start-even and parabolic pacing profiles were most frequently observed. 220 and 182 for males (n=498) and 105 and 135 for females (n=312) respectively.
Mytton et al. (2015)	400m ²	Male (n=48)	n/a	Elite	Freestyle	Real competition (finals)	Races collected at the EC 2006, 2010, 2012. WC 2007, 2010 and CG 2006. Measurements: -50m split times -Velocity per lap -Normalized velocity	-Mann Whitney test -Kruskal-Wallis test -Cohens <i>d</i> effect size: trivial (<0.2), small (0.2-0.6), moderate (0.6-1.2) and large (1.2-2.0).	-Fast-start-even - Parabolic	-Medallists: larger variation in velocity compared to non-medallists -Lap one: normalized velocity medallists < non-medallists (102.2±1.2%, 103.1±1.1%, $p = 0.03, d = 0.75$). Gold medallists = others -Lap two: normalized velocity medallists < non-medallists (97.7±0.8%, 98.2±0.6%, $p < 0.001, d = 0.78$). -Lap three: Normalized velocity medallists = non-medallists (98.5±1.0%, 98.4±0.6%, $p = 0.63$). Gold medallist > 4 th -8 th place ($p = 0.04$ to 0.002). -Lap four: Normalized velocity medallists > medallists

Skorski et al. (2014)	400m ²	Male (n=10), female (n=5)	Male: 19.2±2.0 Female: 16.2±1.8	Sub-Elite	Freestyle	Simulated competition (without opponent)	Self-paced trial (PP _{SS}), trial with first 100m paced 3% slower compared to PP _{SS} (PP _{SLOW}), trial with first 100m paced 3% faster compared to PP _{SS} (PP _{FAST}). Controlled for the dive start. Measurements: -Total racing time -50m split times -La (post-trial) -Normalized velocity	-One-way repeated-measures ANOVA -Two-way ANOVA SR and normalized velocity between trials (with and without start dive). -Post-Hoc (Scheffé) -Cohen d effect (0.2, 0.6, 1.2, 2.0, and 4.0 for trivial, small, moderate, large, very large, and extremely large, respectively)	- Parabolic	(101.8±1.7%, 100.5±1.2%, p≤0.01, d = 0.93 moderate). -Overall performance compared to PP _{SS} : < PP _{FAST} (278.5±16.4s)(P=.05). < PP _{SLOW} (277.5±16.1s)(P=.20). - 7 out of 15 subjects faster time in a manipulated race (3 in PP _{FAST} , 4 in PP _{SLOW}) -Pacing was different between conditions in the first 100m (P<0.001), not in the rest of the trial (P=0.45). -Including the dive start: in all conditions the first 50m were faster compared to the remaining sections of the trial (P<0.001). -Blood lactate (P=.33) and HR (P=.47) were not different between conditions.
Schnitzler, Seifert, & Chollet (2009)	400m ¹	Male (n=6), Female (n=6).	18.2±2.2	Sub-Elite	Freestyle	Simulated competition (without opponent)	100m, 200m, 300m and 400m at 400m velocity. No dive start. Measurements: -HR -La (post-trial) -Mean speed every 50m (V50) -Workload (TWL). By the NASA-TLX questionnaire.	-Three-way ANOVA (fixed factors: swim, gender; random factor: subject) -Three-way ANOVA (fixed factors: swim distance, gender; random factor: subject) -Post-Hoc (Tukey HSD) -CV -One-way ANOVA	-Even	-HR, lactate values and TWL increased with distance for both genders (p<0.05). -HR: increased from 100m to 400m (p<0.05). 200m = 300m. -Lactate: increased from 100m to 400m (p<0.05). 200m = 300m. -Velocity: first 50m > other laps (p<0.05)
Lipinska, Allen, & Hopkins (2016)	800m ²	Female (n=192)	17-34	Elite	Freestyle	Real competition (heats, semi-finals and finals)	Races collected during OG, WC, EC, PPC, Universiades, NC. Measurements: -50m split times -Pacing profiles: linear and quadratic coefficient for the effect of lap number, reductions in time for the first and last laps, and the residual standard error of the estimate.	-Reliability analyses	-Positive	-Mean values of the linear and quadratic coefficients represent a swim with a shallow negative curvature and a slowest lap time in the eleventh lap. -First and last laps were much faster than predicted by the quadratic curve (extremely large and very large reductions in time, respectively).

677

678 1. Short course (25m pool)

679 2. Long course (50m pool)

680

681 Abbreviation list:

682 Competitions: World Championship (WC), European Championship (EC), Olympic Games (OG), Pan-pacific championship (PPC), Commonwealth Games (CG), World record (WR).

684 Measurements: Heart rate (HR), Stroke rate (SR), Rate of Perceived exertion (RPE), Oxygen uptake (VO₂), Blood lactate peak value (La).

686 Statistical analyses: analyses of variance (ANOVA), coefficient of variation (CV), confidence limits (CL), confidence intervals

687 (CI), the standard error of measurement (SEM).

688