

# Northumbria Research Link

Citation: Kordi, Mehdi, Folland, Jonathan, Goodall, Stuart, Haralabidis, Nicos, Maden-Wilkinson, Tom, Patel, Tejal Sarika, Leeder, Jonathan, Barratt, Paul and Howatson, Glyn (2020) Mechanical and Morphological Determinants of Peak Power Output in Elite Cyclists. *Scandinavian Journal of Medicine and Science in Sports*, 30 (2). pp. 227-337. ISSN 0905-7188

Published by: Wiley-Blackwell

URL: <https://doi.org/10.1111/sms.13570> <<https://doi.org/10.1111/sms.13570>>

This version was downloaded from Northumbria Research Link:  
<http://nrl.northumbria.ac.uk/id/eprint/41030/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

**Title: Mechanical and Morphological Determinants of Peak Power Output in Elite Cyclists.**

**Running Title:** Peak Power Output in Sprint Cycling

**Authors:** Mehdi Kordi<sup>1,2,6</sup>, Jonathan Folland<sup>3</sup>, Stuart Goodall<sup>1</sup>, Nicos Haralabidis<sup>4</sup>, Tom Maden-Wilkinson<sup>5</sup>, Tejal Sarika Patel<sup>4</sup>, Jonathan Leeder<sup>6</sup>, Paul Barratt<sup>2</sup>, Glyn Howatson<sup>1,7</sup>

<sup>1</sup> Department of Sport, Exercise and Rehabilitation, Northumbria University, Newcastle, UK

<sup>2</sup> British Cycling, National Cycling Centre, Manchester, UK

<sup>3</sup> School of Sport, Exercise & Health Sciences, Loughborough University

<sup>4</sup> University of Bath, UK

<sup>5</sup> Sheffield Hallam University

<sup>6</sup> English Institute of Sport, UK

<sup>7</sup> Water Research Group, North West University, Potchefstroom, South Africa

**Corresponding Author:**

Mehdi Kordi

[mehdikordi@hotmail.co.uk](mailto:mehdikordi@hotmail.co.uk)

## **Acknowledgements**

This study would not have been possible without the coaches who helped facilitate the study and the riders for taking part. In addition, the authors would also like to thank Karen Voigt and Rebecca Postlethwaite for their help in carrying out the MRI scans and analysis. This work was supported by a collaborative doctoral research project between the English Institute of Sport and Northumbria University.

## ABSTRACT

Mechanical peak power output (PPO) is a determinant of performance in sprint cycling. The purpose of this study was to examine the relationship between PPO and putative physiological determinants of PPO in elite cyclists, and to compare sprint performance between elite sprint and endurance cyclists. Thirty-five elite cyclists (18 endurance; 17 sprint) performed duplicate sprint cycling lab tests to establish PPO and its mechanical components. Quadriceps femoris ( $Q_{VOL}$ ) and hamstrings muscle volume ( $HAM_{VOL}$ ) were assessed with MRI, vastus lateralis pennation angle ( $P\theta_{VL}$ ) and fascicle length ( $FL_{VL}$ ) were determined with ultrasound imaging, and neuromuscular activation of three muscles were assessed using EMG at PPO during sprint cycling. For the whole cohort there was a wide variability in PPO (range 775-2025 W) with very large, positive, bivariate relationships between PPO and  $Q_{VOL}$  ( $r = 0.87$ ),  $HAM_{VOL}$  ( $r = 0.71$ ) and  $P\theta_{VL}$  ( $r = 0.81$ ). Step-wise multiple regression analysis revealed that 87% of the variability in PPO between cyclists was explained by two variables  $Q_{VOL}$  (76%) and  $P\theta_{VL}$  (11%). The sprint cyclists had greater PPO (+61%;  $P < 0.001$  vs endurance), larger  $Q_{VOL}$  ( $P < 0.001$ ) and  $BF_{VOL}$  ( $P < 0.001$ ) as well as more pennate vastus lateralis muscles ( $P < 0.001$ ). These findings emphasise the importance of quadriceps muscle morphology for sprint cycling events.

**Key Words:** Maximum cadence; Maximum power; Maximum torque; Muscle

## 1 INTRODUCTION

Peak power output can be described as the maximum power one can achieve during maximal tasks of a short duration (i.e. < 7 s).<sup>1</sup> Peak power output is a strong predictor of sporting performance during maximal athletic tasks such as sprint running,<sup>2</sup> jumping<sup>3</sup> and cycling.<sup>4</sup> In cycling, mechanical peak power output (PPO) relative to body mass or frontal area has been highly correlated with both acceleration<sup>5,6</sup> and maximum velocity<sup>4</sup>, both of which are related to track sprint cycling performance. Despite these relationships being well established, the underlying physiological determinants of PPO in cycling are poorly researched and understood. Whilst lean leg volume,<sup>4</sup> muscle architecture<sup>7</sup> and neuromuscular activation<sup>8</sup> have been suggested to influence cycling PPO, their quantitative contribution has not been examined. A better understanding of these factors will facilitate exercise prescription targeted more effectively to the key determinants of PPO and may help to maximise performance.

Energy systems and mechanical power output profiles of athletes in sprint cycling disciplines such as BMX and track sprint have been previously documented to have similarly high PPO.<sup>4,9,10</sup> However, physiological comparisons between sprinters (that compete in events that are maximal, 'all-out' and usually last between 9 - 60 s) and endurance riders (events that last from ~4 min to in excess of 4 hours) have not examined the physiological factors that may underpin any differences in PPO.

Theoretically muscle size, specifically muscle volume, is a key predictor of neuromuscular power<sup>11</sup> and there is evidence, that, for example, quadriceps femoris volume explains a high proportion of the variance in single joint knee extension (~80%) and squat jump (90%) power.<sup>12</sup> It has also been suggested that muscle volume is a major predictor of PPO in sprint cycling.<sup>6,13</sup> However, previous work has examined relatively crude estimates of muscle mass/volume (e.g. based on tape measures of superficial anthropometry that are not muscle group specific) in relation to sprint cycling performance.<sup>4,14</sup> Of the lower limb muscle groups,

strength of the knee extensors appears to be the best predictor of cycling PPO, <sup>15,16</sup> and thus, accurate assessment of quadriceps femoris muscle volume e.g. with magnetic resonance imaging (MRI), the gold standard technique, <sup>17</sup> might be expected to be a key determinant of PPO.

Another important component of skeletal muscle mechanics and function is muscle architecture, including pennation angle and fascicle length, that can be assessed *in vivo* with ultrasound imaging. A greater pennation angle is thought to be associated with an improvement of the generation of force output for contractions against high loads by packing more sarcomeres in parallel. <sup>18,19</sup> Whilst fascicle length plays an important role in determining shortening velocity of a muscle. <sup>20</sup>

Furthermore, the ability to develop contractile force, and thus power, rapidly is dependent on neuromuscular activation <sup>21,22</sup> and likely plays a role in cycling PPO. <sup>13</sup> However, the relationship between PPO and neuromuscular activation in a large cohort of elite cyclists has not been investigated.

The primary aim of this study was to examine the relationship of a range of putative neuromuscular determinants (muscle volume, architecture and neuromuscular activation) with cycling PPO. This involved a large cohort of elite cyclists that were all familiar with performing maximum cycling efforts and drawn from different disciplines in order to ensure a wide range of sprint cycling values. The secondary aim was to compare and characterise the sprint performance and physiological measures of elite sprint and endurance cyclists. We hypothesised that muscle volume of the quadriceps femoris would be the primary predictor of sprint cycling PPO.

## 2 METHODS

### *Participants*

Thirty-five elite male cyclists volunteered to take part in the study (mean  $\pm$  SD age, 22  $\pm$  4 yr; stature, 179.1  $\pm$  5.9 cm; mass, 77.4  $\pm$  11.3 kg). The whole cohort included two groups: sprint (n = 17; age, 21  $\pm$  3 yr; stature, 178  $\pm$  4.0 cm; mass, 85.3  $\pm$  9.2 kg) and endurance cyclists (n = 18; age, 22  $\pm$  4 yr; stature, 179.1  $\pm$  5.9 cm; mass, 69.1  $\pm$  5.9 kg). The sprint included disciplines that are 'all-out'/ maximal for  $\leq 60$  s, i.e. BMX (n = 4) and track sprinters (n = 13). Endurance included disciplines that were  $> 4$  mins in duration and are not 'all-out', i.e. track endurance who rode team pursuit (n = 9), road endurance and/or road time trial (n = 7) and mountain bike (n = 2). Twenty-eight of the cyclists were currently competing internationally in their respective Union Cycliste Internationale (UCI) disciplines/categories, as well as training on a full-time basis for at least the past two years. More specifically, their collective experience and success included: 2 Olympic medals, 8 Olympic games representations, 3 Paralympic medals, 3 Paralympic games representations (as a pilot [who are able-bodied and compete individually at national and up to UCI Class 1 level events] or stoker [other than being visually impaired, they are fully able-bodied] of tandem), 10 Senior World Championship medals, 37 Senior World Championships representations, 8 Senior Para-cycling World Championship golds and 6 Senior Para-cycling World Championship medals (as pilot or stoker of tandem). The remaining seven participants who were not competing internationally were competing in 'Elite' category national level road cycling events (n = 4) or had won national level medals on the track (n = 3). Ethical approval was attained from Northumbria University Research Ethics Committee. Following explanation of the study design and protocol, the cyclists provided written, informed consent to prior to their participation in the study.

### *Study Overview*

Before all experimental sessions at the laboratory, cyclists were instructed to avoid caffeine and food for 3 h prior to testing and to avoid strenuous exercise in the 24 h before each session. Cyclists made two identical cycling laboratory visits within 7 days at the same time of day ( $\pm 1$  hour). Firstly, the cyclists assumed their race cycling specific position, on a custom-built ergometer that had the cast flywheel clamped, to ensure the cranks were stationary. In this position, architecture of the vastus lateralis (i.e. pennation angle [ $P\theta_{VL}$ ] and fascicle length [ $FL_{VL}$ ]) was assessed at rest prior to exercise using ultrasound<sup>23</sup>. Subsequently, the cyclists had surface electromyography (EMG) electrodes placed on three muscles of both legs (gluteus maximus [GM], biceps femoris long head [BF], and vastus lateralis [VL]) and mounted another custom-modified isovelocity ergometer (again, the position mirrored their racing position). A standardised warm-up of 10 mins at 80 – 90 revolutions per minute (RPM) and 100 – 150 W followed by a maximal 2 s sprint at 125 RPM was completed by each cyclist. Once this was completed a series of isovelocity sprints (4 s maximal sprints at each of five velocities: 60, 115, 125, 135 and 180 RPM) was performed in this order to assess PPO, as well as torque, whilst surface EMG was simultaneously recorded. On a third occasion, within 7 days of the cycling laboratory visits, MRI was used to assess quadricep femoris and hamstrings muscle volume of both legs of each cyclist.

### *Muscle Architecture*

For the architecture measures a custom-built cycling ergometer (United Kingdom Sports Innovation) was set-up according to the individual cyclist's track or road bike set-up. BMX riders had experience in riding road and track bikes were set-up to their track or road bike position fitted which was typical to a track cycling set-up when assuming the dropped handle bar position (i.e. closed hip angle, flat back parallel with the floor and bent arms). The ergometer could be made isometric, as previously used.<sup>16</sup> Before the cyclist mounted the

ergometer, bib shorts were pulled up to expose their thighs in order to allow mid-thigh to be measured and marked. When the cyclists first mounted the ergometer for the ultrasound imaging, the flywheel was clamped to ensure that the crank position was fixed with the drive-side (right) crank positioned at 90° from top, dead centre (TDC). Once in this position, the cyclists were asked to take their racing position with their hands on the ‘drops’.

An ultrasound (5-10 MHz scanning width 92 mm and depth 65 mm, EUP-L53L; Hitachi EUB-8500) linear array transducer was used to capture B-mode ultrasound images. Water-soluble transmission gel was used to coat the transducer that was positioned with minimal pressure over the skin. Images were captured with the transducer placed on the medial, longitudinal line of the muscle while positioned on the skin over the VL at 50% of femur length (from the knee joint space to the greater trochanter) to correspond with the area of greatest anatomical CSA<sup>24</sup>. The transducer was orientated perpendicular to the skin and parallel to the fascicular path. Parallel fascicle alignment was presumed when transducer orientation produced an image whereby the aponeuroses and the fascicle perimysium trajectory were clearly identified with no visible fascicle distortion at the image edges. Once the images were captured, the cyclists were instructed to switch lead legs and have the non-drive side (left) crank positioned at 90° from TDC and the process was repeated with the left VL. Images were later imported into analysis software (ImageJ, v.1.46; National Institutes of Health, Bethesda, MD, USA) to measure  $FL_{VL}$  and  $P\theta_{VL}$ . The  $FL_{VL}$  was measured as the length of the fascicular path between the superficial and deep aponeurosis. The manual (fascicular line tracing) linear extrapolation approach was adopted when the full fascicle length could not be seen within the ultrasound image. The  $P\theta_{VL}$  was measured as the angle between the fascicular path and the insertion of fascicles into the deep aponeurosis. Three different ultrasound images of each leg were recorded and analysed during each visit before first averaging the measured values from each session, and then averaging across the two sessions. The intra-rater repeatability of

measures  $P\theta_{VL}$  had CV of 4.1% and ICC of 0.86. and  $FL_{VL}$  had a CV of 1.9% and ICC of 0.98 and within-participant repeatability  $P\theta_{VL}$  had CV of 2.9% and ICC of 0.91. and  $FL_{VL}$  had a CV of 1.3% and ICC of 0.97.

### *Electromyography*

A wireless, surface EMG system (Delsys Trigno® Wireless EMG systems, Boston, MA, USA) was used to ascertain muscle activation by measuring EMG amplitude. Once muscle architecture assessment was complete, EMG electrodes were placed (in accordance with standard SENIAM recommendations<sup>25</sup> on each leg over the GM, BF and VL. Each location was shaved, lightly abraded, and then cleaned with a sterilised alcohol wipe. To ensure optimal electrical conductance, the electrodes were then applied using self-adhesive interfaces (Delsys Trigno®, Boston, MA, USA), each site was marked with a semi-permanent marker to ensure consistent placement across sessions. Surface EMG signals were sampled at 2,000 Hz, amplified ( $\times 1000$ ), band-pass filtered (20-450 Hz) using an external analogue-to-digital data acquisition system (Micro 1401, Cambridge Electronic Design, Cambridge, UK) and a PC utilizing Spike2 software (version 7.11, CED, Cambridge, UK).

EMG signal amplitudes were calculated as the root mean square (rmsEMG) over an epoch equivalent to one quarter of the full crank cycle for all isovelocity cadences (e.g. 250 ms at 60 RPM). Peak rmsEMG values of each muscle (GM, VL, BF) were averaged over the first three full crank revolutions (TDC to TDC) of each sprint/velocity. The peak rmsEMG amplitude during the 60 RPM isovelocity sprint was used to normalise peak rmsEMG from the isovelocity sprint with the highest measured PPO (peak rms EMG<sub>PPO</sub>) i.e. peak rmsEMG<sub>PPO</sub>/peak rmsEMG<sub>60</sub> and used as criterion values of activation of each muscle at PPO (GM<sub>ACT</sub>, VL<sub>ACT</sub>, BF<sub>ACT</sub>). From the data previously collected in our lab the coefficient of

variation for between-session peak rmsEMG reliability during isovelocity cycling at 60 RPM as a reference task for GM, VL and BF was 9.9, 13.0 and 9.5%, respectively.

### *Sprint Cycling Performance Test*

Isovelocity sprints were performed on an SRM ergometer (Schoberer Rad Messtechnik, Jülich, Germany) that was modified to have a motor accelerate the flywheel to the prescribed cadence for isovelocity sprints. As with the custom-built ergometer used for muscle architecture, the protocol on isovelocity SRM ergometer was identically set-up to their racing positions. All efforts were performed in the saddle. The cyclists were instructed to perform all the efforts in the saddle. This were only monitored by the investigator and no restraint or strapping was used to ensure they remained attached to the saddle. Each cyclist performed each effort on ‘drop’ handlebars, using clipless pedals and racing attire. The original cranks were replaced with 170 mm instrumented cranks to record instantaneous torque, crank angle, and angular velocity directly as raw data for both right and left crank arms (Factor Cranks, BF1 Systems, Diss, UK), which were sampled at 200 Hz with a separate, wireless electronic data logger (BF1 Systems, Diss, UK). Torque and power was measured over each revolution and was processed offline as previously done.<sup>16</sup> Raw data recorded from the cranks onto the wireless data logger were imported in to Spike2 and analysed offline using custom scripts to calculate mean torque, power and cadence per revolution from TDC to TDC.

Prior to performing the maximal isovelocity efforts to determine PPO, power-cadence and torque-cadence relationship, cyclists undertook a standard 10-min warm-up of submaximal cycling at a self-selected intensity (between 100–150 W) and cadence (between 80–90 RPM) with a 2 s maximal effort at 125 RPM. For the maximal isovelocity efforts, participants performed 4 s sprints at 60, 115, 125, 135 and 180 RPM. The order of cadences was selected at random and every cyclists performed the efforts in the following order: 115, 60, 135, 125

and 180 RPM. Prior to each effort, the motor was brought up to the desired velocity and participants were instructed to pedal below the pre-set cadence and reminded to ‘attack the effort as fast and as hard as possible’. Then the investigator gave a 3 s countdown and the participants performed a 4 s maximal effort at the set cadence. Each isovelocity effort was performed once per lab visit with a 5 min period of passive rest between each isovelocity sprint.

The maximum power output over three consecutive revolutions (from TDC to TDC) at each cadence was used and then averaged over both sessions. From that, power-cadence and torque-cadence relationships were established by fitting a quadratic and linear equation, respectively, by the least square method as previously used.<sup>4,26</sup> The apex of the power-cadence relationship was interpolated to derive PPO (as well as PPO: mass by dividing PPO by body mass [W/Kg]) and cadence at PPO ( $C_{OPT}$ ). Individual torque-cadence relationships, maximal torque ( $T_{MAX}$ ) and maximal cadence ( $C_{MAX}$ ) were extrapolated.

### *MR Imaging*

On a separate occasion, within 7 days of the cycling laboratory visits, muscle volume of both legs was measured via MR imaging (1.5 T Signa HDxt; Alliance Medical Limited, Warwick, UK). T1-weighted axial images of each thigh originating at the anterior-superior iliac spine and finishing at the knee joint space (scan parameters: time of repetition = 600 ms; time to echo = 14 ms; image matrix 512 pixels  $\times$  512 pixels; field of view 260 mm  $\times$  260 mm; slice thickness = 5 mm; and interslice gap = 5 mm). An array of fish-oil capsules were attached using micro-pore surgical tape on and around the anterior-superior iliac spine and knee joint space as previously done<sup>27</sup>. This was to help the operator orientate any overlapping blocks during the analysis stage. Participants were asked to refrain from exercise in the thirty-six hours before the scan and sat quietly for  $> 1$  h before the scan. Participants lay supine with legs fully

extended and strapped in position to discourage any extraneous movement that might cause image distortion.

Muscle volume was measured by an experienced operator, who was blinded to the participant's identity and performance data, using open source software (OsiriX Imaging Software™ version 5.5.1, Geneva, Switzerland). Volume was calculated by measuring anatomical (CSA), in the axial plane, by manual segmentation of VL, vastus intermedius (VI), vastus medialis (VM) and rectus femoris (RF) as well as semitendinosus (ST) and semimembranosus (SM), long and short head BF. In each individual image using the 'closed polygon' tool. Manual outlining started with the most distal slice above the knee, at where the muscles were visible, and ended with the most proximal slice where the muscle was no longer visible. The total number of slices was noted and used to determine the length of the segment (length =  $n \times 15$  mm, where  $n$  is the number of slices, given that MR image slices were 5 mm in thickness). On average, thirty images were analysed per thigh. Consequently, muscle volume was calculated by using Cavalieri formula: <sup>28</sup>

$$\text{Muscle Volume} = \sum_n e_i \times \text{CSA}_i$$

Where  $n$  is the number of slices used, and  $e_i$  is the distance between measured slices.

Knee extensor muscle volume ( $Q_{VOL}$ ) was measured by summing the muscle volume of VL, VM, VI and RF of each leg. Hamstring muscle volume ( $HAM_{VOL}$ ) was measured by summing the muscle volume of long and short head BF, semitendinosus and semimembranosus of each leg. Both  $Q_{VOL}$  and  $HAM_{VOL}$  were averaged over both legs.

### *Statistical Analysis*

All data is presented as mean  $\pm$  SD. A Shapiro-Wilk test of the measures showed that the data was normally distributed and suitable for parametric testing. Data from all thirty-five cyclists were used to perform bivariate correlations and subsequent regression analysis with the physiological measures. Initially, Pearson's product-moment correlations ( $r$ ) were employed to examine the relationship between individual physiological variables and the criterion variable (PPO). The following criteria were adopted to interpret the magnitude of the relationship between test measures:  $<0.1$  trivial,  $0.1$  to  $0.3$  small,  $>0.3$  to  $0.5$  moderate,  $>0.5$  to  $0.7$  large,  $>0.7$  to  $0.9$  very large, and  $>0.9$  to  $1.0$  almost perfect<sup>29</sup>. In addition, the overall coefficient of determination ( $R^2$ ) for the set of physiological measures with PPO was also calculated.

Variables that were significantly correlated with PPO were included in the step-wise regression analysis to predict PPO. With this set of predictors, our collinearity diagnostic exploration resulted in variance inflation factors of  $2.0$  -  $5.0$  and tolerance of  $0.20$  -  $0.80$ , which indicate acceptable levels of multicollinearity.<sup>30</sup>

The sprint and endurance groups within the whole cohort were compared using an independent-samples t-test for sprint performance measures (i.e. PPO, PPO: mass,  $C_{OPT}$ ,  $T_{MAX}$  and  $C_{MAX}$ ) and physiological measures (i.e.  $Q_{VOL}$ ,  $HAM_{VOL}$ ,  $P\theta_{VL}$ ,  $FL_{VL}$ ,  $GM_{ACT}$ ,  $VL_{ACT}$ ,  $BF_{ACT}$ ). In addition, independent-samples t-tests were also used to compare the volume of individual quadriceps femoris (i.e. VL, VI, VM and RF) and hamstrings (bicep femoris short head, bicep femoris long head, ST and SM) muscles. Finally, the relative/proportional volume (percentage) of the whole muscle group accounted for by each individual muscle within the quadriceps and hamstrings muscle groups was compared between sprint and endurance cyclists.

All physiological measures (mentioned above) were averaged over both limbs and then both sessions (with the exception of MR imaging). The level of statistical significance was set at  $P < 0.05$  for all tests. All statistics were calculated using SPSS (IBM Corp. Version 24.0. Armonk, USA).

### 3 RESULTS

Collectively for all thirty-five riders, the average  $\pm$  SD, range (i.e. maximum and minimum) and fold variability (multiple between maximum and minimum value) of the performance and physiological measures are presented in Table 1. Very large, positive bivariate relationships were found between  $Q_{VOL}$  ( $r = 0.87$ ;  $P < 0.001$ ),  $HAM_{VOL}$  ( $r = 0.71$ ;  $P < 0.001$ ) and  $P\theta_{VL}$  ( $r = 0.81$ ;  $P < 0.001$ ) with cycling PPO (Table 2; Figure 1). The remaining measures ( $FL_{VL}$ ,  $VL_{ACT}$ ,  $BF_{ACT}$  and  $GM_{ACT}$ ) were unrelated to PPO. Subsequently, step-wise multiple regression analysis was done using the three significant predictor variables from the bivariate correlations ( $Q_{VOL}$ ,  $HAM_{VOL}$ ,  $P\theta_{VL}$ ) to examine their combined relationship with PPO. The regression analysis found 87% of the variability in PPO between cyclists ( $F_{(2, 28)} = 72.83$ ,  $P < 0.001$ ) was explained by two variables  $Q_{VOL}$  (76%) and  $P\theta_{VL}$  (11%).

The comparison of power-cadence and torque-cadence relationships between the groups of sprint and endurance cyclists (Figure 2) showed that sprint cyclists had substantially higher PPO ( $\sim +579$  W; +47%;  $P < 0.001$ ), PPO: Mass ( $\sim +4.3$  W:Kg; +27%;  $P < 0.001$ ),  $C_{OPT}$  ( $\sim +11$  RPM; 8%;  $P < 0.05$ ),  $T_{MAX}$  ( $\sim +62$  N·m; +35%;  $P < 0.001$ ) and  $C_{MAX}$  ( $\sim +31$  RPM; +11%;  $P < 0.05$ ; Table 3). In terms of the physiological measures sprint cyclists had significantly larger  $Q_{VOL}$ ,  $HAM_{VOL}$ , volume of all the individual muscles of both groups and  $P\theta_{VL}$  (all  $P < 0.001$ ; Table 3; Figure 3). In terms of proportional volume of the individual muscles, all four quadriceps (VL, VM, VI & RF) as well as three of the hamstrings (ST, SM, BF-1), were similar for sprint and endurance cyclists, however biceps femoris short

head was proportionately larger in endurance vs sprint cyclists ( $17.5 \pm 3.1$  vs.  $15.1 \pm 2.8\%$ ;  $P = 0.0219$ ).

No significant differences were seen between groups when  $FL_{VL}$ ,  $VL_{ACT}$ , and  $GM_{ACT}$  were examined whilst the endurance cyclists exhibited higher  $BF_{ACT}$  ( $P < 0.05$ ) during sprint cycling.

#### 4 DISCUSSION

To the author's knowledge, this is the first study to forensically investigate the physiological attributes that determine PPO in an elite, highly-trained cycling cohort. The main finding of this study was the very large, positive relationships between  $Q_{VOL}$ ,  $HAM_{VOL}$  and  $P\theta_{VL}$  with PPO, with multiple regression showing that in combination  $Q_{VOL}$  and  $P\theta_{VL}$ , 87% of the variability in PPO between cyclists could be explained. These findings appear to agree with the hypothesis that muscle volume of the quadriceps femoris are the biggest (but not the only) predictors of PPO in sprint cycling. This demonstrates the importance of muscle morphology for sprint cycling performance. In contrast the remaining variables, particularly neuromuscular activation of three hip and knee joint muscles, showed no relationships with PPO. The secondary finding was that as expected elite sprint cyclists had substantially higher power-cadence and torque-cadence relationships (i.e. PPO, PPO: mass,  $C_{OPT}$ ,  $T_{MAX}$  and  $C_{MAX}$ ) than endurance cyclists and this was underpinned by greater  $Q_{VOL}$  (+52%),  $HAM_{VOL}$  (+52%) and  $P\theta_{VL}$  (+20%).

The sprint performance measures in the current study were similar to previous reports. For example the sprint group recorded PPO of  $1521 \pm 186$  W that was within 80 W of three previous studies using similar although somewhat smaller elite cohorts.<sup>4,31,32</sup> The endurance group in the current study had PPO of  $942 \pm 136$  W, similar to untrained cyclists ( $941 \pm 124$  W)<sup>7</sup> and somewhat lower than a previous elite endurance cohort ( $1122 \pm 65$  W).<sup>32</sup> Of the

physiological measures, the endurance cyclists had muscle volume and  $P\theta_{VL}$  similar to the untrained groups.<sup>27</sup> The endurance group had muscle volume similar to long-term resistance trained participants<sup>27</sup> and  $P\theta_{VL}$  similar to sprint runners.<sup>33</sup>

### *Morphological Determinants of Sprint Cycling Performance*

$Q_{VOL}$ ,  $HAM_{VOL}$  and  $P\theta_{VL}$  all showed very large, positive bivariate relationships with PPO, and multiple regression analysis found  $Q_{VOL}$  and  $P\theta_{VL}$  explained 87% of the variance in PPO, whereas the neuromuscular activation measures were unrelated to PPO. For the first time and in elite cyclists, this study shows that PPO is overwhelmingly determined by muscle morphology, particularly the size and pennation angle of the quadriceps, rather than the ability of the nervous system to activate the muscles.

Using MR imaging, a 'gold standard' method for determining muscle volume,<sup>17</sup> we found  $Q_{VOL}$  i.e. the amount of skeletal muscle, alone explained 76% of the variance in PPO between cyclist, which makes this variable a desirable attribute for competitive (sprint) cyclists. Whilst no previous studies have carefully imaged the quadriceps and hamstrings muscles in relation to sprint cycling performance some crude estimates lower body/thigh muscle mass have been found to be moderately/strongly related to cycling PPO.<sup>4,7,34</sup> Our findings for the predominant influence of  $Q_{VOL}$  on PPO, reinforce the importance of muscle size for neuromuscular power and that cyclists and their coaches be especially attentive to training and nutrition strategies to enhance  $Q_{VOL}$ . In particular resistance training is well known to stimulate hypertrophy and increased muscle volume. Elite sprint cyclists in the current study had slightly smaller  $Q_{VOL}$  than a long-term (mean 4-years) resistance trained, but not elite, cohort assessed with an almost identical MRI protocol.<sup>27</sup> This could be due to the concurrent nature of performing both sprint cycling training in conjunction with resistance exercises that could attenuate hypertrophic responses.<sup>35</sup>

$P\theta_{VL}$  was a strong correlate of PPO in the current study ( $r = 0.81$ ) and given the relationship of muscle volume and PPO this might have been expected as pennation angle is known to be associated with muscle size indices (e.g. <sup>36</sup>) and this was also the case in the current study ( $Q_{VOL}$  vs  $P\theta_{VL}$   $r = 0.78$ ). However, what was perhaps more surprising was that  $P\theta_{VL}$  was an independent predictor of PPO in addition to  $Q_{VOL}$  within the regression analysis. Suggesting that a high  $P\theta_{VL}$  is advantageous for neuromuscular power even after muscle volume has been accounted for. This may reflect a net positive balance of advantages and disadvantages of increasing  $P\theta_{VL}$  at the relatively low angles found in this study ( $< 20^\circ$ ). For PPO, theoretically greater  $P\theta_{VL}$  has the advantage of greater PCSA and thus higher force production capacity but potential disadvantages of loss of force transmission to the tendon and/or reduction in fascicle length and thus shortening velocity. In the current study PPO was unrelated to FL, which is somewhat contrary to the findings of a positive association of FL with sprint running (100 m) performance. <sup>33</sup> Furthermore,  $FL_{VL}$  was also unrelated to  $P\theta_{VL}$  ( $r = -0.23$ ), suggesting no negative consequence of increasing  $P\theta_{VL}$  on  $FL_{VL}$ .

Data in this experiment indicated that peak muscle activation recorded with surface EMG exhibited no relationship with PPO, and thus was not a meaningful determinant of PPO in elite cyclists. Therefore, it is possible that more accurate and sensitive measures of neuromuscular activation, perhaps including surface EMG from more muscles and multiple sites per muscle <sup>37</sup>, as well as alternative normalisation techniques, <sup>38</sup> might reveal a greater role for activation in determining PPO. However, given that muscle morphology explained 87% of the variability in PPO the unexplained variance was relatively small (13%) and the contribution of other independent factors, including neuromuscular activation appears limited for this performance task.

*Magnitude of Morphological measures between Sprint cyclists and endurance cyclists*

Sprint cyclists were greater in every measure of the sprint cycling performance test (i.e. PPO +61%, PPO: Mass +32%,  $T_{MAX}$  +43%,  $C_{OPT}$  +9% and  $C_{MAX}$  +12%) in comparison to endurance riders which is likely to be the consequence of sprint riders being greater in all the morphological measures that had a positive and significant relationship with PPO (i.e.  $Q_{VOL}$ ,  $HAM_{VOL}$  and  $P\theta_{VL}$ ). This perhaps gives further weight that the mechanisms of PPO.

The greater PPO of sprint cyclists (+61% vs endurance cyclists) appeared to be primarily due to their higher  $T_{MAX}$  (+43%), as opposed to a smaller difference in  $C_{OPT}$  (+9%). The greater  $T_{MAX}$  of the sprint cyclists was likely due to their greater  $Q_{VOL}$  and  $HAM_{VOL}$  as a greater muscle volume provides more sarcomeres in parallel that can exert a greater force/torque (the relationship between the sum of  $Q_{VOL}$  and  $HAM_{VOL}$  with PPO:  $r = 0.81$ ). No differences were seen for  $FL_{VL}$  which further suggested that it may not be an important morphological determinant of sprint cycling ability.

The focus of this study has been the relationships between muscle morphology (size and architecture) with cycling PPO. The portion of the PPO differences that were not accounted for in this study could be explained by either, or a combination of, muscle fibre-type composition and muscle fibre contraction speed. With respect to skeletal muscle fibre type composition, it is thought that a higher proportion of type II muscle fibres have a substantial influence on muscle fibre PPO, primarily due to their higher maximum shortening velocity which is underpinned by the higher quantities of ATPase<sup>39</sup>. The higher PPO as well as  $C_{OPT}$  and  $C_{MAX}$  in sprint cyclists could, in part, be attributed to a greater proportion of type II muscle fibres. Whilst there is relatively limited data in humans, one small study ( $n=10$ ) reported a strong positive relationship ( $r = 0.88$ ;  $R^2 = 0.77$  %) between the proportion or number of fast-twitch fibres and  $C_{OPT}$ . As such, it is possible that muscle fibre type composition, and thus maximum shortening velocity, could contribute to higher PPO of sprint cyclists, but needs to be examined further in future studies in athletic cohorts, although this is

unlikely to be achieved in an elite athletic cohort of this level. This idea is underpinned by first principle models from Hill <sup>41,42</sup> and Huxley <sup>43</sup>, suggesting that fibre speeds (rather than fibre type) also could account for some differences in PPO in sprint cycling. Force-velocity characteristics of muscle are based on kinetics of cycling interaction between myosin cross-bridges and actin filaments within sarcomeres of muscle. Higher muscle fibre speeds have higher cross-bridge cycling rates that underpin muscle fibre speeds which could explain the higher torque offsets (and  $C_{OPT}$ ) in sprint cyclists across the cadence ranges in comparison to the endurance cyclists.

The sprint cyclists also measured higher muscle volume in each individual muscle of the quadriceps femoris and hamstrings. Proportional muscle volume was also similar between sprint and endurance cyclists in all four individual quadriceps muscles and three out of four hamstrings muscles. The exception was that the bicep femoris short head, the smallest muscle examined in this study, which was proportionately larger in endurance than sprint cyclists. Therefore, only very limited evidence for muscle-specific regional hypertrophy / muscle mass distribution in one of the 8 muscles examined was seen. A previous study by Ema and colleagues (2016) suggested that in comparison to untrained men, experienced cyclists who were club level track cyclists who competed in sprint events had higher muscle volumes of the bicep femoris short head and semitendinosus. But no difference in the percentage of each individual muscle in the hamstrings were seen. The finding in the current study could suggest that the short head of the bicep femoris short head may have more of a role to play in endurance cycling than in sprint cycling.

### *Limitations of the Study*

Although the data collection within the current study was extensive, there were a number of limitations associated with the methodology. Firstly, the selection of two different,

highly specialised and distinct groups of cyclists may have created coincidental or exaggerated relationships by having big differences between groups for a whole cluster of variables (both assessed variables e.g. PPO and muscle volume, but also potentially unassessed variables e.g. fibre type composition, tendon stiffness). Therefore, the strength of the relationships between predictor and outcome variables in this study may actually be reflective of a range of predictor variables. Secondly, correlation does not demonstrate causality, and therefore whilst there were very strong relationships between  $Q_{VOL}$  and  $P\theta_{VL}$  with PPO, and these factors in combination explained 87% of the variation of PPO, this does not necessarily mean that changes in one or both predictor variables will cause a proportionate increase in PPO. On the basis of the interesting findings of the current work it is recommended that future studies use a wider range of predictor variables (e.g. fibre type composition, or surrogate measures of contractile properties, and tendon stiffness) for cross-sectional analyses and particularly that intervention studies examine the effect of changing muscle morphology on cycling PPO.

This study largely focused on measures in the thigh as previous studies suggest that the physiological determinants of PPO are largely rooted there.<sup>15,16</sup> Other muscle groups that almost certainly contribute to cycling PPO such as gluteus maximus and plantar flexors<sup>31,45</sup> were not assessed. Assessment of other major muscle groups would have given a more complete understanding of physiological determinants of PPO. Furthermore, the VL is a major muscle in PPO production<sup>8</sup> and assessment of VL muscle architecture has been used extensively.<sup>46</sup> However, these images only capture a superficial, two-dimensional representation of the muscle, which may not be representative of the whole muscle or other groups of muscles. A future study could examine multiple muscles involved in cycling), and a range of sites within each muscle in order to further investigate the relationship of muscle architecture and PPO.

## 5 PERSPECTIVES

These new data showed quadriceps femoris muscle volume and pennation angle accounted for 76 and 11 %, respectively, of the variance of PPO in elite cyclists. These findings emphasise the importance of quadriceps muscle morphology for sprint cycling events and reinforce that cyclists and their coaches should be attentive to maximising these characteristics during their preparation and training. In addition, sprint cyclists achieved higher PPO than endurance cyclists, with  $T_{MAX}$  appearing to be the primary explanation for their greater PPO, which was likely because of their greater muscle morphology ( $Q_{VOL}$ ,  $HAM_{VOL}$  and  $P\theta_{VL}$ ). Additionally, future studies would need to measure changes in these morphological measures (throughout the course of a season) and whether these can predict changes in PPO and/or power- and torque-cadence relationships.

## **CONFLICT OF INTEREST**

All authors declare that there is no conflict of interest. The authors alone are responsible for the content and writing of the manuscript.

## REFERENCES

1. Williams JH, Barnes WS, Signorile JF. A constant-load ergometer for measuring peak power output and fatigue. *J Appl Physiol Bethesda Md* 1985 1988;65:2343–2348.
2. Cheetham ME, Boobis LH, Brooks S, Williams C. Human muscle metabolism during sprint running. *J Appl Physiol Bethesda Md* 1985 1986;61:54–60.
3. Bosco C, Luhtanen P, Komi PV. A simple method for measurement of mechanical power in jumping. *Eur J Appl Physiol* 1983;50:273–282.
4. Dorel S, Hautier CA, Rambaud O, Rouffet D, Van Praagh E, Lacour JR, Bourdin M. Torque and power-velocity relationships in cycling: relevance to track sprint performance in world-class cyclists. *Int J Sports Med* 2005;26:739–746.
5. Martin JC, Gardner AS, Barras M, Martin DT. Modeling sprint cycling using field-derived parameters and forward integration. *Med Sci Sports Exerc* 2006;38:592–597.
6. Martin JC, Davidson CJ, Pardyjak ER. Understanding sprint-cycling performance: the integration of muscle power, resistance, and modeling. *Int J Sports Physiol Perform* 2007;2:5–21.
7. Leong CH, McDermott WJ, Elmer SJ, Martin JC. Chronic eccentric cycling improves quadriceps muscle structure and maximum cycling power. *Int J Sports Med* 2014;35:559–565.
8. Akima H, Kinugasa R, Kuno S. Recruitment of the thigh muscles during sprint cycling by muscle functional magnetic resonance imaging. *Int J Sports Med* 2005;26:245–252.
9. Craig NP, Norton KI. Characteristics of track cycling. *Sports Med Auckl NZ* 2001;31:457–468.
10. Gardner SA, Martin TD, Barras M, Jenkins GD, Hahn GA. Power output demands of elite track sprint cycling. *Int J Perform Anal Sport* 2005;5:149–154.
11. Jones D, Round JM, Haan A de. *Skeletal muscle from molecules to movement: a textbook of muscle physiology for sport, exercise, physiotherapy and medicine.* Edinburgh: Churchill Livingstone; 2004. 202 p.
12. O'Brien TD, Reeves ND, Baltzopoulos V, Jones DA, Maganaris CN. Strong relationships exist between muscle volume, joint power and whole-body external mechanical power in adults and children: Development of joint and external mechanical power with adolescence. *Exp Physiol* 2009;94:731–738.
13. Driss T, Vandewalle H. The measurement of maximal (anaerobic) power output on a cycle ergometer: a critical review. *BioMed Res Int* 2013;2013:589361.
14. Rønnestad BR, Hansen EA, Raastad T. Effect of heavy strength training on thigh muscle cross-sectional area, performance determinants, and performance in well-trained cyclists. *Eur J Appl Physiol* 2010;108:965–975.

15. Driss T, Vandewalle H, Le Chevalier JM, Monod H. Force-velocity relationship on a cycle ergometer and knee-extensor strength indices. *Can J Appl Physiol Rev Can Physiol Appl* 2002;27:250–262.
16. Kordi M, Goodall S, Barratt P, Rowley N, Leeder J, Howatson G. Relation between Peak Power Output in Sprint Cycling and Maximum Voluntary Isometric Torque Production. *J Electromyogr Kinesiol Off J Int Soc Electrophysiol Kinesiol* 2017;35:95–99.
17. Engstrom CM, Loeb GE, Reid JG, Forrest WJ, Avruch L. Morphometry of the human thigh muscles. A comparison between anatomical sections and computer tomographic and magnetic resonance images. *J Anat* 1991;176:139–156.
18. Aagaard P, Andersen JL, Dyhre-Poulsen P, Leffers AM, Wagner A, Magnusson SP, Halkjaer-Kristensen J, Simonsen EB. A mechanism for increased contractile strength of human pennate muscle in response to strength training: changes in muscle architecture. *J Physiol* 2001;534:613–623.
19. Blazevich AJ, Coleman DR, Horne S, Cannavan D. Anatomical predictors of maximum isometric and concentric knee extensor moment. *Eur J Appl Physiol* 2009;105:869–878.
20. Lieber RL, Fridén J. Functional and clinical significance of skeletal muscle architecture. *Muscle Nerve* 2000;23:1647–1666.
21. Van Cutsem M, Duchateau J, Hainaut K. Changes in single motor unit behaviour contribute to the increase in contraction speed after dynamic training in humans. *J Physiol* 1998;513 ( Pt 1):295–305.
22. Aagaard P, Simonsen EB, Andersen JL, Magnusson P, Dyhre-Poulsen P. Increased rate of force development and neural drive of human skeletal muscle following resistance training. *J Appl Physiol Bethesda Md* 1985 2002;93:1318–1326.
23. Kordi M, Menzies C, Parker Simpson L. Relationship between power-duration parameters and mechanical and anthropometric properties of the thigh in elite cyclists. *Eur J Appl Physiol* 2018;118:637–645.
24. Erskine RM, Jones DA, Maganaris CN, Degens H. In vivo specific tension of the human quadriceps femoris muscle. *Eur J Appl Physiol* 2009;106:827–838.
25. Hermens HJ, Freriks B, Disselhorst-Klug C, Rau G. Development of recommendations for SEMG sensors and sensor placement procedures. *J Electromyogr Kinesiol Off J Int Soc Electrophysiol Kinesiol* 2000;10:361–374.
26. Gardner AS, Martin JC, Martin DT, Barras M, Jenkins DG. Maximal torque- and power-pedaling rate relationships for elite sprint cyclists in laboratory and field tests. *Eur J Appl Physiol* 2007;101:287–292.
27. Massey GJ, Balshaw TG, Maden-Wilkinson TM, Folland JP. Tendinous tissue properties after short- and long-term functional overload: Differences between controls, 12 weeks and 4 years of resistance training. *Acta Physiol Oxf Engl* 2018;222:e13019.

28. Lund H, Christensen L, Savnik A, Boesen J, Danneskiold-Samsøe B, Bliddal H. Volume estimation of extensor muscles of the lower leg based on MR imaging. *Eur Radiol* 2002;12:2982–2987.
29. Hopkins WG. Spreadsheets for analysis of validity and reliability. *Sportscience* 2015;19:36–42.
30. Hair JF Jr, Anderson R, Tatham R, Black WC. *Multivariate Data Analysis*. New York: Macmillan; 1995.
31. Dorel S, Guilhem G, Couturier A, Hug F. Adjustment of muscle coordination during an all-out sprint cycling task. *Med Sci Sports Exerc* 2012;44:2154–2164.
32. Calbet JAL, De Paz JA, Garatachea N, Cabeza de Vaca S, Chavarren J. Anaerobic energy provision does not limit Wingate exercise performance in endurance-trained cyclists. *J Appl Physiol* 2003;94:668–676.
33. Abe T, Kumagai K, Brechue WF. Fascicle length of leg muscles is greater in sprinters than distance runners. *Med Sci Sports Exerc* 2000;32:1125–1129.
34. Maciejczyk M, Wiecek M, Szymura J, Szygula Z, Brown LE. Influence of increased body mass and body composition on cycling anaerobic power. *J Strength Cond Res* 2015;29:58–65.
35. Coffey VG, Hawley JA. Concurrent exercise training: do opposites distract?: Concurrent exercise training. *J Physiol* 2017;595:2883–2896.
36. Fukunaga T, Kawakami Y, Kuno S, Funato K, Fukashiro S. Muscle architecture and function in humans. *J Biomech* 1997;30:457–463.
37. Balshaw TG, Fry A, Maden-Wilkinson TM, Kong PW, Folland JP. Reliability of quadriceps surface electromyography measurements is improved by two vs. single site recordings. *Eur J Appl Physiol* 2017;117:1085–1094.
38. Lanza MB, Balshaw TG, Massey GJ, Folland JP. Does normalization of voluntary EMG amplitude to  $M_{MAX}$  account for the influence of electrode location and adiposity? *Scand J Med Sci Sports* 2018;
39. Bottinelli R, Pellegrino MA, Canepari M, Rossi R, Reggiani C. Specific contributions of various muscle fibre types to human muscle performance: an in vitro study. *J Electromyogr Kinesiol Off J Int Soc Electrophysiol Kinesiol* 1999;9:87–95.
40. Hautier CA, Linossier MT, Belli A, Lacour JR, Arsac LM. Optimal velocity for maximal power production in non-isokinetic cycling is related to muscle fibre type composition. *Eur J Appl Physiol* 1996;74:114–118.
41. Hill AV. The Heat of Shortening and the Dynamic Constants of Muscle. *Proc R Soc B Biol Sci* 1938;126:136–195.
42. Hill AV. THE EFFECT OF LOAD ON THE HEAT OF SHORTENING OF MUSCLE. *Proc R Soc Lond B Biol Sci* 1964;159:297–318.

43. Huxley AF. Muscle structure and theories of contraction. *Prog Biophys Biophys Chem* 1957;7:255–318.
44. Ema R, Wakahara T, Yanaka T, Kanehisa H, Kawakami Y. Unique muscularity in cyclists' thigh and trunk: A cross-sectional and longitudinal study: Muscle-specific adaptation to cycling. *Scand J Med Sci Sports* 2016;26:782–793.
45. Elmer SJ, Barratt PR, Korff T, Martin JC. Joint-specific power production during submaximal and maximal cycling. *Med Sci Sports Exerc* 2011;43:1940–1947.
46. Kwah LK, Pinto RZ, Diong J, Herbert RD. Reliability and validity of ultrasound measurements of muscle fascicle length and pennation in humans: a systematic review. *J Appl Physiol Bethesda Md* 1985 2013;114:761–769.

**Table 1:** Sprint cycling performance and physiological measures for thirty-five elite cyclists. Data are mean  $\pm$  SD, range and fold variability for minimum to maximum values.: Peak power output (PPO), optimal cadence ( $C_{OPT}$ ), extrapolated maximum torque ( $T_{MAX}$ ), extrapolated maximal cadence ( $C_{MAX}$ ), quadriceps muscle volume ( $Q_{VOL}$ ), hamstring muscle volume ( $HAM_{VOL}$ ), pennation angle of the vastus lateralis ( $PA_{VL}$ ) and fascicle length of vastus lateralis ( $FL_{VL}$ ).


	<b>Mean <math>\pm</math> SD</b>	<b>Range (max - min)</b>	<b>Fold variability</b>
<b>PPO (W)</b>	1240 $\pm$ 335	2025 - 775	x2.6
<b><math>C_{OPT}</math> (RPM)</b>	131 $\pm$ 12	161 - 112	x1.3
<b><math>T_{MAX}</math> (N.m)</b>	175 $\pm$ 37	236 - 117	x2.0
<b><math>C_{MAX}</math> (RPM)</b>	267 $\pm$ 31	362 - 221	x1.6
<b><math>Q_{VOL}</math> (cm<sup>3</sup>)</b>	2268 $\pm$ 582	3343 - 1347	x2.5
<b><math>HAM_{VOL}</math> (cm<sup>3</sup>)</b>	804 $\pm$ 206	1263 - 348	x3.6
<b><math>P\theta_{VL}</math> (°)</b>	15.6 $\pm$ 2.0	18.8 - 11.7	x1.6
<b><math>FL_{VL}</math> (cm)</b>	7.6 $\pm$ 0.7	9.0 - 6.5	x1.4
<b><math>GM_{ACT}</math> (%)</b>	102 $\pm$ 16	128 – 62	x2.1
<b><math>VL_{ACT}</math> (%)</b>	97 $\pm$ 15	137 – 69	x2.0
<b><math>BF_{ACT}</math> (%)</b>	95 $\pm$ 11	120 - 75	x1.6

**Table 2:** Bivariate relationships ( $r$ ) and associated coefficient of determination ( $R^2$ ) for a range of physiological measures and the criterion measure (peak power output) in elite cyclists ( $n = 35$ ). Knee extensor muscle volume ( $Q_{VOL}$ ); knee flexor muscle volume ( $HAM_{VOL}$ ); pennation angle ( $P\theta_{VL}$ ); fascicle length ( $FL$ ); gluteus maximus ( $GM_{ACT}$ ); vastus lateralis ( $VL_{ACT}$ ) and bicep femoris (long head) ( $BF_{ACT}$ ) muscle activation.


	<b>r</b>	<b>R<sup>2</sup></b>	<b>Relationship</b>	<b>p</b>
<b>Q<sub>VOL</sub></b>	0.87	76%	Very large	<0.001
<b>HAM<sub>VOL</sub></b>	0.72	50%	Very large	<0.001
<b>P<math>\theta_{VL}</math></b>	0.81	66%	Very large	<0.001
<b>FL<sub>VL</sub></b>	-0.15	2%	Small	0.933
<b>GM<sub>ACT</sub></b>	0.21	4%	Small	0.276
<b>VL<sub>ACT</sub></b>	-0.01	0%	Trivial	0.977
<b>BF<sub>ACT</sub></b>	-0.29	9%	Small	0.107

**Table 3:** Sprint cycling performance and physiological measures of sprint and endurance cyclists. Performance measures: peak power output (PPO), PPO normalised to body mass (PPO: Mass), optimal cadence ( $C_{OPT}$ ), extrapolated maximal torque ( $T_{MAX}$ ) and extrapolated maximal cadence ( $C_{MAX}$ ). Physiological measures: Knee extensor muscle volume ( $Q_{VOL}$ ); hamstring muscle volume ( $HAM_{VOL}$ ); pennation angle ( $P\theta_{VL}$ ); fascicle length of VL ( $FL_{VL}$ ); gluteus maximus ( $GM_{ACT}$ ); vastus lateralis ( $VL_{ACT}$ ) and bicep femoris (long head) ( $BF_{ACT}$ ) muscle activation. \* denotes significantly higher than endurance ( $p < 0.05$ ); \*\* denotes significantly higher than endurance ( $p < 0.001$ ); † denotes significantly higher than sprint ( $p < 0.05$ ).


	<b>Sprint (n = 18)</b>	<b>Endurance (n = 17)</b>
<b>PPO (W)</b>	1521 ± 186 **	942 ± 136
<b>PPO: Mass (W/kg)</b>	17.9 ± 1.9 **	13.6 ± 1.6
<b><math>C_{OPT}</math> (RPM)</b>	136 ± 14 *	125 ± 7
<b><math>T_{MAX}</math> (N.m)</b>	205 ± 18 **	143 ± 20
<b><math>C_{MAX}</math> (RPM)</b>	282 ± 84 *	251 ± 18
<b><math>Q_{VOL}</math> (cm<sup>3</sup>)</b>	2723 ± 420**	1786 ± 229
<b>VL</b>	961 ± 158**	429 ± 61
<b>VM</b>	662 ± 106**	503 ± 75
<b>VI</b>	762 ± 142**	662 ± 93
<b>RF</b>	337 ± 53**	232 ± 42
<b><math>HAM_{VOL}</math> (cm<sup>3</sup>)</b>	994 ± 176**	655 ± 108
<b>BF-s</b>	141 ± 32*	115 ± 29
<b>BF-l</b>	255 ± 55**	171 ± 30
<b>ST</b>	262 ± 60**	170 ± 29
<b>SM</b>	285 ± 71**	199 ± 47
<b><math>P\theta_{VL}</math> (°)</b>	17.1 ± 1.0**	14.8 ± 1.5
<b><math>FL_{VL}</math> (cm)</b>	7.6 ± 0.6	7.5 ± 0.7
<b><math>GM_{ACT}</math> (%)</b>	103 ± 16	101 ± 16
<b><math>VL_{ACT}</math> (%)</b>	99 ± 18	96 ± 12
<b><math>BF_{ACT}</math> (%)</b>	91 ± 9	98 ± 11†



**Figure 1:** Scatter plots showing the relationships between cycling peak power output (PPO) and different physiological measures: (a) quadriceps muscle volume (b) pennation angle, (c) fascicle length and (d) hamstrings muscle volume.



**Figure 2:** (a) Quadratic relationship of the power-cadence relationships and (b) inverse, linear torque-cadence relationship of sprint (red) and endurance (blue) cyclists. For (a) filled dots represent mean power output at each pre-determined cadence. Peak power output (PPO) and optimal cadence ( $C_{OPT}$ ) are also highlighted. Significant differences were measured between PPO and  $C_{OPT}$  of both groups. For (b), filled dots represent mean torque output at each pre-determined cadence. Extrapolated maximum torque ( $T_{MAX}$ ) and ( $C_{MAX}$ ) are shown for both groups. Significant differences were measured between  $T_{MAX}$  and  $C_{MAX}$  of both groups; Solid lines represent the mean relationship of measured cadences and dotted line represents extrapolation; shaded areas represent standard deviation at each pre-determined cadence which mirror the same relationships of respective relationships.



1

**Figure 3:** A comparison of proportional volume of individual muscles (% of whole muscle group) between sprint (n=17) and endurance cyclists (n=18) for (a) quadriceps femoris muscles: vastus intermedialis (VI), vastus medialis (VM), vastus lateralis (VL) and rectus femoris (RF) and (b) hamstrings muscles: long head bicep femoris (BF (LONG)), short head bicep femoris (BF (SHORT)), semitendinosus (ST) and semimembranosus (SM). Data presented as mean  $\pm$  SD; \* denotes significant difference between proportion of muscle groups between sprinters and endurance riders.