

Northumbria Research Link

Citation: El, Gennady (2006) Erratum: "Resolution of a shock in hyperbolic systems modified by weak dispersion" [Chaos 15, 037103 (2005)]. Chaos: An Interdisciplinary Journal of Nonlinear Science, 16 (2). 029901. ISSN 1054-1500

Published by: American Institute of Physics

URL: <https://doi.org/10.1063/1.2186766> <<https://doi.org/10.1063/1.2186766>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/42797/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

Erratum: “Resolution of a shock in hyperbolic systems modified by weak dispersion” [Chaos 15, 037103 (2005)]

G. A. El

Department of Mathematical Sciences, Loughborough University, Loughborough LE11 3TU, United Kingdom and Institute of Terrestrial Magnetism, Ionosphere and Radio Wave Propagation, Russian Academy of Sciences, Troitsk, Moscow Region 142190, Russia

(Received 11 February 2006; accepted 22 February 2006; published online 11 May 2006)

[DOI: [10.1063/1.2186766](https://doi.org/10.1063/1.2186766)]

1. Formula (23) on p. 5 should be

$$m = 1: \quad V_2 = V_3 = \frac{1}{3}(\beta_1 + 2\beta_3), \quad V_1 = \beta_1.$$

2. On p. 13, where the problem of the decay of a step for the defocusing modified KdV [mKdV(d)] equation is considered, after formula (90), in the case (i), a wrong expression for the velocity/amplitude relationship for the mKdV(d) soliton was used. This led to an incorrect expression for the lead soliton amplitude in the mKdV(d) dispersive shock. The correct expression for the amplitude is derived and discussed below.

For the initial step (88) satisfying the condition of the case (i), $u_+^2 - u_-^2 > 0$, the speeds of the mKdV(d) dispersive shock edges $s^\pm(u_+, u_-)$ are determined by expressions (89). Then the amplitude of the lead soliton a^+ is found from the kinematic condition $c_s(u^+, a^+) = s^+$ [cf. (62); here we use a different set of independent variables for the function c_s], where the relationship between the velocity and the amplitude for the mKdV(d) soliton moving against the background \bar{u} is obtained by an elementary analysis of the traveling wave solution in the soliton limit and has the form $c_s(\bar{u}, a_s) = -\bar{u}^2 \pm 2\bar{u}a_s/3 - a_s^2/6$. Different signs in this expression correspond to solitons of different polarity supported by the mKdV(d) equation. Now, after simple calculation one gets

$a^+ = 2(|u_+| - |u_-|)$. For $u_+u_- \geq 0$ this coincides with the KdV result (31) [to make such a comparison in the case when $u^+ > u^- \geq 0$, the KdV equation should be taken with the negative sign for the nonlinear term, which would result in the change $\Delta \rightarrow -\Delta$ in (31)]. The location of the mKdV(d) dispersive shock given by (89), however, is different from the respective KdV case; for the KdV equation with the same initial data one has $|s^+ - s^-| = \frac{5}{3}(|u_+| - |u_-|)t$ rather than $\frac{5}{3}(u_+^2 - u_-^2)t$.

Of course, these results could have been inferred from the already known periodic solution and the Riemann form of the modulation system for the mKdV(d) equation (see Ref. 1, for instance) but here they were obtained without invoking an integrable structure of the mKdV(d) equation, as an illustration of the effectiveness of the general transition relations for a dispersive shock.

The author is grateful to Tim Marchant for pointing out the error in the expression for the lead soliton amplitude in the mKdV dispersive shock.

¹A. M. Kamchatnov, A. Spire, and V. V. Konotop, “Dissipationless shock wave in a discrete nonlinear Schrödinger equation,” *J. Phys. A* **37**, 5547–5568 (2004).