

Northumbria Research Link

Citation: Bongiovanni, Tindaro, Genovesi, Federico, Nemmer, Monika, Carling, Christopher, Alberti, Giampietro and Howatson, Glyn (2020) Nutritional interventions for reducing the signs and symptoms of exercise-induced muscle damage and accelerate recovery in athletes: current knowledge, practical application and future perspectives. *European Journal of Applied Physiology*, 120 (9). pp. 1965-1996. ISSN 1439-6319

Published by: Springer

URL: <https://doi.org/10.1007/s00421-020-04432-3> <<https://doi.org/10.1007/s00421-020-04432-3>>

This version was downloaded from Northumbria Research Link: <http://nrl.northumbria.ac.uk/id/eprint/43685/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

1 **Nutritional interventions for reducing the signs and symptoms of exercise-induced muscle damage and accelerate**
2 **recovery in athletes: current knowledge, practical application and future perspectives**

3

4 Tindaro Bongiovanni^{1*}, Federico Genovesi², Monika Nemmer³, Christopher Carling⁴, Giampietro Alberti⁵ and Glyn
5 Howatson^{6,7}

6

7

8 ¹Department of Health, Performance and Recovery, Parma Calcio 1913, Parma, Italy

9 ²Medical Department Manchester City Football Club, Manchester, UK

10 ³Nutrition Department Liverpool Football Club, Liverpool, UK

11 ⁴Centre for Elite Performance, French Football Federation, Paris, 75015, France

12 ⁵Department of Biomedical Sciences for Health, Università degli Studi di Milano, Milano, Italy

13 ⁶Department of Sport, Exercise and Rehabilitation, Northumbria University, Newcastle upon Tyne, UK.

14 ⁷Water Research Group, North West University, Potchefstroom, South Africa.

15

16

17

18

19

20

21

22

23

24 *Corresponding author:

25 Tindaro Bongiovanni

26 Department of Health, Performance and Recovery, Parma Calcio 1913, Parma, Italy

27 tindaro.bongiovanni@gmail.com

28

29

30

31 **Keywords**

32 Damage, muscle, supplementation, nutrition, recovery

33

34	Abbreviations	
35	EIMD	Exercise-induced muscle damage
36	ROM	Range of motion
37	CK	Creatine kinase
38	LDH	Lactate dehydrogenase
39	MYO	Myoglobin
40	CRP	C-reactive protein
41	ACTN3	Gene that encodes the α -actinin-3 protein
42	TNF	Tumour necrosis factor
43	IL-6	Interleukin-6
44	IGF2	Insulin-like growth factor 2
45	ECM	Extracellular matrix
46	RBE	Repeated bout effect
47	E-C	Excitation contraction
48	NGF	Nerve growth factor
49	GDNF	Glial cell line-derived neurotrophic factor
50	PLA ₂	Phospholipase A ₂
51	FFA	Free fatty acids
52	EPA	Eicosapentaenoic acid
53	DHA	Docosahexaenoic acid
54	CMJ	Countermovement jump
55	RSI	Reactive strength index
56	BCAA	Branched chain amino acids
57	CMJ	Counter movement jump
58	AST	Aspartate aminotransferase
59	ALT	Alanine aminotransferase
60	1-RM	One-repetition-maximum
61	MDA	Malondialdehyde
62	PC	Protein carbonyls
63	MSM	Methylsulfonylmethane
64	RONS	Reactive oxygen and nitrogen species
65	MVC	Maximal voluntary contraction
66	PNS	Polar-nonpolar-sandwich
67	VAS	Visual analogic scale
68	TBARS	Thiobarbituric acid reactive substances
69		
70		
71		
72		
73		

74 **Abstract**

75 **Purpose** This review provides an overview of the current knowledge of the nutritional strategies to treat the signs and
76 symptoms related to EIMD. These strategies have been organized into the following sections based upon the quality and
77 quantity of the scientific support available: 1) interventions with a good level of evidence; 2) interventions with some
78 evidence and require more research; and 3) potential nutritional interventions with little to-no-evidence to support
79 efficacy.

80 **Method** Pubmed, EMBASE, Scopus and Web of Science were used. The search terms ‘EIMD’ and ‘exercise-induced
81 muscle damage’ were individually concatenated with ‘supplementation’, ‘athletes’, ‘recovery’, ‘adaptation’, ‘nutritional
82 strategies’, ‘hormesis’.

83 **Result** Supplementation with tart cherries, beetroot, pomegranate, creatine monohydrate and vitamin D appear to provide
84 a prophylactic effect in reducing EIMD. β -hydroxy β -methylbutyrate, and the ingestion of protein, BCAA and milk could
85 represent promising strategies to manage EIMD. Other nutritional interventions were identified but offered limited effect
86 in the treatment of EIMD; however, inconsistencies in the dose and frequency of interventions might account for the lack
87 of consensus regarding their efficacy.

88 **Conclusion** There are clearly varying levels of evidence and practitioners should be mindful to refer to this evidence-
89 base when prescribing to clients and athletes. One concern is the potential for these interventions to interfere with the
90 exercise-recovery-adaptation continuum. Whilst there is no evidence that these interventions will blunt adaptation, it
91 seems pragmatic to use a periodised approach to administering these strategies until data are in place to provide and
92 evidence base on any interference effect on adaptation.

93

94

95 **Introduction**

96 Exercise-induced muscle damage (EIMD) is a transient event caused by unaccustomed exercise (Harty et al.
97 2019) or strenuous high intensity or long duration exercise (Jang et al. 2018; Leeder et al. 2014). Moreover, EIMD can
98 be evident after isometric contractions at a long muscle length, even at a low intensity (Chen et al. 2012). Muscle damage
99 is characterized by a primary response from a mechanical challenge during the exercise that results in myofibrillar
100 disruption (Yu et al. 2013) and a secondary inflammatory response that results in cell infiltration into the damaged tissues
101 (Deyhle et al. 2015; Harty et al. 2019) to initiate subsequent tissue repair and remodelling (Peake et al. 2017). Beyond
102 myofibre damage, the extracellular matrix (ECM) of skeletal muscle is also subjected to degradation and remodelling
103 following muscle-damaging exercise that can be directly observed histologically (Hyldahl et al., 2015). Indirect evidence
104 of ECM damage also exists, with a marked increase in the circulation of collagen-specific amino acids, most notably
105 hydroxyproline, in the days following muscle-damaging exercise (Clifford et al. 2019).

106 Although the exact mechanisms responsible for skeletal muscle damage remain unclear, a damage model has
107 been proposed (Figure 1). The damage model can be briefly summarized into two general phases: an initial or *primary*
108 *phase*, which occurs during the exercise bout (Armstrong 1984; Ebbeling and Clarkson 1989; Belcastro et al. 1998;
109 Verburg et al. 2005; Belcastro 1993) that result in an increase in cell membrane permeability causing further influx of
110 Ca^{2+} (for more details see Gissel 2005). The *secondary phase* which is linked with the inflammatory response and
111 increased oxidative stress that further affect cell structure and function (Jackson et al. 1984; Howatson and van Someren
112 2008; Baumert et al. 2016; Owens et al. 2019; Hody et al. 2019). Thus, muscle damage can be considered a tightly
113 regulated process of degradation and repair, where several cell types (i.e. neutrophils, macrophages, cytokines) interact
114 during the proinflammatory and inflammatory stages of muscle damage with the aim of restoring tissue homeostasis.

115 Most of the signs and symptoms of EIMD (i.e. muscle soreness, stiffness, tenderness, pain, impaired muscle
116 function) are evident within a few hours following the exercise insult and can persist for many days following the initial
117 bout, however the magnitude and duration can vary greatly between individuals. Of primary concern is a temporary
118 decrease in muscle function (strength, power, speed, economy of movement) that is accompanied by reduced range of
119 motion (Jamurtas et al. 2005; Howatson et al. 2008; Damas et al. 2016; Qamar et al. 2019). Furthermore, there is often
120 evidence of muscle swelling, muscle soreness, decreased pain threshold, increase of oxidative stress markers (Childs et
121 al., 2001; Lee et al., 2002; Sacheck et al., 2003) and rise of various inflammatory interleukins, C-reactive protein (CRP),
122 tenascin C, which can be accompanied by elevated intramuscular proteins (creatine kinase; CK, myoglobin; MYO, lactate
123 dehydrogenase; LDH, for example); these can be present for several days following the damaging bout of exercise (Byrne
124 et al. 2004; Hyldahl et al. 2014; Warren et al. 1999; Harty et al. 2019; Qamar et al. 2019; Baumert et al. 2016). Delayed-
125 onset of muscle soreness (DOMS), a common symptom of EIMD, typically appears between 8 and 24 hours after the
126 muscle-damaging exercise, peaks between 24 and 48 hours and usually subsides within 96 h (Damas et al. 2016; Jones et
127 al. 1987). The magnitude and time course of signs and symptoms of EIMD and their subsequent impact on performance
128 are variable and depend on the extent of initial muscle damage, which in turn is influenced by the duration, muscle group
129 targeted, mode of exercise, intensity, and volume of the damaging exercise (Bowtell et al. 2011; Peake et al. 2017) and
130 the individual's susceptibility to the damaging stimulus (Owens et al. 2019). Age, sex and physiological training status
131 of the individual (Plowman and Smith 2013), along with genetic variability where certain genotypes are thought to be
132 more susceptible to damage (Baumert et al. 2016), are additional factors that could play a contributing factors in the
133 magnitude to EIMD and hence might require a longer recovery period.

134 Research indicates that inflammation and oxidative stress is evident following muscle damaging exercise
135 (Jamurtas 2018). However, it is widely accepted these processes are necessary for the adaptive remodelling process and

136 hence critical for the resolution of function (Owens et al. 2019). Consequently, there is a dilemma when planning nutrition
137 strategies for interventions that target the post-exercise inflammatory and oxidative stress response. If the principal
138 purpose is to maximize the training stimulus, athletes and practitioners might need to consider a periodised approach to
139 supplementation to adequately support training and competition demands to maximize the balance between the potential
140 for adaptation and optimising recovery. This notion can be applied with the hormesis concept, whereby the biological
141 systems respond in a bell-shaped fashion, where a positive adaptive response is experienced when exposed to an increase
142 in training stimulus (Hawley et al. 2011). However, when the stimulus exposure becomes too great; for example, when
143 EIMD impairs muscular functions for an extended period or the need to accelerate recovery is greater than the need for
144 an adaptive response, a need to intervene to negate the negative effects exists. Consequently, the aim of a recovery
145 intervention is highly dependent upon the purpose of the session and whether it is more important adapt or accelerate
146 recovery of muscle function, ameliorate soreness in preparation for subsequent training or competition (Figure 2).

147 Optimal nutrition plays an important role during post-exercise recovery, where replenishment of endogenous
148 substrate stores and facilitation of muscle-damage-repair process are the main goals (Beelen et al., 2010). The
149 fundamentals of nutrition are extremely important, so ensuring adequate hydration, carbohydrates and protein availability
150 are critical, but these fundamental aspects of nutrition are not covered here. Thus, the aim of this review is to provide an
151 overview of the current evidence of nutritional supplementation strategies to ameliorate the signs and symptoms related
152 to EIMD beyond the fundamentals of good hydration and macronutrient intake. Specifically, we focus on nutritional
153 strategies that have been explored to accelerate the recovery process after damaging exercise. These strategies have been
154 organized into the following sections based on level of available evidence: 1) interventions with good level of research
155 investigating their efficacy; 2) interventions that have a modest evidence base; and 3) nutritional interventions with little
156 or no evidence that might prove worthwhile avenues for future research.

157
158

159 **Supplements with good level of evidence**

160

161 **Vitamin D**

162 It is widely recognized that vitamin D plays an important role of bone health of athletes (Larson-Meyer et al.
163 2019), maintaining a healthy mineralized skeleton (Charoenngam et al. 2019) and regulating the processes of skeletal
164 muscle regeneration (Owens et al. 2015). Vitamin D also has immunomodulatory effects, with increased vitamin D shown
165 to reduce inflammation (Willis et al. 2008); furthermore, there is a strong association between sufficient vitamin D and
166 optimal muscle function (Shuler et al. 2012). Vitamin D₃ (cholecalciferol) can be synthesized endogenously in the skin
167 as a result of exposure to ultraviolet B radiation from sunlight, or obtained from dietary intake in foods that include fatty
168 fish, eggs, and fortified foods (i.e. milk, citrus juice, breakfast cereals), while vitamin D₂ (ergocalciferol) is found only in
169 plants and fungi such as mushrooms (Wilson et al. 2017) and algae (Göring 2018). Dietary sources of vitamin D are
170 thought to have only a small effect on serum 25(OH)D concentration (an index of vitamin D levels) and the combination
171 of limited dietary sources and several others factors (i.e. season, sport, body composition, training location, altitude, air
172 pollution, geographical latitude, athlete's skin pigmentation) could be responsible for the increased risk of vitamin D
173 deficiency in athletes (Larson-Meyer et al. 2019).

174 A previous randomised, placebo-controlled trial has shown the role of vitamin D for muscle recovery, and the
175 interplay between serum vitamin D status and the magnitude of exercise-induced muscle damage (Owens et al., 2015) in
176 20 moderately active, adult males with deficient vitamin D (serum 25(OH)D). Following a six week oral vitamin D₃
177 supplement (4000 IU/day) or a placebo, volunteers completed a damaging bout of exercise. Serum 25(OH)D was

178 increased in the vitamin D₃ group and had improved recovery of muscle function in the days following the damaging
179 exercise compared to the placebo control. Conversely, Shanely et al. (2014) found no effects of vitamin D₂
180 supplementation (600 IU per day, for 6 weeks) on recovery of lower limb muscle function or circulating markers of
181 muscle damage following 1.5 h high-intensity intermittent exercise. The association between vitamin D status, exercise-
182 induced muscle pain and weakness was investigated in 48 young healthy males and females (Ring et al. 2010). Following
183 a damaging bout of eccentric exercise to the elbow flexors (3 x 12, 75% MVC) vitamin D status was not able to predict
184 changes in muscle strength loss or pain outcomes. In contrast, an observational study reported that pre-exercise vitamin
185 D status was influential on muscular weakness after eccentric exercise (Barker et al. 2013a). The study showed a transient
186 increase in circulatory 25(OH)D immediately after the exercise, followed by a decrease. The authors suggested that
187 serum 25(OH)D is temporally sensitive to intense exercise and that a multitude of biomarkers (i.e. cytokines, ALT, AST,
188 albumin) contribute, in part, to the 25(OH)D fluctuations (Barker et al. 2013a) and hence provide some evidence of the
189 interplay between eccentric exercise, circulating vitamin D and the possible role in recovery.

190 Following exercise-induced muscle damage (EIMD), induced by eccentric exercise, Barker et al. (2013b)
191 demonstrated that supplemental vitamin D₃ (4000 IU/d) for 35 days increased serum 25(OH)D concentrations and was
192 associated with a faster restoration of peak force during recovery. Peak force was approximately 8% superior in the
193 supplemented group compared to the placebo at 24 h after the damaging exercise, although recovery between the groups
194 was similar thereafter. Other biomarkers of exercise stress, such as circulating aspartate amino-transferase (68 h post-
195 exercise) was attenuated by 29%, while alanine amino-transferase (24 h post-exercise) was attenuated by 40%, compared
196 to placebo. In contrast, Nieman et al. (2013) showed that vitamin D₂ supplementation in professional race car pit-crew
197 had no effect on muscle function tests following 90 min eccentric exercise.

198 Potential reasons for these discrepancies across research might be related to outcome measures, populations,
199 mode/type, length and dose of supplementation and follow-up or different experimental protocols, including the muscle
200 group(s) exercised, damaged protocol, contraction modalities, volume, duration, intensity of the exercise protocol. In
201 addition, differences between studies exist with respect to baseline levels of 25(OH)D. More work is necessary to clarify
202 the benefit of vitamin D for athletic muscle recovery. Athletes should aim to maintain appropriate vitamin D status
203 through regular sun exposure, supplementation, and/or diet. The target level of 25(OH)D has not been identified for an
204 endpoint of muscle repair; however, it seems prudent for athletes to aim for the clinical cut-off for sufficient vitamin D
205 status, which is 30-50 nmol/L (Owens et al., 2015). The available data suggest vitamin D might play a role in the muscle
206 repair and recovery process (Heaton et al. 2017) but based on the available evidence there is no clear benefit in improving
207 recovery following damaging exercise.

208

209 **Omega-3 polyunsaturated fatty acids**

210 Omega-3 fatty acids (eicosapentaenoic acid or EPA and docosahexaenoic acid or DHA) are essential nutrients;
211 there is no mechanism in humans for producing these fats however, they can be consumed through oily fish, selected
212 vegetables, nuts and seeds. It is widely known that omega-3 are effective for improving cardiac function, depression,
213 cognitive function and a relationship with improved endurance performance (Ochi and Tsuchiya 2018). Omega-3
214 polyunsaturated fatty acids (PUFA) are able to modulate inflammation, decrease swelling, and reduce sensitivity to pain
215 (Jouris et al. 2011). Moreover, EPA and DHA have been shown to play a role in the regulation of skeletal muscle protein
216 synthesis and immune function (Heaton et al. 2017). Many studies have highlighted that intake of omega-3 PUFA could
217 modulate EIMD outcomes, though results are generally equivocal because of high variability in study designs,
218 participants, dosing and timing protocol exists in the literature (Tartibian et al. 2009; Philpott et al. 2018; Tsuchiya et al.,

219 2016). For example, some studies have reported attenuated the increase in plasma IL-6 up to 72 h after strenuous exercise
220 (DiLorenzo et al. 2014; Phillips et al. 2003; Tartibian et al. 2011) and others have not (Lenn et al. 2002; Nieman et al.
221 2009).

222 Lenn et al. (2002) examined the effect of 30 days fish oils (1.8 g of EPA and DHA per day) supplementation on
223 delayed onset of muscle soreness following a damaging bout of exercise. The authors reported that fish oils were not
224 effective in ameliorating DOMS in comparison to a placebo or isoflavones (120 mg of soy isolate) supplementation.
225 Similarly, others (Bloomer et al., 2009) found no effect of 6 weeks of omega-3 PUFA (2224 mg EPA and 2208 mg DHA
226 per day) on attenuation of oxidative stress and inflammation induced by a 60-min treadmill climb, in 15 well-trained
227 participants. In contrast, Tartibian et al. (2009) showed a beneficial treatment effect of omega-3 (32 days of 1.8 g of EPA
228 and DHA per day versus a placebo) on perceived pain and a minor decrement of range of motion 48 h after damaging
229 exercise (40 min of bench stepping exercise) in 27 healthy untrained males. In a subsequent randomized, double-blinded
230 study (Tartibian et al., 2011), 45 healthy untrained men completed an eccentric exercise program (40 minutes of bench
231 stepping) following 30 days supplementation (324 mg of EPA and 216 mg of DHA per day, which continued 48 h after
232 exercise. The authors showed a reduction in the plasma TNF- α , PGE₂, LDH, IL-6, CK immediately, 24, and 48 h after
233 the damaging exercise compared to the placebo (soybean/corn oil mixture) and control group. A recent double-blind,
234 placebo-controlled, parallel design study showed that daily supplementation for 8 weeks of fish oil rich in omega-3 (EPA;
235 600mg and DHA; 260 mg) has a positive role in inhibiting muscle stiffness after eccentric contractions and was able to
236 inhibit the loss of muscle strength, reduce ROM, development of DOMS, and to reduce muscle swelling in comparison
237 to a placebo (Tsuchiya et al. 2019). Likewise, Ramos-Campo et al. (2020) reported that the ingestion of 2.1 g of DHA
238 and 240 mg of EPA every day for 10 weeks, promoted lower concentrations of IL1 β , IL6, CK, LDH and muscle soreness
239 after a damaging eccentric exercise in endurance athletes, when compared to placebo treatment (olive oil).

240 In eleven healthy men and women, Jouris et al. (2011) tested the effects of 7 days of EPA and DHA
241 supplementation, in a ratio 2:1 (2000 mg of EPA and 1000 mg of DHA) on inflammation induced by two sets of eccentric
242 elbow flexion exercise (120% 1RM) and showed a reduction of post-exercise soreness at 48 h post exercise. In contrast,
243 Houghton and Onambele (2012) investigated the effects of a dose of 360 mg of EPA supplementation per day for 3 weeks
244 on muscle soreness and basal inflammation induced by acute and chronic resistance exercise in a three-week, double-
245 blind placebo-controlled study in 20 participants. Data showed no effect in reducing DOMS and attenuating increases in
246 IL-6 levels in comparison to a placebo. The authors speculated that these results might have been associated by increased
247 EPA-mediated contractile capacity, resulting in increased glucose metabolism, glycogen depletion and hence elevations
248 in the IL-6 concentrations. Conversely, in a study by DiLorenzo et al. (2014), 41 untrained volunteers received 28 days
249 (2 g/day) of DHA or a placebo, before a 17-day resistance exercise program. The DHA supplemented group showed a
250 reduction of CK and IL-6 response in the first 4 days of the program, while no differences in muscle strength and soreness
251 were observed between groups. Similarly, Gray et al. (2014) examined the effect of a six-week fish oil supplementation
252 (1.3 g EPA, 0.3 g DHA and 45 I.U. d- α tocopherol) or a placebo on exercise induced muscle soreness and oxidative stress
253 markers in 20 trained males following eccentric exercise. There were no differences between groups for indices of muscle
254 damage and soreness. However, the fish oil group showed lower oxidative stress markers (thiobarbituric acid reactive
255 substances; TBARS and H₂O₂ stimulated DNA damage) at 48 and 72 h after the exercise bout of eccentric exercise.
256 Nevertheless, it is important to treat these results with caution, because TBARS and H₂O₂ have methodological limitations
257 and should be considered surrogate measures, at best. In another study (Lembke et al. 2014), 63 healthy male and female
258 college students were supplemented for 30 days with omega-3 (2.7 g per day) or placebo (sunflower oil) prior to forearm
259 extension eccentric exercise. Results showed a decrease in muscle soreness at 72 h and 96 h after damaging exercise in

260 the supplemented group compared to placebo. A more recent double-blind, placebo-controlled, parallel-group trial
261 (Tsuchiya et al. 2016), conducted on 24 healthy Japanese men, concluded that EPA and DHA supplementation (600 mg
262 EPA and 260 mg DHA per day) for eight weeks prior to, and 5 days after maximal unilateral isokinetic elbow flexor
263 exercise was more beneficial than a placebo to attenuate strength loss and limit reductions in ROM during the recovery
264 period. Jakerman et al. (2017) first examined the effect of an acute intake of two fish oil supplements, with different
265 concentrations of EPA and DHA in a double-blind, placebo-controlled study conducted in 27 physically active males.
266 Immediately following a damaging bout of exercise (100 drop jumps), volunteers ingested either a low-EPA fish oil (150
267 mg EPA, 100 mg DHA), high-EPA fish oil (750 mg EPA, 50 mg DHA) or a placebo. Results showed that acute ingestion
268 of high-EPA fish oil was beneficial on some markers of performance compared to low-EPA fish oil and placebo.

269 Some have reported beneficial effects of omega-3 supplementation in females. In 42 healthy women, Corder et
270 al. (2016) examined the effect of DHA supplementation (7 days before and 2 days after exercise with 3000 mg per day)
271 on DOMS and local muscle inflammation after an eccentric elbow flexion exercise. Results showed DHA
272 supplementation reduced muscle soreness and stiffness after 48 h, but no effects on swelling compared to a placebo.
273 Later, Tinsley et al. (2017) supplemented both EPA and DHA in 17 non-resistance trained females. Participants were
274 randomized into fish oil (6 g/d; 5:1 EPA:DHA) or placebo groups and took the supplement for one week prior to, and one
275 week after a damaging elbow flexion and extension exercise. Results showed that at 48 h post-exercise, the supplemented
276 group reported less muscle soreness in the lower body than the placebo group.

277 In some recent investigations the benefits of combining omega-3 with other nutrients have been
278 examined. For example, Philpott et al. (2018) added 1100 mg EPA and 1100 mg DHA to a ready-to-drink beverage (15
279 g whey protein, 1.8 g leucine and 20 g carbohydrate) and supplemented this blend in ten male soccer players each day for
280 6 weeks. The other two groups (10 participants per group) ingested a beverage containing 15 g whey protein, 1.8 g leucine
281 and 20 g carbohydrate, or a beverage containing only carbohydrate (20g). After 6 weeks of supplementation, an eccentric
282 damaging exercise was performed, and muscle soreness and blood markers of muscle damage were measured. Notable
283 results showed that muscle soreness, during the 72 h recovery, and blood CK was less in participants that ingested the
284 beverage with EPA and DHA, compared to other two groups. No differences in muscle function, soccer performance or
285 blood CRP were observed between groups. In a further recent double-blind study, Black et al. (2018) investigated the
286 effectiveness of consuming, twice daily, 551 mg EPA and 551 mg DHA added to a protein-carbohydrate supplement (15
287 g protein, 14.5 g carbohydrate and 8.4 g fat) or placebo (15 g protein, 14.5 g carbohydrate and 8.4 g fat, without omega-
288 3) in 33 professional Rugby Union players during 5 weeks of pre-season training. Reduced muscle soreness and fatigue
289 and a better maintenance of jump performance was reported in athletes that consumed the enriched-omega 3 beverage
290 compared to placebo group during all time points. The efficacy of a dietary supplement that contained, in part, omega 3
291 PUFA to modulate inflammation following EIMD was evaluated in 40 healthy, untrained males (Phillips et al., 2014).
292 Researchers supplemented a blend consisting of 300 mg mixed tocopherols, 800 mg DHA and 300 mg of flavonoids (100
293 mg hesperetin and 200 mg quercetin) or placebo (high-oleic sunflower oil) for 14 days. Although blood indices of damage
294 (LDH and CK) was not different between groups, the supplemented group showed a reduction in IL-6 and CRP after
295 eccentric exercise, compared to the placebo. Although these show some promise, the blended supplement makes it
296 difficult to attribute solely to omega-3 PUFA.

297 The divergent results in this section could be explained, in part, by differences in supplement dose, type of fatty
298 acid ingested, fitness level of participants (trained vs untrained) duration of supplementation and intensity and/or modality
299 of exercise. Albeit there is a potential for omega-3 PUFA supplementation to improve muscle adaptation and muscle
300 recovery, supplementation might yet prove to be effective to attenuate muscle damage (Hennigar et al. 2017) and further

301 study in to the optimal dose of EPA:DHA ratio, the synergistic effect with other nutrients and the timing areas for further
302 enquiry.

303

304 **Tart Montmorency cherries (*Prunus cerasus*)**

305 Montmorency tart cherries (*Prunus cerasus L.*) have been proposed as a recovery supplement because of their
306 high concentrations of phytochemicals, and in particular, the flavonoids anthocyanins and other phenolic compounds
307 (Bell et al. 2015; Coelho et al. 2015; Howatson et al. 2010). These phenolic compounds have been shown to reduce
308 oxidative stress and inflammation by inhibiting the action of cyclooxygenase to a similar extent as nonsteroidal anti-
309 inflammatory drugs (Seeram et al. 2001; Wang et al., 1999). These findings have led to numerous lines of enquiry that
310 applied Montmorency cherries in exercise recovery (Connolly et al., 2006; Howatson et al., 2010; Kuehl et al., 2010;
311 Bowtell et al. 2011; Bell et al., 2014).

312 The first study (Connolly et al., 2006) examined the efficacy of a tart Montmorency cherry juice blend in
313 preventing the symptoms of exercise induced muscle damage in a randomised, placebo-controlled trial. Results showed
314 that 12 oz of tart Montmorency cherry juice taken twice a day for 4 d before and 4 d after heavy eccentric contractions of
315 the elbow flexors, reduced pain and limited post-exercise strength loss (4% versus 22%) compared to a placebo. In a
316 subsequent applied study, Howatson et al. (2010) use the same cherry juice blend used previously (Connolly et al., 2006)
317 and showed improved recovery of strength and reduced inflammation (IL-6 and C-reactive protein) and oxidative stress
318 (TBARS) following a marathon run. This placebo controlled, parallel group study supplemented with 8 fl oz of cherry
319 juice twice a day, ingested for 5 consecutively days before, the day of race and 2 days after running a marathon or a
320 placebo group (8 fl oz of fruit flavoured cordial). A further study using a randomized, double blind, placebo-controlled
321 trial design (Kuehl et al., 2010) in a cohort of 54 runners, examined the effect of tart Montmorency cherry juice following
322 a strenuous trail running event. They showed that ingestion of 335 mL tart Montmorency cherry juice consumed for 7
323 days prior to, and during, the event reduced post-run muscle pain compared to placebo. Collectively, these studies indicate
324 that the intervention is effective to reduce pain and restore function in damaging running activity.

325 Bowtell et al. (2011) showed that consumption of a tart Montmorency cherry concentrate (30 mL twice per day
326 for 10 days) in ten well-trained males reduced symptoms of EIMD. Force recovery was greater in the cherry juice group
327 after a damaging eccentric knee extensor exercise compared to an isoenergetic fruit concentrate placebo. This study
328 illustrated the potential application in a well-trained cohort following strength-based training. In a further study (Kastello
329 et al. 2014) using a double-blind, placebo controlled, cross-over design, the effects of tart cherry supplementation (freeze
330 dried powder tablets – one tablet, twice per day for 16 days prior to, and 3 days following) was examined in an untrained
331 mixed sex cohort (10 females and 4 males) following damaging exercise. Findings suggested showed cherries to reduce
332 C reactive protein levels, range of motion and perceived pain more than a placebo (Kastello et al. 2014).

333 The aforementioned studies show that tart Montmorency cherries are able to provide enhanced recovery
334 following exercise with a mechanical stressor where there is myofibrillar disruption from high intensity eccentric
335 contractions. In an attempt to investigate the possible application in a solely metabolic challenge, Bell et al. (2015)
336 investigated if a tart Montmorency cherries concentrate would facilitate recovery on signs and symptoms of EIMD in 16
337 well-trained cyclists. In this parallel group design, volunteers were given the cherry concentrate or caloric matched
338 placebo for 8 days. On the fifth day of supplementation, a strenuous bout of cycling designed to replicate a road race was
339 completed and indices of inflammation, and muscle damage were taken. Results showed that the Montmorency tart cherry
340 group showed lower levels of IL-6 and hsCRP, and attenuated reductions in muscle function (Bell et al. 2015). In a
341 similar study with the same research group, Bell et al. (2014) examined the effect of a tart Montmorency cherry

342 concentrate in trained cyclists conducting three repeated days of simulated road racing and showed reductions in
343 inflammation during the 3 days of cycling compared to a placebo. Collectively, these data show that tart Montmorency
344 cherries can support recovery following both mechanically and metabolically challenging exercise (Bell et al., 2014).

345 Levers et al. (2015) investigated 10 days supplementation (480 g/d) with tart cherry skin powder on indices on
346 EIMD. After 7 days, of supplementation, resistance trained males conducted a damaging bout of resistance training
347 followed by 3 days supplementation for three days. They showed reduced creatinine, total protein and perceived soreness
348 in comparison to a placebo in the 48 h after the exercise insult. In a successive study (Levers et al. 2016), using the same
349 powdered tart cherries, examined in 27 endurance-trained runners or triathletes (18 males and 9 females) following a half
350 marathon. The groups received a similar 7 day loading phase and dose (480 g/d) followed by two days of supplementation
351 or a placebo. The dried cherry skins attenuated post-run muscle soreness, urea/blood urea nitrogen, total protein, and
352 cortisol compared to the placebo group (Levers et al. 2016). In a related study to Howatson et al. (2010), Dimitriou et al.
353 (2015) investigated the effects of Montmorency cherry juice on muscle inflammation, immunity and stress. Using the
354 same cohort and supplementation to the aforementioned study (Howatson et al., 2010), the cherry juice group showed no
355 upper respiratory symptoms in the period following the marathon compared to the placebo group indicating some potential
356 for immune function.

357 Montmorency tart cherry supplementation was also studied in sixteen semi-professional male soccer players in
358 a double blind, placebo-controlled design with independent groups following prolonged intermittent sprint exercise (Bell
359 et al. 2016). Participants were supplemented with 30 mL twice per day of tart Montmorency cherry concentrate or caloric
360 matched placebo for seven consecutive days. This study showed an attenuation in performance declines and a reduction
361 of muscle damage indices, for example IL-6 at 24 and 48 h post exercise. In addition, Brown et al. (2019) investigated
362 the effect of tart cherry in twenty female dancers in a randomised, double-blind, placebo-controlled study. Participants
363 were allocated tart Montmorency cherry concentrate (30 mL twice a day) or placebo (fruit flavoured concentrate) for
364 eight days. After 4 days of supplementation, participants completed a damaging repeated sprint exercise protocol. Results
365 showed that dancers that ingested tart cherry juice had a faster recovery of muscle function, as well as a lower reported
366 DOMS compared to the placebo group. Likewise, Quinlan and Hill (2019) reported that male and female team-sport
367 players who ingested tart Montmorency cherry juice concentrate accelerated recovery after an intermittent shuttle run
368 test, compared to the placebo group. Collectively, these data show the application of tart Montmorency cherries in
369 repeated sprint activity like soccer and other field sports for both males and females.

370 Tart cherry was investigated in different doses, formulation and different sports and trained and untrained
371 athletes. Overall, studies have typically reported beneficial effects on signs and symptoms of EIMD, however, some work
372 has shown no benefit (Beals et al. 2017; McCormick et al. 2016). Beals et al. (2017) saw no beneficial effect of
373 supplementation with a beverage containing freeze-dried tart cherry powder on the recovery of DOMS or muscle strength
374 in recreationally active men and women following damaging eccentric exercise. Similarly, McCormick et al. (2016) found
375 no effect of supplementation with tart cherry juice on DOMS and markers of inflammation and oxidative stress in well-
376 trained male water polo players who undertook a simulated game of water polo. Furthermore, in a recent study (Lamb et
377 al. 2019) the supplementation with tart cherry juice (2 x 250 mL) for 9 days did not enhance recovery in non-resistance
378 trained men after maximal eccentric elbow flexor exercise. Likewise, Abbott et al. (2019) have reported that academy
379 soccer players who ingested tart cherry juice at the dose of 30 mL before and 30 mL after a 90-minute match and at 12
380 and 36 hours after failed to hasten recovery of muscle function, when compared who ingested an isocaloric cherry-
381 flavoured control drink, although this study was not able to control diet and did not include a loading phase which might
382 explain the null effects. In many of the cases, tart Montmorency cherries reduce the inflammatory response following

383 exercise, which is an important signally cascade for the adaptative response. Whilst there is no evidence that any
384 functional foods affect the adaptive response, it is clear that some inflammatory processes could be modulated. One
385 caveat to this is that the inflammatory responses are not absent, but rather reduced by the intervention, so some
386 inflammatory signalling is still present; the question for researchers and practitioners is “how much inflammation is
387 enough?”. Pragmatically, once the athletes are in a condition where further adaptations are not a priority, and restoration
388 of function is the focus, tart Montmorency cherries have a beneficial role (Vitale et al. 2017).

389

390 **Creatine monohydrate**

391 Beyond the well-known effects of creatine monohydrate to improve exercise performance such as power (Kim
392 et al. 2015), body composition, anaerobic, aerobic and strength performance (Fernandez-Landa et al. 2019), several
393 studies suggested that creatine supplementation might also reduce muscle damage and/or enhance recovery following
394 intense exercise (Kreider et al. 2017; Kim et al. 2015). Creatine supplementation has been shown to play an important
395 role in muscle damage; one mechanism could be through cell membrane stabilization. The molecular structure of
396 phosphocreatine, allows it to bind to the phospholipid heads of the cell membrane and stabilize the membrane, thereby
397 reducing membrane fluidity and thus potentially decrease the degradation associated with muscle damage (McKinnon et
398 al. 2012). Vierck et al. (2003) and Olsen et al. (2006) demonstrated that creatine supplementation might influence post-
399 exercise muscle recovery by enhancing muscle satellite cell proliferation. Since satellite cells are myogenic stem cells
400 that fuse with damaged muscle fibres to affect repair (Olsen et al. 2006), increased satellite cell proliferation could result
401 in an increase recovery in muscle fibre integrity (McKinnon et al. 2012). Other potential mechanisms provide some
402 rationale for the use of creatine on EIMD, including modulation of the inflammatory response (Santos et al. 2004; Bassit
403 et al. 2008; Deminice et al. 2013), oxidative stress (Lawler et al. 2002; Deminice and Jordao 2012) and regulation of
404 calcium homeostasis (Beaton et al., 2002; Cooke et al., 2009). According to a human study by Rahimi (2011) ingesting
405 20 g/day of creatine for 7 days decreased oxidative stress (malondialdehyde; MDA and 8-hydroxy-2-deoxyguanosine; 8-
406 OHdG) after resistance exercise compared to a placebo. Regulation of calcium homeostasis is another candidate
407 mechanism of creatine supplementation. Cooke et al. (2009) and Korge et al. (1993) reported that creatine assists in
408 maintaining the sarcoplasmic reticulum calcium pump function by phosphorylating ADP to ATP, which decreases
409 cytosolic calcium levels. Minajeva et al. (1996) suggested that increasing muscle Phosphocreatine levels accelerates ATP
410 homeostasis, leading to reduced secondary damage due to an increase of calcium concentration in sarcoplasmic reticulum.
411 However, the evidence for this idea in a muscle damage paradigm is thin and requires greater evidence.

412 In early work in this area, Willoughby and Rosene (2003) gave creatine to 22 untrained males (6 g/day for 12
413 weeks) followed by a bout of unaccustomed exercise. Results showed that the perception of muscle soreness was reduced
414 in the supplemented group. It was speculated that creatine helped to direct gene transcription and upregulate protein
415 synthesis. In a later study, Cooke et al. (2009) investigated the effects of five days of creatine supplementation (21 g per
416 day) and carbohydrate (84 g day) or only carbohydrate in 14 untrained healthy males on muscle function and muscle
417 damage indices after intense eccentric exercise. They showed that plasma CK levels were lower (by an average of 84%)
418 in the days following the exercise in the creatine group; furthermore, this group showed a greater isokinetic (+10%) and
419 isometric (+21%) knee extension strength following EIMD than carbohydrate group. Rosene et al. (2009) showed that
420 both shorter and longer-term creatine supplementation (20 g/day for the first 7 days followed by 6 g/day for 23 days)
421 improves maximal strength after exercise in 20 physically active males. Veggi et al. (2013) also showed that ingesting
422 20/day of creatine for 6 days between exercise bouts attenuates rises in CK, improved range of motion and decreased
423 muscle soreness.

424 Santos et al. (2004) also studied the effects of creatine supplementation (20 g/day for 5 days prior to the 30 km
425 race) on inflammatory markers and muscle soreness in experienced marathon runners. Results showed creatine
426 supplementation attenuated the increase of prostaglandin E2 (-60.9%), TNF- α (-33.7%), CK (-19%) and abolished the
427 increase of LDH levels after the run, compared to controls. Further, Bassit et al. (2010) in a small cohort (8 triathletes)
428 randomized into a placebo or creatine monohydrate supplementation group (20 g/day for 5 days). The authors reported
429 that creatine supplementation decreased plasma creatine kinase, LDH and aldolase, and prevented the rise of glutamic
430 pyruvic acid transaminase and glutamic oxaloacetic acid transaminase, when compared to placebo group. Likewise,
431 Deminice et al. (2013) supported previous data showing that seven days of creatine supplementation (0.3 g/kg of body
432 mass/d) in 25 healthy, well-trained male soccer players reduced the increase of TNF- α , CRP and LDH, after a repeated
433 sprint exercise protocol. In contrast, in a double-blind study where creatine supplementation (0.6 g/kg of body mass/d
434 for 7 days) or placebo was examined in sedentary individuals following damaging resistance exercise, no difference
435 between groups was observed (Machado et al. 2009).

436 Although several studies reported that creatine attenuated indices of EIMD (Willoughby and Rosene 2003;
437 Santos et al. 2004; Bassit et al. 2010, Cooke et al. 2009; Rosene et al. 2009; Deminice et al. 2013; Veggi et al. 2013) other
438 studies have reported that creatine had no effect (Rawson et al. 2001; Rawson et al. 2007; Mckinnon et al. 2012). For
439 example, Rawson et al. (2001) investigated the effects of creatine supplementation (20 g/day taken for 5 days before and
440 after exercise) in 23 non-weight-trained men, following eccentric contractions and showed no differences in muscle
441 damage indices the creatine and placebo groups. In a later study, Rawson et al. (2007) examined the effects of short-term
442 creatine supplementation on markers of muscle damage in 22 healthy, resistance-trained males. Participants consumed
443 creatine (0.3 g/kg of body mass for the first 5 days and 0.03 g/kg for the next 5 days) or a placebo. Following the initial
444 5 days of supplementation, subjects performed damaging resistance exercise. There were no differences between groups
445 suggesting that oral creatine supplementation did not reduce indices of EIMD. In further support of the null findings,
446 Mckinnon et al. (2012) reported that ingesting creatine (loading dose: 40 g split into two servings/day for the first 5 days,
447 followed by a maintenance period of 10 g/day split into two servings/day next 5 days) did not reduce indices of muscle
448 damage or speed recovery following eccentrically induced muscle damage. Finally, a recent study (Fernández-Landa et
449 al. 2020) failed to find any positive effect of combined creatine monohydrate and β -hydroxy β -methylbutyrate
450 consumption (creatine 0.04 g/day per kg of body mass, plus 3g/day of HMB) in elite male rowers on reducing indirect
451 markers of EIMD. Despite the mixed results, athletes might benefit from creatine supplementation to restore muscle
452 function and assisting to reduce delayed onset muscle soreness.

453

454 **Beetroot juice**

455 Red Beetroot (*Beta vulgaris rubra*) is a functional food rich in nitrate, and other phytochemicals that include
456 bioactive compounds such as betalains, ascorbic acid, carotenoids, phenolic acids and flavonoids. Several published
457 works have reported beneficial effects of beetroot ingestion in managing hypertension, atherosclerosis, type 2 diabetes
458 and dementia (Clifford et al. 2015; Siervo et al., 2018; Gilchrist et al., 2013). In addition, recent investigations have
459 showed that the consumption of beetroot juice after eccentric exercise could reduce the magnitude of EIMD. In the first
460 of a series of studies by the same laboratory, beetroot juice was used to manage signs and symptoms of exercise-induced
461 muscle damage (Clifford et al., 2016a). Using a double-blind, independent groups design, 30 recreationally active males
462 were randomized to consume a higher dose (250 mL per serving), or a lower dose of beetroot juice (125 mL per serving)
463 or an isocaloric placebo (250 mL). Three servings were given immediately following damaging drop jump exercise, and
464 2 servings after 24 and 48 h. Results showed a faster recovery of pressure pain threshold during a 72-h recovery period

465 with beetroot juice in comparison to the placebo. However, muscle function was only improved in the higher-dose group
466 compared to placebo. The same group adopted the same supplementation protocol in 29 recreationally active males that
467 performed two bouts of 100-drop jumps, separated by 14-21 days. In this trial, authors reported that supplementation with
468 beetroot juice did not adversely affect acute adaptations (repeated bout effect) to eccentric exercise (Clifford et al. 2017a).
469 In a further study (Clifford et al. 2016b) beetroot juice was applied as a post-exercise recovery strategy to attenuate EIMD
470 in 20 collegiate team-sports players. In this trial, the effect of 500 mL of beetroot juice or a placebo for 3 days was
471 examined after performing an initial repeated sprint test (20 x 30 m) followed by a second repeated sprint test, performed
472 72 h later. The supplemented group showed a faster recovery of countermovement jump (CMJ) and reactive strength
473 index (RSI) after the first repeated sprint test. After 72 h following damaging exercise, the supplemented group showed
474 an increased restoration of CMJ and RSI performance (7.6% and 13.8%, respectively) compared to placebo group
475 (Clifford et al. 2016b).

476 In an attempt to understand the role of dietary nitrate in the beetroot juice, a later study investigated the effects
477 of a nitrate only drink (sodium nitrate), beetroot juice or placebo drink on indices of muscle damage caused by exercise.
478 In this trial, Clifford et al. (2017b) recruited 30 recreationally active males that were equally randomized in to the 3 groups
479 and ingested the supplement immediately, 24 and 48 h (3 x 250 mL servings the first day, 2 x 250 mL second and third
480 day) following 100 drop jumps. Pressure-pain threshold was attenuated in the beetroot juice group, which was $104 \pm 26\%$
481 of baseline values at 72 h post exercise, compared to sodium nitrate ($94 \pm 16\%$) and the placebo ($91 \pm 19\%$) suggesting that
482 beetroot was more effective than sodium nitrate for attenuating exercise-induced muscle pain; thus, any analgesic effect
483 could be attributable to phytonutrients other than nitrate, or interactions between nitrate and other constituent compounds.
484 Similarly, Montenegro et al. (2017) supported previous results of Clifford et al. (2016a; 2016b; 2017b) reporting that a
485 betalain-rich beetroot concentrate, not containing nitrates (50 mg twice a day, before breakfast and evening, per 7 days),
486 attenuated the increase of creatine kinase after a 10 km running time trial in triathletes. Most recently, these previous
487 beneficial effects of beetroot juice supplementation were mirrored by Daab et al. (2020) in a randomized double-blind
488 cross-over design in thirteen soccer players. Seven days of beetroot juice supplementation (2 servings of 150 mL per day
489 for 3 days pre-exercise, day of test and 3 days after damaging exercise) was able to reduce muscle soreness and attenuate
490 the decrease of MVC and CMJ. One contrasting study (Clifford et al., 2017c) reported no effect of beetroot juice
491 consumption (3 x 250 mL servings/day for 3 days) on muscle damage indices exercise-induced, in 34 well trained runners
492 who completed a marathon.

493 Existing evidence has shown some potential beneficial effects of beetroot supplementation on preserving muscle
494 function and reducing exercise-induced pain; however, it should be made clear that any benefit appears to be very small
495 and is largely around perceptual measures. Whilst an optimal dosing strategy does not currently exist, beetroot is easily
496 applied for those who can tolerate the taste.

497

498

499 **Pomegranate juice**

500 Pomegranate (*Punica granatum* L.) is categorized as a fleshy berry that is rich in polyphenols and has shown
501 some potential for promoting recovery from EIMD. It contains malic and citric acid, flavanols, anthocyanins,
502 hydrolysable and condensed tannins (Kandylis and Kokkinomagoulos, 2020), in particular ellagitannins has anti-
503 inflammatory and antioxidant properties (Danesi and Ferguson, 2017) and precursors of urolithin A, which has been
504 associated with mitophagy and increased muscle function (Martinez-Sanchez et al. 2017). Consequently, a number of
505 studies have suggested that consumption of pomegranate juice could be an intervention to manage the effects of exercise-

506 induced muscle damage.

507 Trombold et al. (2010) investigated the effects of ellagitannin extract (from pomegranate) on recovery of muscle
508 strength after an eccentric exercise. In a crossover, placebo-controlled trial, 16 recreationally active males were
509 randomized either pomegranate extract (480 mL twice a day) or a placebo for 9 days. Participants ingesting the extract
510 had greater preservation of muscle function following damaging eccentric exercise compared to an isocaloric placebo.
511 The same group (Trombold et al. 2011) supported these data in a subsequent study conducted in 17 resistance-trained
512 subjects. Participants were assigned either 250 mL/day of pomegranate juice or a placebo for 15 days. On the eighth day
513 of supplementation, damaging eccentric exercise of the elbow flexors and knee extensors was performed. Findings
514 showed that pomegranate supplementation attenuated upper body strength loss and soreness compared to a placebo. No
515 effect was reported on lower body and the researchers hypothesised that knee extensors are more active during activities
516 of daily living and hence more accustomed to the exercise stimulus. It would, be prudent to consider the results arising
517 from cross-over design studies in EIMD paradigms, because of potential issues relating to the contralateral RBE
518 (Howatson and van Someren 2007), which show that the magnitude of the damage response on the contralateral limb can
519 be significantly attenuated and hence mask possible effects of an intervention.

520 Other work (Machin et al., 2014) determined if there was a dose response effect of pomegranate supplementation
521 on muscular strength recovery following a bout of combined upper and lower body eccentric exercise. Forty-five non-
522 resistance trained volunteers were randomized into: (1) once-daily (30 mL of pomegranate juice concentrate); (2) twice-
523 daily (60 mL of pomegranate juice concentrate), or (3) placebo, consumed for 8 days. After 3 days of supplementation,
524 subjects performed 20 minutes of downhill running and 40 eccentric contractions of the elbow flexors. During the 4-day
525 recovery period, both lower and higher doses facilitated the recovery of elbow flexion and knee extension strength when
526 compared to the placebo. Other positive responses were also observed with pomegranate juice by Ammar et al. (2016) in
527 professional male weightlifters. Subjects that consumed pomegranate juice had lower soreness during the 48-h recovery
528 period and smaller increases in CK and LDH post-exercise, compared to a placebo.

529 In contrast, Lamb et al. (2019) observed that resistance trained men who ingested 500 mL/day of tart cherry juice
530 or pomegranate juice, for 5 days before and 4 days after a damaging exercise, failed to enhance recovery from high-force
531 eccentric exercise of the elbow flexors. In addition, Torregrosa-Garcia et al. (2019) failed to show significant differences
532 for force restoration after pomegranate extract supplementation (225 mg/day of punicalgins) or placebo in 26 amateur
533 cyclists. Recent evidence suggested that pomegranate juice showed promise as a strategy to improve muscle recovery and
534 reduce indirect biomarkers of exercise-induced muscle damage.

535

536

537 **Interventions with a modest evidence base**

538

539 **L-citrulline and Watermelon juice**

540 L-citrulline is a colourless, water-soluble non-essential amino acid (Yang et al. 2019), an endogenous precursor
541 of L-Arginine and an indirect precursor of nitric oxide (Martinez-Sanchez et al. 2017). Furthermore, L-citrulline is an
542 essential component of the urea cycle in the liver and kidneys, being responsible for detoxification of ammonia via
543 conversion to urea (Martinez-Sanchez et al. 2017).

544 Pérez-Guisado and Jakerman (2010) investigated the acute effects of citrulline on muscle soreness in 41 men.
545 Participants who ingested l-citrulline (8 g of citrulline malate) 1 hour before the performing 8 sets of bench press exercise
546 to fatigue at 80% 1RM. They showed a decrease of 40% in muscle soreness at 24 and 48 hours after the damaging session,

547 compared to placebo or isocaloric drink. A further study (Tarazona-Diaz et al., 2013) investigated the effect of watermelon
548 juice containing 1.17 g or 6 g of L-citrulline or a placebo on muscle soreness in 7 men. Subjects who ingested the placebo
549 1 h prior to exercise session (8 x 30 s, with 1 min rest of cycle interval sprints) had greater soreness after 24 h of recovery
550 period compared to participants that consumed watermelon juice. After these initial studies, Martinez-Sanchez et al.
551 (2017a) reported that healthy male runners who consumed watermelon juice (3.45 g of L-citrulline per 500 mL) 1 h prior
552 to a half-marathon race, showed an attenuation of muscle soreness during 24-72 h of recovery and preserved muscle
553 function compared to the placebo. The same research group (Martinez-Sanchez et al. 2017b) investigated the effects of
554 two watermelon juices with different concentration of L-citrulline (0.5 g and 3.3 g per serving, respectively) or a mix of
555 watermelon juice and a concentrate of pomegranate enriched in L-citrulline (3.3 g per serving) and ellagitannins (22.0
556 mg per serving) compared to a placebo beverage in nineteen healthy males. Subjects consuming a serving of watermelon
557 juice with a concentration of 3.3 g of L-citrulline or the beverage mix containing 3.3 g L-citrulline and 22.0 mg
558 ellagitannins, prior to a half-squat exercise (8 sets of 8 repetitions) reported a reduced muscle soreness during 48 h of
559 recovery, exhibited lower increase of myoglobin and showed a significant maintenance of muscle function compared to
560 the placebo group.

561 In contrast to these promising results with watermelon juice enriched with L-citrulline other studies have reported
562 no benefit (Shanely et al. 2016; da Silva et al. 2017; Chappell et al. 2018). For example, Shanely et al. (2016) reported
563 no effect of watermelon puree consumption (980 mL/day for 14 days) on markers of inflammation in trained cyclists after
564 a 75 km time trial. However, it is important to indicate that cycling (a predominantly concentric contraction sport) is not
565 considered to be a damaging *per se*. Similarly, da Silva et al. (2017) found no effect of supplementation with 6 g of
566 citrulline malate on muscle soreness or serum CK during the 72 h recovery period after a leg press and hack squat exercise
567 when compared to placebo control. Finally, a more recent study (Chappell et al., 2018) reported that acute citrulline
568 malate supplementation (4.21 g citrulline and 3.79 g malate) actually *increased* total muscle soreness during the 72-h
569 period after isokinetic knee extensions (10 sets of 10 repetitions, 70% MVC) compared to placebo (6 g of citric acid) in
570 5 females and 13 males resistance-trained. This is one of the very few examples where a functional food has been shown
571 to have a negative effect as opposed to a positive or null effect. On balance, L-citrulline and watermelon show some
572 benefit to decreasing muscle soreness and preserving muscle function after EIMD.

573

574 **β -hydroxy β -methylbutyrate (HMB)**

575 β -hydroxy β -methylbutyrate (HMB) is a metabolite of the amino acid leucine and a potent stimulus for translation
576 initiation and protein synthesis (Fitschen et al. 2013). Research during the last 20 years has investigated the effects of
577 HMB supplementation on muscle soreness, indirect markers of muscle damage (Knitter et al. 2000; Paddon-Jones et al.
578 2001; Hoffman et al. 2004; van Someren et al. 2005; Nunan et al., 2010; Wilson et al. 2009; Wilson et al. 2013; Wilson
579 et al. 2014) and circulating levels of pro-inflammatory cytokines (Portal et al. 2011; Gonzalez et al. 2014; Hoffman et al.
580 2016; Arazi et al. 2018b; Arazi et al. 2019) and showed mixed results. The first study (Knitter et al., 2000) to investigate
581 the effects of HMB on EIMD used six weeks supplementation (3 g/day) prior to a 20 km run. The HMB supplementation
582 was shown to reduce CK and LDH response after a prolonged run. Further work (van Someren et al., 2005) showed that
583 HMB supplementation (3 g/day) associated with alpha ketoisocaproic acid (0.3 g/day) in eight male volunteers, for 14 d
584 prior a damaging exercise bout of biceps curls at 70% of 1-RM, resulted in less muscle soreness at 24 h post bout, lower
585 CK increases and overall in a greater muscle function, compared to a placebo. Similarly, beneficial effects of
586 supplementation with the free acid form of HMB (3 g per day) was shown by Wilson et al. (2014) over a 12 weeks period.
587 In addition, Hoffman et al. (2016) showed that 23 days of HMB supplementation in combat soldiers attenuated the

588 inflammatory response (IL-8, IL-10, TNF-alpha, granulocyte colony-stimulating factor, fractalkine and interferon- γ) to
589 intense military training, and preserved muscle quality. This suggested that a longer duration supplementation period
590 might be beneficial to influence inflammatory indices. Promising data also comes from the acute consumption of the free
591 acid form of HMB (3 g) that when ingested prior to a maximal eccentric exercise to attenuate LDH levels (Wilson et al.
592 2009), and reducing the increase of CK and improved the perceived readiness to train after a muscle-damaging resistance-
593 training (Wilson et al. 2013). More recent investigations have suggested that free acid form of HMB has greater
594 bioavailability in a shorter time when compared with calcium salt form of HMB (Arazi et al. 2019) and its
595 supplementation might represent a useful strategy to modulate exercise-induced muscle damage (Asadi et al. 2017) and
596 attenuate exercise-induced oxidative stress (Arazi et al. 2018b; Arazi et al. 2019).

597 In contrast to the aforementioned beneficial effects of HMB ingestion, Paddon-Jones et al. (2001) reported no
598 effects of 6 days of HMB supplementation (40 mg/kg body mass per day) prior to a bout of maximal isokinetic eccentric
599 contractions of the elbow flexors in non-resistance trained subjects. Additionally, Hoffman et al. (2004) studied the effects
600 of 10 days of HMB supplementation (3 g/day) and revealed that short duration of HMB supplementation did not provide
601 any benefit in reducing indirect markers of muscle damage (CK and MYO) during a preseason football training camp. In
602 a later study, Nunan et al. (2010) did not find positive effects of 14 days (11 days before and 3 days after a damaging
603 exercise) of 3 g/day of HMB associated with 0.3 g/day of alpha ketoisocaproic acid on muscle function, CK levels or
604 DOMS following a 40-min bout of downhill running, in fourteen untrained male subjects. This contradicted their previous
605 work (van Someren et al. 2005) that did show some benefit using the same supplement in an isolated muscle group.
606 Further, Portal et al. (2011) reported that seven-week consumption of HMB (3 g/day) did not change serum levels of IL-
607 6 and IL-1 during the early phase of training in elite volleyball players. Similarly, to those find null effects, Gonzalez et
608 al. (2014) showed that the acute administration of the free acid form of HMB (3 g) in recreationally resistance trained
609 males did not attenuate post-exercise markers of muscle damage, inflammation or performance following lower body
610 resistance exercise, when compared to placebo or a combination of cold water immersion and HMB-FA or placebo plus
611 cold water immersion. Studies have shown that both acute and chronic supplementation of HMB prior to exercise might
612 improve functional and biochemical markers of EIMD, however, it should be noted that this conclusion is based on a
613 limited number of studies, which are largely done in untrained individuals who are unaccustomed to eccentric exercise,
614 so the applicability to well-trained athletes is not clear.

615

616 **Protein and amino acids**

617 It is well known that protein intake plays a crucial role in the regulation of muscle protein turnover, particularly
618 in response to exercise (Phillips and Van Loon 2011), and that adequate intake should never be compromised in the diet.
619 However, there are doubts about the role of protein and amino acids supplementation in the prevention and treatment of
620 EIMD (Wojcik et al. 2001; Tipton et al. 2003; Blacker et al. 2010, Tipton 2015, Eddens et al. 2017). A systematic review
621 (Pasiakos et al. 2014) of the effects of protein ingestion on EIMD-related outcomes concluded that protein
622 supplementation had little effect on muscle damage which was attributable to large variations in study design and the
623 biomarkers selected. In addition, Poulos et al. (2019) confirmed there was no clear evidence to support a relationship
624 between changes in muscle damage indices with protein-based supplementation. Nevertheless, other studies have
625 suggested that hydrolysate whey protein (25 g) ingested immediately after the completion of maximal eccentric
626 contractions could alleviate muscle damage and enhances recovery of force following damaging exercise in sedentary
627 males (Buckley et al. 2010) cyclists (Eddens et al., 2017) and physically active females (Brown et al. 2018), or after
628 ingestion of isolated soy protein (25g) from boxers and cyclists (Shenoy et al. 2016). More recently, Ma et al. (2020)

629 reported that young runners who ingested ‘immune protein’ (10 g/day over a period of 2 months) obtained from cows
630 immunised against specific antigens, had reduced inflammatory markers (TNF- α and IL-1 β) post exercise, when
631 compared to placebo treatment.

632 A systematic review by Fouré and Bendahan (2017) concluded that at least 200 mg per kg of body mass per day
633 of branched chain amino acids (BCAA) for at least 10 days could be effective prophylactic to combat moderate muscle
634 damage. In addition, a recent meta-analysis indicated that BCAA supplementation reduced delayed onset muscle soreness
635 following exercise training (Fedewa et al. 2019). Likewise, Rahimi et al. (2017) reviewed randomized clinical trials and
636 concluded that BCAA supplementation, before and during recovery days after damaging exercise, for at least one week,
637 is able to attenuate muscle strength and muscle power loss. Previously, Nosaka et al. (2006) suggested that the ingestion
638 of 9 essential and 3 non-essential amino acids before and after damaging exercise (7.2 g/day and for 5 days) attenuated
639 indices of EIMD. Similar results were shown (Howatson et al., 2012) where BCAA supplementation for a total of 12 days
640 (10 g twice a day) before and following damaging resistance exercise reduced indices of muscle damage in resistance-
641 trained males. Similarly, Waldron et al. (2017) investigated the acute supplementation of BCAA (0.087 g/Kg of body
642 mass, prior exercise) or placebo on EIMD outcomes in resistance trained athletes and showed improved rates of recovery
643 in muscle function indices compared to placebo. Osmond et al. (2019) also reported the beneficial effect of BCAA (10 g
644 twice a day per 11 days enriched with 5 g of additional leucine) on recovery following 100 drop jumps and afford no
645 recovery benefits.

646 Collectively, despite the heterogeneity in study design and biomarkers, athletes might benefit from a BCAA
647 supplementation (ratio 2:1:1 of l-leucine, valine, isoleucine) and hydrolysate whey protein to combat signs and symptoms
648 of EIMD. However, further research is required to better understand the implications of different dosing and timing
649 strategies on EIMD in addition to gaining a greater understanding of such interventions in well trained athletes.

650

651 **Milk**

652 Cows’ milk contains ~80% casein and ~20% whey protein, thus providing a good balance of slow and fast digested amino
653 acids to the muscle, which some researchers have hypothesized would have a beneficial effect on muscle recovery. Milk
654 is the combination of a protein-carbohydrate beverage that has the potential to resolve EIMD (Cockburn et al., 2013). In
655 a series of studies (Cockburn et al., 2008; 2010; 2012; 2013) showed that milk ingestion consumed post damaging exercise
656 attenuated signs and symptoms of EIMD, such as muscle soreness and muscle function in team and individual sports. A
657 further study (Rankin et al., 2015), male and female team sport athletes ingested 500 ml of milk immediately following a
658 damaging eccentric-concentric exercise and showed smaller increases in soreness and serum CK and skeletal muscle
659 troponin I compared to an energy-matched carbohydrate solution. Some athletes are intolerant to lactose and/or the A1
660 beta-casein protein, but A2 milk addresses these issues by offers an alternate, but otherwise identical milk (Kirk et al.
661 2017). Similarly, to whey hydrolysate protein and BCAA, milk could represent a potentially efficacious intervention,
662 although the mechanism(s) underlying its positive effects is not fully understood; critically, milk is readily available to
663 most individuals and is inexpensive,

664

665 **L-glutamine**

666 Glutamine is the most abundant amino acid in skeletal muscles and plasma and is an important fuel for immune
667 system cells (Street et al. 2011). It has beneficial antioxidant and anti-inflammatory properties, so supplementation is
668 considered potentially useful for athletes; for example, and recent research suggested that L-glutamine ingestion can
669 restore plasma concentrations of glutamine (Nemati et al. 2019). In the setting of EIMD, L-glutamine was supplemented

670 for a short period, using a dosing protocol of 0.3g/Kg/day for up to 7 days, with beneficial effects on muscle function
671 (Street et al. 2011) and on muscle soreness (Legault et al. 2015). Using higher doses (1.5g/kg/day for 7 days), Nakhostin-
672 Roohi et al. (2016) showed beneficial effects on the CK response following strenuous exercise. However, at lower doses
673 taken for longer periods (0.1g/kg/day for 28 days) L-glutamine did not improve muscle function or ameliorate muscle
674 soreness (Rahmani et al., 2013). Although some rationale exists for the use this supplement to sustain immune function,
675 further studies are needed to understand the application of recovery from EIMD.

676

677 **L-Carnitine**

678 L-Carnitine has the potential to ameliorate muscle damage signs and symptoms. In a recent review and meta-
679 analysis, Yarizadh et al. (2020) described that L-carnitine supplementation resulted in significant improvements in muscle
680 soreness following damaging exercise in both resistance-trained and untrained populations, when compared to a placebo.
681 An earlier study (Giamberardino et al., 1996) showed that supplementation of L-carnitine (3g/day) alleviated pain,
682 tenderness and appearance of CK in blood, indicating the potential role of the nutrient in reducing tissue disruption and
683 leakage of cytosolic proteins. Other studies (Spiering et al., 2008; Kraemer et al., 2003; Volek et al., 2002; Spiering et al.
684 2007) also reported the ability of L-carnitine to reduce indices of muscle damage. For example, a daily intake of 2 g of
685 L-carnitine compared to a placebo was accompanied by a reduction in released CK, MDA as well as hypoxanthine and
686 xanthine oxidase (Spiering et al., 2008; Volek et al., 2002). Spiering et al. (2007) reported that both 1 g/day and 2g/day
687 of L-carnitine supplementation, provided comparable benefits to modulate muscle soreness and indirect markers of
688 muscle damage. Kraemer et al. (2006) investigated the effect of L-carnitine and L-tartrate supplementation over a period
689 of 3 weeks and showed an increase of the level of androgen receptors on muscle cells, thus improving protein signalling
690 that is necessary for recovery after exercise. Further, Parthimos et al. (2008) showed one month of L-carnitine
691 supplementation (2 g/day) was able to maintain total antioxidant status, but it was unaltered post-exercise in basketball
692 players. Similarly, Parandak et al. (2014) reported that 2g/day of L-carnitine over 14 days increased total antioxidant
693 capacity before and 24 h post-exercise, when compared to a placebo, but no differences in muscle damage markers and
694 lipid peroxidation were observed. Although these promising benefits, the exact mechanism(s) through L-carnitine
695 improve outcomes of EIMD remain unknown and further investigations are warranted.

696

697 **Curcumin**

698 Curcumin is a natural compound and bioactive polyphenol found in turmeric (2%-5% by weight). It could
699 represent a promising and beneficial strategy for attenuating EIMD because of the potential antioxidant and anti-
700 inflammatory effects (Fernández-Lázaro et al. 2020). In line with this, some studies (Gaffey et al. 2017; Harty et al. 2019)
701 have shown the antioxidants and anti-inflammatory effect of curcumin. It has can also attenuate muscle damage from
702 thermal shock (Dunsmore et al. 2001) and reduce the increased activity of biomarkers of muscle damage such as CK
703 (Kocaadam and Sanlier 2017). Supplementation of curcumin on sport performance and EIMD has been applied in elite
704 rugby players (Delecroix et al. 2017), in moderately active subjects (Sciberras et al. 2015; Drobnic et al. 2014; Jäger et
705 al. 2019; Nicol et al. 2015; Basham et al. 2019) and relatively inactive participants (Tanabe et al. 2019a; Tanabe et al.
706 2015; Nakhostin-Roohi et al. 2016; Tanabe et al. 2019b; McFarlin et al. 2016), but there are inconsistencies in findings
707 across these studies.

708 Nicol et al. (2015) described that curcumin supplementation (5 g/day) taken 48 h before, and 72 h after eccentric
709 exercise, reduced muscle pain (VAS scale -1.4 to -1.7) and blunted the CK activity (-22-29%) following damaging
710 exercise (7 sets of 10 eccentric single-leg press repetitions), although paradoxically the post-exercise (at 0-h) IL-6 was

711 elevated relative to placebo (+31%). McFarlin et al. (2016) reported that CK, TNF- α and IL-8 responses were blunted (-
712 48%, -25%, -21%, respectively) in subjects who consumed curcumin (400 mg/day for 2 days) prior to, and for 4 days
713 after, damaging eccentric leg-press exercise; however, there were no differences in muscle soreness between
714 supplemented and placebo groups. A similar response (Tanabe et al. (2015) of curcumin (150 mg 1 h prior to and 12 h
715 after damaging eccentric elbow exercise) was observed, where declines in MVC were lower and recovery was faster (e.g.,
716 4 days post-exercise: $-31 \pm 13\%$ vs. $-15 \pm 15\%$) and peak CK was also smaller (peak: 7684 ± 8959 IU/L vs. 3398 ± 3562
717 IU/L) than the placebo condition. Two studies from the same group (Tanabe et al. 2019a; Tanabe et al. 2019b) showed a
718 significant reduction in muscle pain measured after 3-4 days (VAS scale) in subjects that ingested curcumin (90 mg twice
719 daily) when administered four (Tanabe et al. 2019a) and seven (Tanabe et al. 2019b) days after eccentric exercise. In
720 addition, compared to a placebo group, MVC torque and ROM were higher 3-7 days and 2-7 days after exercise,
721 respectively (Tanabe et al. 2019b). These findings were supported by Jäger et al. (2019). who showed that 200 mg of
722 curcumin was effective in preventing the decrease in MVC at 1 and 24 h after damaging exercise and showed less muscle
723 soreness compared to placebo or lower dose of curcumin (50 mg). Similarly, Nakhostin-Roohi et al. (2016) showed that
724 150 mg of curcumin taken immediately after exercise reduced muscle pain at 48 and 72 h after eccentric exercise and was
725 able to lower maximum CK, AST and ALT. Further evidence (Basham et al., 2019) for curcumin supplementation (1.5
726 g/day for 28 total days) abating EIMD exists, where 19 males completed 15 min of continuous sitting with one leg (a total
727 of 225 repetitions). They showed that curcumin blunted CK (199.62 U/L vs 287.03 U/L) and DOMS (2.88 vs 3.36)
728 compared to the placebo. Most recently, Amalraj et al. (2020) reported that oral consumption of curcumin (Cureit™),
729 which was developed based on the restructuring of the turmeric matrix with active curcumin using polar/nonpolar-
730 sandwich (PNS) technology (Amalraj et al., 2017), decreased the pain VAS score (-59.77%) and the CK levels (-15.55%)
731 following eccentric exercise. The PNS technology could be a strategy to overcome the limitations of curcumin
732 bioavailability by increasing concentrations of free curcuminoids in the blood plasma and hence be beneficial in managing
733 pain and DOMS (Jude et al., 2018).

734 In contrast with previous data, other researches failed to demonstrate benefits of curcumin supplementation
735 (Delecroix et al. 2017; Sciberras et al. 2015; Drobnic et al. 2014) on EIMD outcomes. For example, Drobnic et al. (2014)
736 reported a moderate reduction in pain at 48 h after downhill running (total score: 23.3 ± 7.9 vs. 30.6 ± 7.9 , curcumin and
737 placebo group, respectively) with 200 mg curcumin supplementation taken 48 h before and 24 h after exercise. Delecroix
738 et al. (2017) showed that curcumin (6g/day) and piperine (60 mg/day) supplementation (48 h before and 48 h after
739 exercise) in elite rugby players had a limited effect on power loss during the one leg 6 s sprint, without any effect on other
740 aspects of muscle damage. Sciberras et al. (2015) showed that curcumin supplementation failed to produce differences in
741 inflammatory markers after 2 hours of endurance cycling. The authors suggested that the lack of significance could be
742 related to the sample size, mode and intensity of exercise or dose of curcumin used.

743 The mixed findings in studies could be the result of several limitations, including functional tests used, study
744 settings and timing of supplement ingestion. Importantly, curcumin has poor aqueous solubility, has low absorption
745 volume from the gut, is rapidly metabolised, and is excreted quickly (Anand et al., 2007). While conjugated curcumin
746 achieves greater plasma concentrations around 1 hour after ingestion, a substantial amount of conjugated curcumin (up
747 to a third) can be detected in plasma 24 hours post administration compared to only trace levels of non-conjugated
748 curcumin (Asai and Miyazawa 2000). Although studies support that curcumin conjugates maintain similar molecular
749 properties as the parent curcumin (Sandur et al. 2007; Pfeiffer et al. 2007; Kim et al. 1998; Pari and Murugan 2006),
750 whether the magnitude of their bioactivity is comparable to parent curcumin remains unclear. Importantly, curcumin, like
751 many other polyphenols, has poor bioavailability that can be improved with piperine co-ingestion. Piperine is the major

752 component of black and long peppers and has been shown to inhibit enzymatic conjugation of curcumin allowing greater
753 levels of unconjugated curcumin to be absorbed into blood (Shoba et al. 1998) and increase curcumin tissue retention
754 time. Notwithstanding, it appears that curcumin has good potential as an intervention to attenuate EIMD, particularly if
755 metabolites are bio-available in a 24 h window.

756

757 **Bromelain and Proteases**

758 Bromelain is a mixture of proteases obtained from the stems of the developing pineapple fruit. Castell et al.
759 (1997) showed that bromelain is absorbed and remains active following digestion with an estimated half-life of 6-9 h and
760 that supplementation in small doses with other proteases has the potential to attenuate reductions in muscle function after
761 damaging eccentric exercise (Buford et al. 2009; Miller et al. 2004). Miller et al. (2004) investigated the effect a combined
762 protease supplement (4.2 g per day, one day and 4 days) prior to a 30 min downhill run. They reported a faster recovery
763 of muscle function and reduced muscle soreness during the 72 h recovery period compared to a placebo group. Similarly,
764 Beck et al. (2007) showed that strength loss immediately following eccentric elbow flexor contractions was reduced
765 following acute protease supplementation. In addition, Buford et al. (2009) showed that protease supplementation (5.8 g
766 per day for 21 days before downhill running and 3 days after) was able to attenuate decrements in MVC, a rise in basophil
767 and eosinophil count and lower pro-inflammatory cytokines (IL-6, IL-12), cyclooxygenase-2. Conversely, Udani et al.
768 (2009) found that a protease-containing supplement (258 mg of a proteolytic enzyme blend that included bromelain and
769 other proteases) for 30 days prior eccentric exercise in untrained subjects did not modulate CK, CRP or cytokine activity.
770 However, it did result in less tenderness 24 h post-exercise and less pain after 48 h of recovery compared to placebo. A
771 more recent study by Shing et al. (2016) examined if markers of muscle damage and testosterone was influenced by acute
772 bromelain in fifteen competitive cyclists that participated in a 10-stage cycle race over six consecutive days. Participants
773 who ingested 1000 mg of bromelain (1500 GDU) twice a day for 6 days, reported lower sensations of fatigue and tended
774 to maintain serum-free testosterone concentration across the race period compared to a placebo group. The exact
775 mechanism(s) underlying the potential ergogenic effect of protease supplementation is poorly understood, but it might be
776 related to reduced inflammation and hence improve recovery of strength loss after exercise (Beck et al. 2007). Thus,
777 bromelain in isolation might have a limited effect, but there could be benefit when used in combination with other protease
778 inhibitors. However, additional research is needed to understand the effects of proteases on EIMD in athletes and the
779 underlying physiological mechanisms.

780

781 **Caffeine**

782 The effects of caffeine in reducing fatigue and increasing wakefulness, alertness (Burke 2008) and modulating
783 pain perception through the action on adenosine receptors is well documented (Baratloo et al. 2016); however, limited
784 research has examined caffeine's effects on exercise-induced muscle damage. Hurley et al. (2013) reported that caffeine
785 consumption (5 mg/kg) 1 h before a damaging elbow flexion and extension exercise resulted in lower muscle soreness at
786 48 h and 72 h of a 5-day recovery period in resistance-trained males when compared to placebo group. This analgesic
787 effect was later supported by Caldwell et al. (2017) with a lower dose of caffeine (3 mg/kg body mass) in males and
788 females cyclists involved in a 164 km cycle ride. Furthermore, previous work (Chen et al. 2019) showed that acute
789 caffeine (6 mg/kg body mass) facilitated recovery and attenuated muscle soreness after 30-min downhill running in a
790 mixed sex cohort of college athletes.

791 Conversely, a series of studies conducted by Machado et al. (2009a; 2009b) found no effect following caffeine
792 ingestion (5.5 mg/kg body mass) on biomarkers of muscle damage (serum CK and LDH) in professional soccer players.

793 The same group (Machado et al. 2010) reported that ingestion of caffeine at 4.5 mg/kg body mass, before exercise, did
794 not reduce markers of muscle damage or leukocytes following moderate intensity resistance exercise session. Previously,
795 Vimercatti et al. (2008) reported no benefit of caffeine supplementation on immune responses or muscle cell integrity. In
796 this study, physically active males performed 60-min of treadmill exercise at 65% of VO_{2max} following caffeine ingestion
797 (4.5 or 5.5 mg/kg body mass 1 h before exercise) or placebo. There were no differences in blood biomarkers of damage
798 and immune function between groups.

799 These data suggest caffeine could have an analgesic effect through the inhibition of adenosine on the central
800 nervous system (CNS) by blocking the pain perception from the peripheral nerves to the CNS via adenosine receptors
801 (Hurley et al. 2013). Thus, caffeine ingestion could be an intervention for individuals who are particularly susceptible to
802 muscle soreness following training sessions or competitions. Although some recent studies demonstrated that caffeine
803 ingestion was able to modulate others indices of EIMD such as muscle soreness (Hurley et al. 2013; Karabalaefar et al.
804 2013, Maridakis et al. 2007) and performance (Hurley et al. 2013; Maridakis et al. 2007), there is a need for a greater
805 understanding of the effects of time of day ingestion, habituation of use, training status and prolonged use of caffeine on
806 post-exercise recovery.

807

808 **Taurine**

809 Taurine is a β -amino acid classified as a conditionally essential nutrient found in most human cells, with levels
810 particularly high in excitable tissues such as skeletal and cardiac muscle (Schaffer and Kim 2018; De Carvalho et al.
811 2017). This amino acid is commonly known for its effects as an energizer and anti-fatigue properties; beyond this, it can
812 exert other physiological functions that including membrane stabilization, osmoregulation and cytoprotection (De Luca
813 et al. 2015). Regarding supplementation, taurine consumption is linked to diminished body mass index and reduced
814 inflammation in obese women (Schaffer and Kim 2018). It is also reported to have antioxidant effects *in vivo* and a
815 regulatory effect of calcium homeostasis on ion channel function in cardiac and skeletal muscle (De Carvalho et al. 2017).
816 Recently, it has been reported that the ingestion of taurine might improve muscle function and reduce indices of EIMD.
817 Specifically, Ra et al. (2013) investigated whether a combination of taurine and BCAA could attenuate indices of muscle
818 damage in untrained males. The authors reported no effect of taurine supplementation alone (2 g/day, three times daily,
819 for 14 days prior to, and for the 3 day recovery period), but when taurine was combined with BCAA (2 g/day and 3.2
820 g/day three times per day, respectively) a reduction of soreness and other damage indices were reduced in comparison to
821 a placebo. Although the authors reported a synergic effect between taurine and BCAA, based on the evidence, it might
822 be that 9.6 g of BCAA ingested per day could expedite signs and symptoms EIMD. However, to reinforce the potential
823 use of taurine alone, Ra et al. (2015) showed that (2 g/day, three times daily, for 14 days) attenuated muscle soreness after
824 an eccentric elbow flexion exercise during the 4-days recovery period. These data were supported by McLeay et al. (2017)
825 who reported a faster recovery of eccentric torque in healthy active males who consumed taurine supplement (0.1g/kg
826 body mass per day, 30 min prior exercise and during 3 days after exercise) after a bout of damaging elbow flexors exercise.
827 These data concur with da Silva et al. (2014) who employed taurine a daily dose of 50 mg/kg body mass for 14 days
828 before damaging exercise and for 7 days after, in 21 males. Finally, De Carvalho et al. (2017) administered a combination
829 of taurine (3 g) plus 400 mL of chocolate milk ingested post exercise for 8 weeks. Results of this study indicated decreases
830 in oxidative stress markers in triathletes, compared to a placebo. Taurine could be a useful tool, but data are far from
831 conclusive. Notwithstanding, taurine might be a useful synergist supplement to other nutrients, but requires further
832 investigation.

833

834 **Ginger**

835 Ginger (*Zingiberofficinale Roscoe; Zingiberaceae*) is a popular tropical spice used as a dietary supplement and
836 food ingredient (Sellami et al. 2018). Several scientific reviews and meta-analyses have examined the use of ginger for
837 the treatment of vomiting and nausea, cancer prevention, pain management and inflammation (Wilson 2015). Although
838 the exact analgesic mechanism(s) is unclear, it seems that ginger blocks the activity of COX enzymes, leukotriene and
839 prostaglandin synthesis. In addition, ginger inhibits the release of proinflammatory cytokines that play a role in
840 exacerbating EIMD (Wilson 2015).

841 Black et al. (2010) showed in untrained participants, who ingested 2 g raw or heat-treated ginger per day for 11
842 days, less pain during the 24 h after eccentric elbow flexor exercise, relative to placebo. Matsumura et al. (2015) showed
843 beneficial effects of ginger powder supplementation (4 g/day for 5 days) on a faster recovery of elbow flexor 1-RM
844 strength during the 48 h recovery period after a damaging exercise, compared to placebo. Hoseinzadeh et al. (2015)
845 reported that untrained females who completed 20 minutes of stepping exercise following 60 mg of ginger extract 1 h
846 before showed a reduction of pain and IL-6 during the 48 h recovery period, when compared with consuming ginger
847 immediately after the exercise and a placebo. This suggested that the timing of the ginger supplementation is an important
848 consideration. Furthermore, Wilson et al. (2015) evaluated the effects of ginger root supplementation (2.2 g/day for 3
849 days before, the day of, and the day after exercise) on muscle function following a 20-22 mile training run, among 20
850 college students. Although median muscle soreness during jogging at 24 h after the training run was lower in the ginger
851 root group compared to placebo, muscle function recovery was not different between groups.

852 Conversely, Black and O'Connor (2008) reported that an acute dose of ground ginger (2 g) ingested 30 minutes
853 before 30 minutes cycling at 60% VO₂ peak in untrained females and males showed no differences in pain perception
854 between treated and placebo groups. A subsequent study by the same authors (Black and O'Connor, 2010) noted that an
855 acute dose of 2 g of ginger consumed by untrained participants 24 h after eccentric elbow exercise did not induce any
856 improvements on pain intensity, range of motion, arm volume and metabolic rate the following day, relative to placebo.
857 Overall, the limited evidence indicates no clear benefit of ginger on indices of EIMD.

859 **Ginseng**

860 There are several species of ginseng (Asian, Korean, Chinese, Canadian, Siberian and American (Sellami et al.
861 2018) with varying levels of active phytochemical compounds, the most important of which, is probably ginsenosides.
862 Ginsenosides are thought to modulate immune function, glucose metabolism, enhance cognitive performance and affect
863 oxidative stress (Bach et al. 2016). Recent evidence suggested that ginseng might have an important role in modulating
864 markers of EIMD. Hsu et al. (2005) examined the influence of American ginseng (1.6 g/day for 4 weeks) on a damaging
865 exercise bout of treadmill running in active college males. The authors showed lower levels of CK, compared to placebo,
866 in the days after exercise. Likewise, Jung et al. (2011) showed that post exercise CK and IL-6 were reduced in male
867 college students who ingested Korean red ginseng extract (20 g per 3 times/day for seven days prior, and for four days
868 after eccentric exercise) compared to placebo group. Further, Pumpa et al. (2013) had well trained males complete
869 intermittent downhill running treadmill exercise (5 bouts of 8 min running at -10% gradient) following consumption of
870 4g of Chinese ginseng 1 h before plus another 4g immediately after the exercise bout. They showed that muscle function
871 was preserved and muscle soreness and pro-inflammatory cytokines (IL-6 and TNF- α) were reduced with a ginseng.
872 Finally, Caldwell et al. (2018) investigated the effects of two doses of ginseng on perceived muscle soreness after
873 resistance exercise in males and females. The authors showed that both lower (160 mg/day) and higher doses (960
874 mg/day) of Korean ginseng ingested 14 days prior to strenuous resistance exercise reduced changes in muscle soreness

875 at 24 post exercise. Whilst these preliminary data are promising, this strategy warrants further investigation.

876

877 **Vitamin C and Vitamin E**

878 Vitamin C (e.g. L-ascorbic acid) and vitamin E (α -tocopherol) are two essential micronutrients, with redox-
879 dependent and independent biochemical functionality (Owens et al., 2019). Several studies have investigated the effects
880 on EIMD indices and save few exceptions (Gabrial et al. 2018; Righi et al., 2020), the general consensus is that both
881 vitamins show limited ability to positively impact the signs and symptoms of EIMD (Thompson et al. 2001; Thompson
882 et al. 2004; Shafat et al., 2004; Connolly et al., 2006; McGinley et al., 2009; Sousa et al., 2014; Candia-Luján et al., 2014;
883 de Oliveira et al. 2019; Owens et al. 2019).

884 Gabrial et al. (2018) examined the role of vitamin C intake in attenuating markers of muscle damage, oxidative
885 stress and inflammatory responses in 20 recreationally active males that were regularly involved (2-3 times per week) in
886 strenuous recreational exercise. The authors showed that longer term vitamin C ingestion (500 mg/day for 90 days)
887 reduced CRP and uric acid when compared to a placebo group. Most recently, Righi et al. (2020) systematically reviewed
888 the effects of vitamin C supplementation on oxidative stress, indirect markers of muscle damage and musculoskeletal
889 functionality after a single bout of exercise; they reported that vitamin C supplementation attenuated lipid peroxidation
890 and IL-6 elevation following a single session of damaging exercise.

891 Conversely, de Oliveira et al. (2019) showed no effects of vitamin C (500 mg/day) and vitamin E (400 UI/day)
892 supplementation 7 days before and 7 days after plyometric exercise on EIMD indices. Notably the dosing strategy was
893 much shorter than the aforementioned studies. In addition, a Cochrane review suggested a moderate to low quality of
894 scientific evidence for the use of high dose of “antioxidant” supplements in reducing delayed onset muscle soreness
895 (Ranchordas et al. 2017). Similarly, a previous systematic review reported the low effectiveness of vitamin C and E
896 supplementation in reducing DOMS (Candia-Luján et al. 2014) that was echoed by other work (Connolly et al., 2006)
897 that also found no benefit of vitamin C supplementation (1000 mg/day) ingested 3 days before and 5 days after damaging
898 exercise. Two studies (Thompson et al., 2001; 2004) suggested that supplementation with vitamin C 400 mg/day for 14
899 days did not affect IL-6 levels after a downhill running exercise (Thompson et al. 2004); nor did a higher dose of vitamin
900 C (1000 mg/day for 14 days prior to, and 2 h before) attenuate variables following 90 min of intermittent shuttle-running
901 test (Thompson et al. 2001). In a recent review, Owens et al. (2019) highlighted the issues of measuring free radical and
902 non-radical species to quantify redox balance in humans and suggested that vitamin C and vitamin E are unlikely to
903 directly scavenge free radicals in the phagosome that interfere with the inflammatory responses. Thus, the use of vitamin
904 E and vitamin C to modulate EIMD outcomes appears to lack support and further well-designed studies are warranted to
905 better understand if lower doses for a longer period or a periodized approach to supplementation with either vitamin C or
906 E alone, or in combination, in addition to a balanced and varied diet, can be used to reduce signs and symptoms of EIMD.

907

908 **Nutritional interventions with little or no evidence to support efficacy**

909 Exercise scientists have identified other potential interventions as solutions to manage the negative effects
910 associated with strenuous physical activity. This section of the review offers a brief overview of a wide assortment of
911 nutritional options with little evidence to support efficacy but that in the next future could attract researchers for additional
912 investigations.

913 One of the most studied rich sources of polyphenols is green tea. Previous studies showed that green tea extract
914 from *Camelia sinensis* supplementation reduced oxidative stress (Sugita et al., 2016), reduced muscle soreness (Herrlinger
915 et al., 2015), decreased other indirect markers of muscle damage after eccentric exercise (da Silva et al., 2018), and

916 maintained neuromuscular function after a submaximal bout of exercise (Machado et al., 2018). Some promising
917 beneficial effects were also observed following consumption of flavin-enriched black tea extract (Arent et al., 2010) and
918 mate tea (Panza et al. 2016). Nevertheless, although there are some promising findings that green, black and mate tea
919 polyphenols improve some outcomes of EIMD, research in this topic still is relatively weak and definitive conclusions
920 about efficacy cannot yet be drawn because of the variations in study design, intervention protocols, eligibility criteria,
921 statistical power, and timing of the intervention.

922 The exogenous antioxidants, Quercetin (Q), a polyphenol present in grapes, berries, tomatoes, apples, onions as
923 well as in some medicinal plants such as *Hypericum perforatum* and *Gingko biloba* have been demonstrated to exert a
924 variety of bioactive effects that are related to its anti-inflammatory and antioxidant properties (Abbey and Rankin, 2011;
925 Malaguti et al., 2013; Bazzucchi et al., 2019; Askari et al., 2012). In addition, Martin-Rincon et al. (2020) investigated
926 competition recovery in males and females following ingestion quercetin (140 mg) in combination with mango leaf extract
927 that was rich in the polyphenol mangiferin (140 mg) one hour before exercise, followed by three additional doses every
928 eight hours and reported an attenuation of muscle pain and a faster recovery of muscle performance. Conversely, other
929 lines of enquiry failed to support the use of quercetin in managing EIMD in trained cyclists (Nieman et al. 2007a;
930 McAnulty et al. 2008), ultramarathoners (Nieman et al., 2007b), healthy men and women (O’Fallon et al. 2012), young
931 men (Patrizio et al. 2018) and male and female runners (Konrad et al. 2011). In summary, quercetin requires further
932 research to clarify the most appropriate dose, timing and implication for longer term supplementation. Astaxanthin is also
933 a powerful antioxidant naturally found in algae, fish and birds, has been shown to be effective against lipid peroxidation
934 and oxidative stress *in vivo* and *in vitro* models. Some authors suggest that astaxanthin has small beneficial effect on
935 managing EIMD (Djordjevic et al. 2012), conversely other studies (Bloomer et al. 2005; Klinkenberg 2013) demonstrated
936 no benefit on muscle function nor antioxidant capacity.

937 In relation to EIMD, there is a growing interest from scientists and practitioners in the use of foods rich in
938 phytochemicals, for example black chokeberry (Kim et al., 2019), acai (Carvalho-Peixoto et al., 2015; Sadowska-Krępa
939 et al., 2015), grape (Kim and So, 2019; Dalla Corte et al., 2013), trans-resveratrol and polyphenol-enriched extracts (Jo
940 et al., 2019) and cacao (Morgan et al., 2018; Peschek et al., 2013) but to date, there is very limited evidence to support
941 efficacy on EIMD outcomes. Lyall et al. (2009) examined the consumption of a New Zealand Blackcurrant extract
942 (equivalent to 48 g of blackcurrants) in male and female rowers and showed a blunted CK response at 24 h post-exercise
943 and suppressed plasma protein carbonyls levels (0.9 ± 0.1 vs. 0.6 ± 0.1 nmol/mg) compared to a placebo. Later, Hutchison
944 et al. (2016) supported these findings and confirmed that consumption of a drink of blackcurrant nectar (16 oz, twice a
945 day), reduced CK levels at 48 and 96 h after a session of eccentric knee extensions in college students, compared to
946 placebo. Similarly, Coelho et al. (2017) reported that females who completed an maximum eccentric exercise of the
947 bicep brachii following consumption of New Zealand blackcurrant extract (7 days before and 4 days following), reported
948 less muscle soreness at 24 and 48 h post exercise, compared to a placebo, but no differences between groups in CK efflux
949 or IL-6. The preliminary data from these studies show that blackcurrants might attenuate indices of EIMD, although the
950 mechanisms have not been clarified. Like blackcurrants, blueberries and bilberries are also known for the anthocyanin
951 and other phenolic compounds (Wu et al. 2004) that could be important modulators of oxidative stress and inflammation
952 (McAnulty et al., 2004; Hurst et al., 2010 McLeay et al. 2012) in an EIMD scenario.

953 Tomato juice, rich in lycopene, vitamin C and other vitamins and minerals, could represents a protective food
954 against oxidative stress and inflammation (Jacob et al. 2008). Previous studies investigated the effects of tomato juice on
955 LDH, CK, C-RP and homocysteine (Tsitsimpikou et al. 2013) and oxidative stress exercise-induced (Harms-Ringdahl et
956 al., 2012; Samaras et al. 2014) showing potential use of tomato juice as a strategy to modulate the negative outcomes of

957 EIMD. While tomato juice consumption has proven effective in ameliorating LDH and CK responses to anaerobic training
958 (Tsitsimpikou et al. 2013) and reducing serum levels of 8-oxo-dG (Harms-Ringdahl et al., 2012), further studies must
959 clarify doses, duration and better explain the mechanism(s) of action.

960 An area of growing interest is diet modulation of gut microbiota and, in particular, probiotics supplementation
961 to influence immune functions (Gleeson et al. 2011), support healthy digestive and prevent respiratory illness or persistent
962 common cold (Mach et al. 2017) in subjects who follow exercise programs. Other studies have shown potential beneficial
963 effects between probiotics use, muscle damage, repair (Jäger 2016a; Jäger 2016b; Gepner 2017; Martarelli et al. 2011),
964 fatigue (Shing et al. 2014) and inflammation (Lamprecht et al. 2012). Although the use of probiotics in athlete populations
965 has attracted interest and is promising, the small number of studies and considerable differences in experimental
966 approaches represents important limitations. There is little doubt that gut health will impact on whole system physiology,
967 but the interaction between alterations in gut health and microbial populations and exercise performance, recovery and
968 adaptation requires a great deal of work (Francavilla et al., 2017).

969 A growing interest arise for supplements containing collagen-specific peptides, or gelatine (partially hydrolysed
970 collagen), due to its potential role on extra cellular matrix collagen synthesis supporting the remodelling process (Shaw
971 et al. 2017; Li and Wu, 2018), on reduction of muscle soreness (Clifford et al. (2019) and on attenuation of CK activity
972 following muscle-damaging exercise (Lopez et al. 2014). A recently published study showed that ingestion of 5 or 15 g
973 of gelatine augmented bone collagen synthesis following acute mechanical loading (Shaw et al., 2017). As a result, it is
974 tenable that collagen hydrolysate supplementation could attenuate some of signs and symptoms associated with EIMD.

975 Other nutritional strategies to improve recovery from exercise-induced muscle damage include
976 methylsulfonylmethane (Barmaki et al. 2012; van der Merwe and Bloomer, 2016) selenium (Miliadis et al. 2006), allicin
977 (Su et al. 2008), lemon verbena extract (Buchwald-Werner et al., 2018), flavanol-rich lychee (Nishizawa et al. 2011)
978 saffron (Meamarbashi and Rajabi 2015), sesame (Barbosa et al. 2017), spinach (Bohlooli et al. 2015) melatonin (Cheikh
979 et al. 2020) and sucrosomial® iron (Bongiovanni et al. 2019) specialized water (Borsa et al. 2013) and the leucine
980 metabolite alfa-hydroxy-isocaproic acid (HICA) studied by Mero et al. (2010) and Teixeira et al. (2019) for its proposed
981 ability to reduce training-induced inflammation and DOMS.

982 The aforementioned studies have tested the consumption of foods and supplements to reduce exercise-induced
983 outcomes, but other options, as such caloric restriction and/or intermittent fasting, to modulate inflammation and oxidative
984 stress was object of investigations (Dannecker et al. 2013; Pons et al. 2018). Fasting and caloric restriction are intriguing
985 for their potential effects on markers of inflammation and oxidative status (Golbidi et al., 2017) but could they represent
986 a possible application to modulate signs and symptoms of EIMD, particularly in athletic populations? Future studies
987 should attempt to explore this fascinating topic.

988

989 **Limitations, practical applications and future perspectives**

990 Importantly, all supplements that contain antioxidants might be of limited benefit where the athlete's
991 diet has a rich variety of fruits, vegetables, whole grains and nuts. A "food first" approach is highly recommended (Close
992 et al., 2019), rather than processed capsules. Many foods contain sufficient phytochemicals to support the stressors
993 associated with training and competition and aid recovery (Peternelj and Coombes 2011; Myburgh 2014). Exercise-
994 induced muscle damage inflammation and oxidative stress are important elements in the activation of appropriate adaptive
995 pathways that underpin repair of structures and exercise-induced improvements in health and immune function. Indeed,
996 a growing body of evidence indicates that RONS production are a necessary response for exercise adaptations. Therefore
997 supplements (e.g. antioxidant, anti-inflammatory) that completely mitigate or stop oxidative stress and inflammation

998 could alter these adaptive events. It is only when the exercise stressors might outweigh the benefits that whole foods
999 can afford that these supplements could be used to facilitate the recovery process. Critical to this mantra, is the need to
1000 assess the balance between the benefits these supplements might provide against the potential to interfere with
1001 exercise-induced adaptations. Consequently, practitioners should consider a periodised approach where identification of
1002 higher-stress training and competition needs to be managed to facilitate timely recovery. Equally researchers should
1003 attempt to investigate the longer-term effect of the use interventions on the adaptive response and hence the ability to
1004 interfere or facilitate long term athlete development. Notwithstanding, there is currently no evidence to suggest that any
1005 functional food interventions (e.g., fruits, berries, cherries) affect the adaptive response. However, until this is
1006 systematically examined, the aforementioned idea of hormesis should be pragmatically applied to use interventions at an
1007 appropriate time (Figure 2).

1008 From an applied perspective, how might this be applied? For example, in a preseason training phase, the best
1009 nutritional advice might be to follow a regular and varied diet and allow adaptations to occur naturally without the addition
1010 of supplements. However, when recovery is the priority, then the use of carefully planned interventions can be very
1011 helpful to restore function and abate soreness. One scenario could be during congested competition phases and busy travel
1012 schedules, such as those experienced in professional soccer, ice hockey, rugby, basketball and so on, where the main goal
1013 is to restore athletes back to the basal state as quickly as possible.

1014 In the most-part, a majority of studies use heterogenous populations and variables, where fitness level, age,
1015 gender, sample size, outcome measures, short or longer term period of supplementation and study designs. For example,
1016 an important, but nonetheless often-overlooked limitation of some studies, was the use of a cross-over design in a
1017 damaging paradigm, which has been shown to confer a contralateral repeated bout effect and hence influence the damage
1018 in the contralateral limb (Howatson and van Someren, 2007). In addition, measures of REDOX balance at the systemic
1019 level often reveal limited mechanistic insights on the real activity of polyphenols with regard to oxidative stress and
1020 defence. Most measures reported in the exercise literature are fraught with limitations and should be treated cautiously to
1021 infer what might be occurring at the skeletal muscle level (Cobley et al. 2017).

1022 Furthermore, other important issues to consider include bioavailability and pharmacokinetics: is the dose enough,
1023 too much or too little? Is it taken for long enough? When do levels peak in the body? Are the bioactive compounds stored
1024 or metabolised quickly? Should it be before or after bout of exercise? With this in mind, it is difficult to univocally
1025 translate science into practice. Overall, we classified supplements in three main groups, based on the current level of
1026 evidence (Figure 3) and the most frequently studied micronutrients to support EIMD, so called group “good level of
1027 evidence”, are summarized in Table 1. Importantly, this group of supplements are suggested to have a good deal research
1028 completed on them as opposed to the supplement being unequivocally effective. Despite this, there is still much to be
1029 learned regarding these interventions and how they might relate to EIMD, recovery and remodelling.

1030 It is important that further studies be conducted, in male and female recreationally active and elite athletes. Other
1031 aspect that must be taken in account is that placebo-controlled, randomized control studies are difficult to design and
1032 conduct in elite athletes. Therefore, the importance of high-quality case studies can become increasing vital to inform the
1033 evidence base. It is also important that future research replicates how athletes actually utilize supplements and daily
1034 dietary programs in a real-world scenario in order to inform practice. Finally, there is a growing need to ensure that
1035 professional sports nutritionists receive the right information and education on what to use and when to use it, to correctly
1036 translate the evidence into practice.

1037
1038 **Conclusion**

1039 This review provided an overview of EIMD and explored the nutritional strategies that could manage the
1040 negative signs and symptoms associated with EIMD. Very often there is little or no evidence-based practice in applying
1041 these nutritional interventions and hence it is difficult to ascertain an appropriate strategy to combat the negative effects
1042 of EIMD. Further research is warranted to elucidate the most appropriate dose, frequency and intensity of these
1043 interventions in order to determine “best evidence-based practice”. Importantly, it is unclear how these interventions
1044 affect the adaptation process and it might transpire that treatments are being prescribed that attenuate adaptations;
1045 therefore, there is a need to ascertain the full implication of these interventions; 1) to reduce the negative effects of EIMD,
1046 and 2) understand the impact of these interventions on adaptation. With this in mind, a periodized approach to nutritional
1047 supplementation seems a sensible approach to adopt until further evidence is available, particularly for well-trained
1048 athletes.
1049
1050





Fig. 1 Proposed mechanisms of Exercise-induced muscle damage

Fig. 1

1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064

Proposed mechanisms of Exercise-induced muscle damage

The potential damage model can be summarized into two general phases: an initial or *primary phase* (left-hand side), triggered by eccentric contractions, beyond this, an increased permeability of the muscle cellular membrane, causing extracellular influx of Ca^{2+} into the muscle fibre activates different Ca^{2+} -sensitive proteases (calpains). Calpain activation leads to proteolysis of cytoskeletal and costameric proteins. Furthermore, a failure of excitation-contraction coupling also seems to play an important role in strength loss following strenuous exercise. The *secondary phase* (right-hand side) is characterized by neutrophils infiltrate to damaged muscle fibres and production of reactive oxygen and nitrogen species to degrade cellular debris. Neutrophils are substituted by macrophages M1 and in the latter stage of muscle damage, a shift from M1 to M2 macrophages is associated with the activation of satellite cells and the subsequent regeneration of muscle fibres. Neutrophils and macrophages also express tumour necrosis factor (TNF), which activates the ubiquitin-proteasome pathway that regulates proteolysis.



Increase in inflammation and oxidative stress for the adaptive remodeling process is a positive effect of exercise stress


Conceptual region to intervene with nutritional recovery strategy:
 1) when exercise stress is high enough to be detrimental to adaptation
 2) when exercise stress impairs timely return to training & competition and recovery is prioritized over adaptation

Dose of exercise stress

Impair of adaptation is a negative effect of exercise stress

1065
 1066
 1067
 1068
 1069
 1070
 1071
 1072
 1073

Fig. 2 Model of hormesis in exercise recovery. Figure adapted from Howatson et al. (2016)
 Hormesis theory in the context of nutritional interventions for the management of EIMD. This framework suggests that the adaptive response to EIMD presents as a bell-shaped curve. A positive effect of the exercise stress exists to a point when the exposure becomes too great, thereafter there is an impaired adaptive response. Using this theory, we suggest a conceptual region for intervention (yellow text box) where the exercise stress impairs timely return to training and competition or is detrimental to adaptation.



1074
 1075
 1076
 1077

Fig. 3 Proposed strategies for reducing signs and symptoms of Exercise-induced muscle damage
 Overview of nutritional strategies proposed for manage EIMD outcomes and classified based on level of evidence.

1078 **REFERENCES**

- 1079 Abbey EL, Rankin JW (2011) Effect of quercetin supplementation on repeated-sprint performance, xanthine oxidase
1080 activity, and inflammation. *Int J Sport Nutr Exerc Metab.* 21:91-6.
- 1081 Abbott W, Brashill C, Brett A, Clifford T (2019) Tart cherry juice: no effect on muscle function loss or muscle soreness
1082 in professional soccer players after a match. *Int J Sports Phys. Perform.* 1-6.
- 1083 Abbott W, Brett A, Cockburn E, Clifford T (2019) Presleep casein protein ingestion: acceleration of functional recovery
1084 in professional soccer players. *Int J Sports Physiol Perform.* 14:385-391.
- 1085 Anand P, Kunnumakkara AB, Newman RA, Aggarwal BB (2007) Bioavailability of curcumin: problems and promises.
1086 *Mol Pharm.* 4:807-18.
- 1087 Amalraj A, Divya C, Gopi S (2020) The effects of bioavailable curcumin (Cureit) on delayed-onset muscle soreness
1088 induced by eccentric continuous exercise: a randomized, placebo-controlled, double-blind clinical study. *J Med Food.*
1089 doi: 10.1089/jmf.2019.4533 [Epub ahead of print].
- 1090 Amalraj A, Jude S, Varma K, Jacob J, Gopi S, Oluwafemi OS, Thomas S (2017) Preparation of a novel bioavailable
1091 curcuminoid formulation (Cureit™) using polar-nonpolar-sandwich (PNS) technology and its characterization and
1092 applications. *Mater Sci Eng C Mater Bio Appl.* 75:359-367.
- 1093 Ammar A, Turki M, Chtourou H, Hammaouda O, Trabelsi K, Kallel C, Abdelkarim O, Hoekelmann A, Bouaziz M, Ayadi
1094 F, Driss T, Souissi N (2016) Pomegranate supplementation accelerates recovery of muscle damage and soreness and
1095 inflammatory markers after a weightlifting training session. *PLoS One* 11:e0160305.
- 1096 Arazi H, Asadi A, Suzuki K (2018b) The effects of Beta-hydroxy Beta-methylbutyrate-free acid supplementation and
1097 resistance training on oxidative stress markers: a randomized, double-blind, placebo-controlled study. *Antioxidants*
1098 (Basel). 7(6).
- 1099 Arazi H, Hosseini Z, Asadi A, Ramirez-Campillo R, Suzuki K (2019) β -hydroxy β -methylbutyrate-free acid attenuates
1100 oxidative stress induced by a single bout of plyometric exercise. *Front Physiol.* 10:776.
- 1101 Arazi H, Taati B, Suzuki K (2018a) A review of the effects of leucine metabolite (β -hydroxy β -methylbutyrate)
1102 supplementation and resistance training on inflammatory markers: a new approach to oxidative stress and cardiovascular
1103 risk factors. *Antioxidants (Basel).* 7(10).
- 1104 Arent SM, Senso M, Golem DL, McKeever KH (2010) The effect of theaflavin-enriched black tea extract on muscle
1105 soreness, oxidative stress, inflammation, and endocrine responses to acute anaerobic interval training: a randomized,
1106 double-blind, crossover study. *J Int Soc Sports Nutr.* 7:11.
- 1107 Armstrong RB (1984) Mechanisms of exercise-induced delayed onset muscular soreness: a brief review. *Med Sci*
1108 *Sports Exerc.* 16:529-38.

1109 Arnold L, Henry A, Poron F, Baba-Amer Y, van Rooijen N, Plonquet A, Gherardi RK, Chazaud B (2007) Inflammatory
1110 monocytes recruited after skeletal muscle injury switch into inflammatory macrophages to support myogenesis. *J Exp*
1111 *Med.* 204:1057-69.

1112 Asadi A, Arazi H, Suzuki K (2017) Effects of β -hydroxy β -methylbutyrate-free acid supplementation on strength, power
1113 and hormonal adaptations following resistance training. *Nutrients.* 9(12).

1114 Asai A, Miyazawa T (2000) Occurrence of orally administered curcuminoid as glucuronide and glucuronide/sulphate
1115 conjugates in rat plasma. *Life Sci.* 67:2785-93.

1116 Askari G, Ghiasvand R, Karimian J, Feizi A, Paknahad Z, Sharifirad G, Hajishafiei M (2012) Does quercetin and vitamin
1117 C improve exercise performance, muscle damage, and body composition in male athletes? *J Res Med Sci.* 17:328-31.

1118 Bach HV, Kim J, Myung SK, Cho YA (2016) Efficacy of ginseng supplements on fatigue and physical performance: a
1119 meta-analysis. *J Korean Med Sci.* 31:1879-1886.

1120 Baratloo A, Rouhipour A, Forouzanfar MM, Safari S, Amiri M, Negida A (2016) The role of caffeine in pain
1121 management: a brief review. *Anesth Pain Med.* 6:e33193.

1122 Barbosa CV, Silva AS, de Oliveira CV, Massa NM, de Sousa YR, da Costa WK, Silva AC, Delatorre P, Carvalho R,
1123 Braga VA, Magnani M (2017) Effects of Sesame (*Sesamum indicum L.*) supplementation on creatine kinase, lactate
1124 dehydrogenase, oxidative stress markers, and aerobic capacity in semi-professional soccer players. *Front Physiol.* 8:196.

1125 Barker T, Henriksen VT, Martins TB, Hill HR, Kjeldsberg CR, Scheneider ED, Dixon BM, Weaver LK (2013a) Higher
1126 serum 25-hydroxyvitamin D concentrations associate with a faster recovery of skeletal muscle strength after muscular
1127 injury. *Nutrients.* 5:1253-75.

1128 Barker T, Scheneider ED, Dixon BM, Henriksen VT, Weaver LK (2013b) Supplemental vitamin D enhances the
1129 recovery in peak isometric force shortly after intense exercise. *Nutr Metab.* 10:69.

1130 Barmaki S, Bohlooli S, Khoshkharesh F, Nakhostin-Roohi B (2012) Effect of methylsulfonylmethane supplementation
1131 on exercise - Induced muscle damage and total antioxidant capacity. *J Sports med Phys Fitness.* 52:170-4.

1132 Basham SA, Waldman HS, Krings BM, Lamberth J, Smith JW, McAllister MJ (2019) Effect of curcumin supplementation
1133 on exercise-induced oxidative stress, inflammation, muscle damage, and muscle soreness. *J Diet Suppl.* 26:1-14.

1134 Bassini-Cameron A, Sweet E, Bottino A, Bittar C, Veiga C, Cameron LC (2007) Effect of caffeine supplementation on
1135 haematological and biochemical variables in elite soccer players under physical stress conditions. *Br J Sports Med.*
1136 41:523-30.

1137 Bassit RA, Curi R, Costa Rosa LF (2008) Creatine supplementation reduces plasma levels of pro-inflammatory cytokines
1138 and PGE2 after a hal-ironman competition. *Amino Acids.* 35:425-31.

1139 Bassit RA, Pinheiro CH, Vitzel KF, Sproesser AJ, Silveira LR, Curi R (2010) Effect of short-term creatine
1140 supplementation on markers of skeletal muscle damage after strenuous contractile activity. *Eur J Appl Physiol.* 108:945-
1141 55.

1142 Baumert P, Lake MJ, Stewart CE, Drust B and Erskine RM (2016) Genetic variation and exercise-induced muscle
1143 damage: implications for athletic performance, injury and ageing. *Eur J Appl Physiol.* 116:1595-1625.

1144 Bazzucchi I, Patrizio F, Ceci R, Duranti G, Sgrò P, Sabatini S, Di Luigi L, Sacchetti M, Felici F (2019) The effects of
1145 quercetin supplementation on eccentric exercise-induced muscle damage. *Nutrients.* 11 (1). pii:E205.

1146 Beals K, Allison KF, Darnell M, Lovalekar M, Baker R, Nieman DC, Vodovotz Y, Lephart SM (2017) The effects of a
1147 tart cherry beverage on reducing exercise-induced muscle soreness. *Isokinetics and Exercise Science* 25:53-63.

1148 Beaton LJ, Tarnopolsky MA, Phillips SM (2002) Contraction-induced muscle damage in humans following calcium
1149 channel blocker administration. *J Physiol.* 544:849-859.

1150 Beck TW, Housh TJ, Johnson GO, Schimidt RJ, Housh DJ, Coburn JW, Malek MH, Mielke M (2007) Effect of a protease
1151 supplement on eccentric exercise-induced markers of delayed-onset muscle soreness and muscle damage. *J Strength Cond*
1152 *Res.* 21:661-7.

1153 Beelen M, Burke LM, Gibala MJ, van Loon LJC (2010) Nutritional strategies to promote postexercise recovery. *Int J*
1154 *Sport Nutr Exerc Metab.* 20:515-32.

1155 Belcastro AN (1993) Skeletal muscle calcium-activated neutral protease (calpain) with exercise. *J Appl Physiol* 74:1381-
1156 6.

1157 Belcastro AN, Shewchuk LD, Raj DA (1998). Exercise-induced muscle injury: a calpain hypothesis. *Mol Cell Biochem.*
1158 179:135-45.

1159 Bell PG, McHugh MP, Stevenson E, Howatson G (2014) The role of cherries in exercise and health. *Scand J Med Sci*
1160 *Sports* 24:477-90.

1161 Bell PG, Stevenson E, Davison GW, Howatson G (2016) The effects of Montmorency tart cherry concentrate
1162 supplementation on recovery following prolonged, intermittent exercise. *Nutrients.* 8(7).

1163 Bell PG, Walshe IH, Davison GW, Stevenson E, Howatson G (2014) Montmorency cherries reduce the oxidative stress
1164 and inflammatory responses to repeated days high-intensity stochastic cycling. *Nutrients.* 6:829-43.

1165 Bell PG, Walshe IH, Davison GW, Stevenson EJ, Howatson G (2015) Recovery facilitation with montmorency cherries
1166 following high-intensity, metabolically challenging exercise. *Appl Physiol Nutr Metab* 40:414-23.

1167 Bieuzen F, Bleakley CM, Costello JT (2013) Contrast water therapy and exercise induced muscle damage: a systematic
1168 review and meta-analysis. *PLoS One.* 8:e62356.

1169 Borsa PA, Kaiser KL, Martin JS (2013) Oral consumption of electrokinetically modified water attenuates muscle damage
1170 and improves postexercise recovery. *J Appl Physiol.* 114:1736-42.

1171 Black CD, Herring MP, Hurley DJ, O'Connor PJ (2010) Ginger (*Zingiber officinale*) reduces muscle pain caused by
1172 eccentric exercise. *J Pain.* 11:894-903.

1173 Black CD, O'Connor PJ (2010) Acute effects of dietary ginger on muscle pain induced by eccentric exercise. *Phyther*
1174 *Res.* 24:1620-6.

1175 Black CD, Oconnor PJ (2008) Acute effects of dietary ginger on quadriceps muscle pain during moderate-intensity
1176 cycling exercise. *Int J Sport Nutr Exerc Metab.* 18:653-64.

1177 Black KE, Witard OC, Baker D, Healey P, Lewis V, Tavares F, Christensen S, Pease T, Smith B (2018) Adding omega-
1178 3 fatty acids to a protein-based supplement during pre-season training results in reduced muscle soreness and the better
1179 maintenance of explosive power in professional Rugby Union players. *Eur J Sport Sci.* 18:1357-1367.

1180 Blacker SD, Williams NC, Fallowfield JL, Bilzon JLj, Willems ME (2010) Carbohydrate vs protein supplementation for
1181 recovery of neuromuscular function following prolonged moad carriage. *J Int Soc Sports Nutr.* 7:2.

1182 Bleakley CM, Bieuzen F, Davison GW, Costello JT (2014) Whole-body cryotherapy: empirical evidence and theoretical
1183 perspectives. *Open Access J Sports Med.* 5:25-36.

1184 Bloomer RJ, Fry A, Schilling B, Chiu L, Hori N, Weiss L (2005) Astaxanthin supplementation does not attenuate muscle
1185 injury following eccentric exercise in resistance-trained men. *Int J Sport Nutr Exerc Metab.* 15:401-12.

1186 Bloomer RJ, Larson DE, Fisher-Wellman KH, Galpin AJ, Schilling BK (2009) Effect of eicosapentaenoic and
1187 docosahexaenoic acid on resting and exercise-induced inflammatory and oxidative stress biomarkers: a randomized,
1188 placebo controlled, cross-over study. *Lipids Health Dis.* 8:36.

1189 Bohlooli S, Barmaki S, Khoshkharesh F, Nakhostin-Roohi B (2015) The effect of spinach supplementation on exercise-
1190 induced oxidative stress. *J Sports Med Phys Fitness.* 55:609-14.

1191 Bolling BW, Chen CY, McKay DL, Blumberg JB (2011) Tree nut phytochemicals: composition, antioxidant capacity,
1192 bioactivity, impact factors. A systematic review of almonds, brazils, cashews, hazelnuts, macadamias, pecans, pine nuts,
1193 pistachios and walnuts. *Nutr Res Rev.* 24:244-75.

1194 Bongiovanni T, Pasta G, Tarantino G (2019) Sucrosomial® iron and folic acid supplementation is able to induce IL-6
1195 levels variation in healthy trained professional athletes, regardless of the hemoglobin and iron values. *Science & Sport.*
1196 34:165-172.

1197 Bowtell JL, Summers DP, Dyer A, Fox P, Mileva KN (2011) Montmorency cherry juice reduces muscle damage caused
1198 by intensive strength exercise. *Med Sci Sports Exerc.* 43:1544-51.

1199 Brown MA, Stevenson EJ, Howatson G (2018) Whey protein hydrolysate supplementation accelerates recovery from
1200 exercise-induced muscle damage in females. *Appl Physiol Nutr Metab.* 43:324-330.

1201 Brown MA, Stevenson EJ, Howatson G (2019) Montmorency tart cherry (*Prunus cerasus* L.) supplementation accelerates
1202 recovery from exercise-induced muscle damage in females. *Eur J Sport Sci.* 19:95-102.

1203 Buchwald-Werner S, Naka I, Wilhelm M, Schütz E, Schoen C, Reule C (2018) Effects of lemon verbena extract
1204 (Recoverben®) supplementation on muscle strength and recovery after exhaustive exercise: a randomized, placebo-
1205 controlled trial. *J Int Soc Sports Nutr.* 15:5.

1206 Buckley JD, Thomson RL, Coates AM, Howe PR, DeNichilo MO, Rowney MK (2010) Supplementation with a whey
1207 protein hydrolysate enhances recovery of muscle force-generating capacity following eccentric exercise. *J Sci Med Sport*
1208 13:178-81.

1209 Buford TW, Cooke MB, Redd LL, Hudson GM, Shelmadine BD, Willoughby DS (2009) Protease supplementation
1210 improves muscle function after eccentric exercise. *Med Sci Sports Exerc.* 41:1908-14.

1211 Burke LM (2008) Caffeine and sports performance. *Appl Physiol Nutr Metab.* 33:1319-34.

1212 Byrne C, Twist C, Eston R (2004) Neuromuscular function after exercise-induced muscle damage: theoretical and applied
1213 implications. *Sports Med* 34:49-69.

1214 Caldwell AR, Tucker MA, Butts CL, McDermott BP, Vingren JL, Kunces LJ, Lee EC, Munoz CX, Williamson KH,
1215 Armstrong LE, Ganio MS (2017) Effect of caffeine on perceived soreness and functionality following an endurance
1216 cycling event. *J Strength Cond Res.* 31:638-643.

1217 Caldwell LK, DuPont WH, Beeler MK, Post EM, Barnhart EC, Hardesty VH, Anders JP, Borden EC, Volek JS, Kraemer
1218 WJ (2018) The effects of a Korean Ginseng, GINST15, on perceptual effort, psychomotor performance, and physical
1219 performance in men and women. *J Sports Sci Med.* 17:92-100.

1220 Candia-Luján R, De Paz Fernández JA, Costa Moreira O (2014) Are antioxidant supplements effective in reducing
1221 delayed onset muscle soreness? A systematic review. *Nutr. Hosp* 31:32-45.

1222 Carvalho-Peixoto J, Moura MR, Cunha FA, Lollo PC, Monteiro WD, Carvalho LM, Farinatti Pde T (2015) Consumption
1223 of açai (*Euterpe oleracea* Mart.) functional beverage reduces muscle stress and improve effort tolerance in elite athletes:
1224 a randomized controlled intervention study. *Appl Physiol Nutr Metab.* 40:725-33.

1225 Castell JV, Friedrich G, Kuhn CS, Poppe GE (1997) Intestinal absorption of undegraded proteins in men: presence of
1226 bromelain in plasma after oral intake. *Am J Physiol.* 273:G139-46.

1227 Chappell AJ, Allwood DM, Johns R, Brown S, Sultana K, Anand A, Simper T (2018) Citrulline malate supplementation
1228 does not improve german volume training performance or reduce muscle soreness in moderately trained males and
1229 females. *J Int Soc Sports Nutr.* 15:42.

1230 Charoengam N, Shirvani A, Holick MF (2019) Vitamin D for skeletal and non-skeletal health: what we should know.
1231 J Clin Orthop Trauma. 10:1082-1093.

1232 Chazaud B (2016) Inflammation during skeletal muscle regeneration and tissue remodeling: application to exercise-
1233 induced muscle damage management. Immunol Cell Biol. 94:140-5.

1234 Chazaud B, Sonnet C, Lafuste P, Bassez G, Rimaniol AC, Authier FJ, Dreyfus PA, Gherardi RK (2003) Satellite cells
1235 attract monocytes and use macrophages as a support to escape apoptosis and enhance muscle growth. J Cell Biol.
1236 163:1133-43.

1237 Cheikh M, Makhoulouf K, Ghattassi K, Graja A, Ferchichi S, Kallel C, Houada M, Souissi N, Hammouda O (2020)
1238 Chronobiol Int. 37:236-247.

1239 Chen TC, Chen HL, Pearce AJ, Nosaka K (2012) Attenuation of eccentric exercise-induced muscle damage by
1240 preconditioning exercises. Med Sci Sports Exerc 44:2090-2098.

1241 Chen HY, Chen YC, Tung K, Chao HH, Wang HS (2019) Effects of caffeine and sex on muscle performance and delayed-
1242 onset muscle soreness after exercise-induced muscle damage: a double-blind randomized trial. J Appl Physiol. 27:798-
1243 805.

1244 Childs A, Jacobs C, Kaminski T, Halliwell B, Leeuwenburgh C (2001) Supplementation with vitamin C and N-acetyl-
1245 cysteine increases oxidative stress in humans after an acute muscle injury induced by eccentric exercise. Free Radic Biol
1246 Med. 31:745-753.

1247 Clarkson PM, Hubal MJ (2002) Exercise-induced muscle damage in humans. Am J Phys Med Rehabil. 81:52-69.

1248 Clarkson PM, Sayers SP (1999) Etiology of exercise-induced muscle damage. Can J Appl Physiol. 24:234-48.

1249 Clifford T, Allerton DM, Brown MA, Harper L, Horsburgh S, Keane KM, Stevenson EJ, Howatson G (2017c) Minimal
1250 muscle damage after a marathon and no influence of beetroot juice on inflammation and recovery. Appl Physiol Nutr
1251 Metab. 42:263-270.

1252 Clifford T, Bell O, West DJ, Howatson G, Stevenson EJ (2016a) The effects of beetroot juice supplementation on indices
1253 of muscle damage following eccentric exercise. Eur J Appl Physiol 116:353-62.

1254 Clifford T, Bell O, West DJ, Howatson G, Stevenson EJ (2017a) Antioxidant-rich beetroot juice does not adversely affect
1255 acute neuromuscular adaptation following eccentric exercise. J Sports Sci. 35:812-819.

1256 Clifford T, Berntzen B, Davison GW, West DJ, Howatson G, Stevenson EJ (2016b) Effects of beetroot juice on recovery
1257 of muscle function and performance between bouts of repeated sprint exercise. Nutrients. 8(8).

1258 Clifford T, Constantinou CM, Keane KM, West DJ, Howatson G, Stevenson EJ (2017d) The plasma bioavailability of
1259 nitrate and betain from beta vulgaris rubra in humans. Eur J Nutr. 56:1245-1254.

1260 Clifford T, Howatson G, West DJ, Stevenson EJ (2015) The potential benefits of red beetroot supplementation in health
1261 and disease. *Nutrients*. 7:2801-22.

1262 Clifford T, Howatson G, West DJ, Stevenson EJ (2017b) Beetroot juice is more beneficial than sodium nitrate for
1263 attenuating muscle pain after strenuous eccentric-bias exercise. *Appl Physiol Nutr Metab*. 42:1185-1191.

1264 Clifford T, Ventress M, Allerton DM, Stansfield S, Tang JCY, Fraser WD, Vahnoecke B, Prawitt J, Stevenson E (2019)
1265 The effects of collagen peptides on muscle damage, inflammation and bone turnover following exercise: a randomized,
1266 controlled trial. *Amino Acids*, 51:691-704.

1267 Close LC, Kasper AM, Morton JP (2019) From paper to podium: quantifying the translational potential of performance
1268 nutrition research. *Sports Med*. 49:25-37.

1269 Cobley JN, Close GL, Bailey DM, Davison GW (2017) Exercise redox biochemistry: Conceptual, methodological and
1270 technical recommendations. *Redox Biology*. 12:540-548.

1271 Cockburn E, Hayes PR, French DN, Stevenson E, Gibson ASC (2008) Acute milk-based protein-CHO supplementation
1272 attenuates exercise-induced muscle damage. *Appl Physiol Nutr Metab*. 33:775-83.

1273 Cockburn E, Bell PG, Stevenson E (2013) Effect of milk on team sport performance after exercise-induced muscle
1274 damage. *Med Sci Sports exerc*. 45:1585-92.

1275 Cockburn E, Robson-Ansley P, Hayes PR, Stevenson E (2012) Effect of volume of milk consumed on the attenuation of
1276 exercise-induced muscle damage. *Eur J Appl Physiol* 112:3187-94.

1277 Cockburn E, Stevenson E, Hayes PR, Robson-Ansley P, Howatson G (2010) Effect of milk-based carbohydrate-protein
1278 supplement timing on the attenuation of exercise-induced muscle damage. *Appl Physiol Nutr Metab*. 35:270-7.

1279 Coelho M, Buxton S, Butcher R, Foran D, Manders R, Hunt J (2017) The effect of New Zealand blackcurrant extract on
1280 recovery following eccentric exercise induced muscle damage. *Proceedings of the nutrition society*. 76(OCE2), E34.

1281 Coelho RL, Oliveira AC, Prestes J, Sérgio Denadai B (2015) Consumption of cherries as a strategy to attenuate exercise-
1282 induced muscle damage and inflammation in humans. *Nutr Hosp*. 32:1885-93.

1283 Connolly DA, Lauzon C, Agnew J, Dunn M, Reed B (2006) The effects of vitamin C supplementation on symptoms of
1284 delayed muscle soreness. *J Sports Med Phys Fitness*. 46:462-7.

1285 Connolly DA, McHugh MP, Padilla-Zakour OI, Carlson L, Sayers SP (2006) Efficacy of a tart cherry juice blend in
1286 preventing the symptoms of muscle damage. *Br J Sports Med*. 40:679-83.

1287 Cooke MB, Rybalka E, Williams AD, Cribb PJ, Hayes A (2009) Creatine supplementation enhance muscle force recovery
1288 after eccentrically-induced muscle damage in healthy individuals. *J Int Soc Sports Nutr*. 6:13.

1289 Corder KE, Newsham KR, McDaniel JL, Ezekiel UR, Weiss EP (2016) Effects of short-term docosahexaenoic acid
1290 supplementation on markers of inflammation after eccentric strength exercise in women. *J Sports Sci Med*. 15:176-83.

1291 da Silva DK, Jacinto JL, de Andrade WB, Roveratti MC, Estoche JM, Balvedi MCW, de Oliveira DB, da Silva RA,
1292 Aguiar AF (2017) Citrulline malate does not improve muscle recovery after resistance exercise in untrained young adult
1293 men. *Nutrients*. 9(10).

1294 da Silva LA, Tromm CB, Bom KF, Mariano I, Pozzi B, da Rosa GL, Tuon T, da Luz G, Vuolo F, Petronilho F, Cassiano
1295 W, De Souza CT, Pinho RA (2014) Effects of taurine supplementation following eccentric exercise in young adults. *Appl*
1296 *Physiol Nutr Metab*. 39:101-104.

1297 da Silva W, Machado AS, Souza MA, Mello-Carpes PB, Carpes FP (2018) Effect of green tea extract supplementation
1298 on exercise-induced delayed onset muscle soreness and muscular damage. *Physiol Behav*. 194:77-82.

1299 Daab W, Bouzid MA, Lajri M, Bouchiba M, Saafi MA, Rebai H (2020) Chronic beetroot juice supplementation
1300 accelerates recovery kinetics following simulated match play in soccer players. *J Am Coll Nutr*. 3:1-9.

1301 Dalla Corte CL, de Carvalho NR, Amaral GP, Puntel GO, Silva LF, Retamoso LT, Royes LF, Bresciani GB, da Cruz IB,
1302 Rocha JB, Barrio Lera JP, Soares FA (2013) Antioxidant effect of organic purple grape juice on exhaustive exercise.
1303 *Appl Physiol Nutr Metab*. 38:558-65.

1304 Damas F, Nosaka K, Libardi CA, Chen TC, Ugrinowitsch C (2016) Susceptibility to exercise-induced muscle damage: a
1305 cluster analysis with a large sample. *Int J Sports Med*. 37:633-40.

1306 Danesi F, Ferguson LR (2017) Could pomegranate juice help in the control of inflammatory diseases? *Nutrients*. 9(9).

1307 Dannecker EA, Liu Y, Rector RS, Thomas TR, Sayers SP, Leeuwenburgh C, Ray BK (2013) The effect of fasting on
1308 indicators of muscle damage. *Exp Gerontol*. 48:1101-6.

1309 De Carvalho FG, Galan BSM, Santos PC, Pritchett K, Pfrimer K, Ferrioli E, Papoti M, Marchini JS, de Freitas EC (2017)
1310 Taurine: a potential ergogenic aid for preventing muscle damage and protein catabolism and decreasing oxidative stress
1311 produced by endurance exercise. *Front Physiol* 8:710.

1312 De Luca A, Pierno S, Camerino DC (2015) Taurine: the appeal of a safe amino acid for skeletal muscle disorders. *J Transl*
1313 *Med*. 13:243.

1314 Delecroix B, Abaïdia AE, Leduc C, Dawson B, Dupont G (2017) Curcumin and piperine supplementation and recovery
1315 following exercise induced muscle damage: a randomized controlled trial. *J Sports sci Med*. 16:147-153.

1316 Deminice R, Jordao AA (2012) Creatine supplementation reduces oxidative stress biomarkers after acute exercise in rats.
1317 *Amino Acids*. 43:709-15.

1318 Deminice R, Rosa FT, Franco GS, Jordao AA, de Freitas EC (2013) Effects of creatine supplementation on oxidative
1319 stress and inflammatory markers after repeated-sprint exercise in humans. *Nutrition*. 29:1127-32.

1320 Deng B, Wehling-Henricks M, Villalta SA, Wang Y, Tidball JG (2012) IL-10 triggers changes in macrophage phenotype
1321 that promote muscle growth and regeneration. *J Immunol* 189:3669-80.

1322 Deyhle MR, Gier AM, Evans KC, Eggett DL, Nelson WB, Parcell AC, Hyldahl RD (2015) Skeletal muscle inflammation
1323 following repeated bouts of lengthening contractions in humans. *Front Physiol.* 6:424.

1324 DiLorenzo FM, Drager CJ, Rankin JW (2014) Docosahexaenoic acid affects markers of inflammation and muscle
1325 damage after eccentric exercise. *J Strength Cond Res.* 28:2768-74.

1326 Dimitriou L, Hill A, Jehnali A, Dunbar J, Brouner J, McHugh MP, Howatson G (2015) Influence of a montmorency
1327 cherry juice blend on indices of exercise-induced stress and upper respiratory tract symptoms following marathon running
1328 – a pilot investigation. *J Int Soc Sports Nutr.* 12:22.

1329 Dimitrov VG, Arabadzhiev TI, Dimitrova NA, Dimitrov GV (2012) The spectral changes in EMG during a second bout
1330 eccentric contraction could be due to adaptation in muscle fibres themselves: a stimulation study. *Eur J Appl Physiol.*
1331 112:1399-409.

1332 Djordjevic B, Baralic I, Kotur-Stevuljevic, Stefanovic A, Ivanisevic J, Radivojevic N, Andjelkovic M, Dikic N (2012)
1333 Effect of astaxanthin supplementation on muscle damage and oxidative stress markers in elite young soccer players. *J*
1334 *Sports Med Phys Fitness.* 52:382-92.

1335 Dobashi S, Takeuchi K, Koyama K (2020) Hydrogen-rich water suppresses the reduction in blood total antioxidant
1336 capacity induced by 3 consecutive days of severe exercise in physically active males. *Med Gas Res.* 10:21-26.

1337 Drobnic F, Riera J, Appendino G, Togni S, Franceschi F, Valle X, Pons A, Tur J (2014) Reduction of delayed onset
1338 muscle soreness by a novel curcumin delivery system (Meriva®): a randomised, placebo-controlled trial. *J Int Soc Sports*
1339 *Nutr.* 11:31.

1340 Dunsmore KE, Chen PG, Wong HR (2001) Curcumin, a medicinal herbal compound capable of inducing the heat shock
1341 response. *Crit Care Med.* 29:2199-204.

1342 Dupuy O, Douzi W, Theurot D, Bosquet L, Dougué B (2018) An evidence-based approach for choosing post-exercise
1343 recovery techniques to reduce markers of muscle damage, soreness, fatigue, and inflammation: a systematic review with
1344 meta-analysis. *Front Physiol* 9:403.

1345 Ebbeling CB, Clarkson PM (1989) Exercise-induced muscle damage and adaptation. *Sports Med.* 7:207-34.

1346 Eddens L, Browne S, Stevenson EJ, Sanderson B, van Someren K, Howatson G (2017) The efficacy of protein
1347 supplementation during recovery from muscle-damaging concurrent exercise. *Appl Physiol Nutr Metab.* 42:716-724.

1348 Effects of dietary supplementation with vitamin C and E on muscle function during and after eccentric contractions in
1349 humans. *Eur J Appl Physiol.* 93:196-202.

1350 Fedewa MV, Spencer SO, Williams TD, Becker ZE, Fuqua CA (2019) Effect of branched-chain amino acid
1351 supplementation on muscle soreness following exercise: a meta-analysis. *Int J Vitam Nutr. Res.* 89:348-356.

1352 Fernández-Landa J, Calleja-González J, León-Guereño P, Caballero-García A, Córdova A, Mielgo-Ayuso J (2019) Effect
1353 of combination of creatine monohydrate plus HMB supplementation in sports performance, body composition, markers
1354 of muscle damage and hormone status: a systematic review. *Nutrients*. 11(10).

1355 Fernández-Landa J, Fernández-Lázaro D, Calleja-González J, Caballero-García, Córdova A, León-Guereño P, Mielgo-
1356 Ayuso J (2020) Long-term effect of combination of creatine monohydrate plus β -Hydroxy β -Methylbutyrate (HMB) on
1357 Exercise-Induced Muscle Damage. *Biomolecules*. 10(1):140.

1358 Fernández-Landa J, Fernández-Lázaro D, Calleja-González J, Caballero-García A, Córdova A, León-Guereño P, Mielgo-
1359 Ayuso J (2020) Long-term effect of combination of creatine monohydrate plus β -hydroxy β -methylbutyrate (HMB) on
1360 exercise-induced muscle damage and anabolic/catabolic hormones in elite male endurance athletes. *Biomolecules* 10.

1361 Fernández-Lázaro D, Mielgo-Ayuso J, Seco Calvo J, Córdova Martínez A, Caballero García A, Fernandex-Lazaro CI
1362 (2020). Modulation of exercise-induced muscle damage, inflammation, and oxidative markers by curcumin
1363 supplementation in a physically active population: a systematic review. *Nutrients*. 12(2).

1364 Fitschen PJ, Wilson GJ, Wilson JM, Wilund KR (2013) Efficacy of β -hydroxy β -methylbutyrate supplementation in
1365 elderly and clinical populations. *Nutrition* 29:29-36.

1366 Fouré A, Bendahan D (2017) Is branched-chain amino acids supplementation an efficient nutritional strategy to alleviate
1367 skeletal muscle damage? A systematic review. *Nutrients*. 9(10).

1368 Francavilla VC, Bongiovanni T, Todaro L, Di Pietro V, Francavilla G (2017) Probiotic supplements and athletic
1369 performance: a review of the literature. *Medicina dello Sport*. 70:247-59.

1370 Fridén J, Seger J, Sjöström M, Ekblom B (1983) Adaptive response in human skeletal muscle subjected to prolonged
1371 eccentric training. *Int J Sports Med*. 4:177-83.

1372 Gabrial SGN, Shakib MR, Gabrial GN (2018) Protective role of vitamin C intake on muscle damage in male adolescents
1373 performing strenuous physical activity. *Open Access J Med Sci*. 6:1594-1598.

1374 Gaffey A, Slater H, Porritt K, Campbell JM (2017) The effects of curcuminoids on musculoskeletal pain: a systematic
1375 review. *JBI Database System Rev Implement Rep*. 15:486-516.

1376 Gault ML, Willems ME (2013) Aging, functional capacity and eccentric exercise training. *Aging Dis*. 4:351-63.

1377 Gepner Y, Hoffman JR, Shemesh E, Stout JR, Church DD, Varanoske AN, Zelicha H, Shelef I, Chen Y, Frankel H,
1378 Ostfeld I (2017) Combined effect of *Bacillus coagulans* GBI-30, 6086 and HMB supplementation on muscle integrity
1379 and cytokine response during intense military training. *J Appl Physiol*. 123:11-8.

1380 Giamberardino MA, Dragani L, Valente R, Di Lisa F, Saggini R, Vecchiet L (1996) Effects of prolonged L-carnitine
1381 administration on delayed muscle pain and CK release after eccentric effort. *Int J Sports Med*. 17:320-4.

1382 Gissel H (2005) The role of Ca^{2+} in muscle cell damage. *Ann N Y Acad Sci*. 1066:166-80.

1383 Gilchrist M, Winyard PG, Aizawa K, Anning C, Shore A, Benjamin N (2013) Effect of dietary nitrate on blood pressure,
1384 endothelial function, and insulin sensitivity in type 2 diabetes. *Free Radic Bio Med.* 60:89-97.

1385 Gleeson M, Bishop NC, Oliveira M, Tauler P (2011) Daily probiotic's (*Lactobacillus casei* Shirota) reduction of infection
1386 incidence in athletes. *Int J Sport Nutr Exerc Metab.* 21:55-64.

1387 Golbidi S, Daiber A, Korac B, Li H, Essop MF, Laher I (2017) Health benefits of fasting and caloric restriction. *Curr*
1388 *Diab Rep.* 17:123.

1389 Gonzalez AM, Stout JR, Jajtner AR, Townsend JR, Wells AJ, Beyer KS, Boone CH, Pruna GJ, Mangine GT, Scanlon
1390 TM, Bohner JD, Oliveira LP, Fragala MS (2014) Effects of β -hydroxy β -methylbutyrate free acid and cold water
1391 immersion on post-exercise markers of muscle damage. *Amino Acids.* 46:1501-11.

1392 Göring H (2018) Vitamin D in nature: a product of synthesis and/or degradation of cell membrane components.
1393 *Biochemistry (Mosc).* 83:1350-1357.

1394 Gray P, Chappell A, Jenkinson AM, Thies F, Gray SR (2014) Fish oil supplementation reduces markers of oxidative
1395 stress but not muscle soreness after eccentric exercise. *Int J Sport Nutr Exerc Metab.* 24:206-14.

1396 Guigoz Y (2011) Dietary proteins in humans: basic aspects and consumption in Switzerland. *Int J Vitam Nutr Res.* 81:87-
1397 100.

1398 Harms-Ringdahl M, Jenssen D, Haghdoost S (2012) Tomato juice intake suppressed serum concentration of 8-oxodG
1399 after extensive physical activity. *Nutr J.* 11:29.

1400 Harty PS, Cottet ML, Malloy JK, Kerksick CM (2019) Nutritional and supplementation strategies to prevent and attenuate
1401 exercise-induced muscle damage: a brief review. *Sports Med Open.* 5:1.

1402 Hawley JA, Burke LM, Phillips SM, Spriet LL (2011) Nutritional modulation of training-induced skeletal muscle
1403 adaptations. *J Appl Physiol.* 110:834-45.

1404 Heaton LE, Davis JK, Rawson ES, Nuccio RP, Witard OC, Stein KW, Baar K, Carter JM, Baker LB (2017) Selected in-
1405 season nutritional strategies to enhance recovery for team sport athletes: a practical overview. *Sports Med* 47:2201-
1406 2218.

1407 Hennigar SR, McClung JP, Pasiakos SM (2017) Nutritional interventions and the IL-6 response to exercise. *FASEB J.*
1408 31:3719-3728.

1409 Herrlinger KA, Chirouzes DM, Ceddia MA (2015) Supplementation with a polyphenolic blend improves post-exercise
1410 strength recovery and muscle soreness. *Food Nutr Res.* 59:30034.

1411 Highman B, Altland PD (1963) Effect of exercise and training on serum enzyme and tissue changes in rats. *Am J Physiol.*
1412 205:162-6

1413 Hill J, Howatson G, van Someren K, Leeder J, Pedlar C (2014) Compression garments and recovery from exercise-
1414 induced muscle damage: a meta-analysis. *Br J Sports Med.* 48:1340-6.

1415 Hyldahl RD, Nelson B, Xin L, Welling T, Groscost L, Hubal MJ, Chipkin S, Clarkson PM, Parcell AC (2015)
1416 Extracellular matrix remodeling and its contribution to protective adaptation following lengthening contractions in human
1417 muscle. *FASEB J.* 29:2894–904.

1418 Hody S, Croisier JL, Bury T, Rogister B and Leprince P (2019) Eccentric muscle contractions: risks and benefits. *Front*
1419 *Physiol.* 10:536.

1420 Hoffman JR, Cooper J, Wendell M, Im J, Kang J (2004) Effect of β -hydroxy β -methylbutyrate on power performance
1421 and indices of muscle damage and stress during high-intensity training. *J Strength Cond. Res.* 18:747-52.

1422 Hoffman JR, Gepner Y, Stout JR, Hoffman MW, Ben-Dov D, Funk S, Daimont I, Jajtner AR, Townsend JR, Church DD,
1423 Shelef I, Rosen P, Avital G, Chen Y, Frankel H, Ostfeld I (2016) β -hydroxy β -methylbutyrate attenuates cytokine
1424 response during sustained military training. *Nutr Res.* 36:553-63.

1425 Hoseinzadeh K, Daryanoosh F, Baghdasar PJ, Alizadeh H (2015) Acute effects of ginger extract on biochemical and
1426 functional symptoms of delayed onset muscle soreness. *Med J Islam Repub Iran* 29:261.

1427 Hosseinzadeh M, Andersen OK, Arendt-Nielsen L, Samani A, Kamavuako EN, Madeleine P (2015) Adaptation of local
1428 muscle blood flow and surface electromyography to repeated bouts of eccentric exercise. *J Strength Cond Res.* 29:1017-
1429 26.

1430 Houghton D, Onambele GL (2012) Can a standard dose of eicosapentaenoic acid (EPA) supplementation reduce the
1431 symptoms of delayed onset of muscle soreness? *J Int Soc Sports Nutr.* 9:2.

1432 Howatson G, van Someren KA (2007) Evidence of a contralateral repeated bout effect after maximal eccentric
1433 contractions. *Eur J Appl Physiol.* 101:207-14.

1434 Howatson G, Hoad M, Goodall S, Tallent J, Bell PG, French DN (2012) Exercise-induced muscle damage is reduced in
1435 resistance-trained males by branched chain amino acids: a randomized, double-blind, placebo controlled study. *J Int Soc*
1436 *Sports Nutr.* 9:20.

1437 Howatson G, McHugh MP, Hill JA, Brouner J, Jewell AP, van Someren KA, Shave RE, Howatson SA (2010) Influence
1438 of tart cherry juice on indices of recovery following marathon running. *Scand J Med Sci Sports.* 20:843-52.

1439 Howatson G, van Someren KA (2007) Evidence of a controlateral repeated bout effect after maximal eccentric
1440 contractions. *Eur J Appl Physiol.* 101:207-14.

1441 Howatson G, van Someren KA (2008) The prevention and treatment of exercise-induced muscle damage. *Sports Med.*
1442 38:483-503.

1443 Howatson G, van Someren KA, Leeder JDC (2016) BASES Expert Statement on Athletic Recovery Strategies. *The Sport*
1444 *and Exercise Scientist*, Issue 48, Summer, 2016.

1445 Hsu CC, Ho MC, Lin LC, Su B, Hsu MC (2005) American ginseng supplementation attenuates creatine kinase level
1446 induced by submaximal exercise in human beings. *World J Gastroenterol*. 11:5327-31.

1447 Hurley CF, Hatfield DL, Riebe DA (2013) The effect of caffeine ingestion on delayed muscle soreness. *J Strength Cond*
1448 *Res*. 27:3101-9.

1449 Hurst RD, Wells RW, Hurst SM, McGhie TK, Cooney JM, Jensen DW (2010) Blueberry fruit polyphenolics suppress
1450 oxidative stress induced skeletal muscle cell damage in vitro. *Mol Nutr Food Res*. 54:353-363.

1451 Hutchison AT, Flieller EB, Dillon KJ, Leverett BD (2016) Black currant nectar reduces muscle damage and inflammation
1452 following a bout of high-intensity eccentric contractions. *J Diet Suppl*. 13:1-15.

1453 Hyldahl RD, Hubal MJ (2014) Lengthening our perspective: morphological, cellular, and molecular responses to eccentric
1454 exercise. *Muscle Nerve*. 49:155-70.

1455 Institute of Medicine Food and Nutrition Board (2000) Dietary Reference Intakes: Vitamin C, vitamin E, selenium, and
1456 carotenoids. Ed. National Academy Press, Washington.

1457 Jackson MJ, Jones DA, Edwards RH (1984) Experimental skeletal muscle damage: the nature of the calcium-activated
1458 degenerative processes. *Eur J Clin Invest*. 14:369-74.

1459 Jäger R, Purpura M, Kerksick CM (2019) Eight weeks of a high dose of curcumin supplementation may attenuate
1460 performance decrements following muscle-damaging exercise. *Nutrients*. 11(7).

1461 Jäger R, Purpura M, Stone JD, Turner SM, Anzalone AJ, Eimerbrink MJ, Pane M, Amoroso A, Rowlands DS, Oliver JM
1462 (2016b) Probiotic *Streptococcus thermophilus* FP4 and *Bifidobacterium breve* BR03 supplementation attenuates
1463 performance and range-of-motion decrements following muscle damaging exercise. *Nutrients*. 8(10). pii: E642.

1464 Jäger R, Shields KA, Lowery RP, De Souza EO, Partl JM, Hollmer C, Purpura M, Wilson JM (2016a) Probiotic *Bacillus*
1465 *coagulans* GBI-30, 6086 reduces exercise-induced muscle damage and increases recovery. *Peer J*. 4:e2276.

1466 Jajtner AR, Hoffman JR, Townsend JR, Beyer KS, Varanoske AN, Church DD, Oliveira LP, Herrlinger KA, Random-
1467 Aizik S, Fukuda DH, Stout JR (2016) The effect of polyphenols on cytokine and granulocyte response to resistance
1468 exercise. *Physiol Rep*. 4(24). pii:e13058.

1469 Jakerman JR, Lambrick DM, Wooley B, Babraj JA, Faulkner JA (2017) Effect of an acute dose of omega-3 fish oil
1470 following exercise-induced muscle damage. *Eur J Appl Physiol*. 117:575-582.

1471 Jacob K, Periago MJ, Böhm V, Berruezo GR (2008) Influence of lycopene and vitamin C from tomato juice on biomarkers
1472 of oxidative stress and inflammation. *Br J Nutr*. 99:137-46.

1473 Jamurtas AZ (2018) Exercise-induced muscle damage and oxidative stress. *Antioxidants (Basel)*. 7:E50.

1474 Jamurtas AZ, Theocharis V, Tofas T, Tsiokanos A, Yfanti C, Paschalis V, Koutedakis Y, Nosaka K (2005) comparison
1475 between leg and arm eccentric exercises of the same relative intensity on indices of muscle damage. *Eur J Appl Physiol.*
1476 95:179-85.

1477 Jang HJ, Lee JD, Jeon HS, Kim AR, Kim S, Lee HS, Kim KB (2018) Metabolic profiling of eccentric exercise-induced
1478 muscle damage in human urine. *Toxicol Res.* 34:199-210.

1479 Jo E, Bartosh R, Auslander AT, Directo D, Osmond A, Wong MW (2019) Post-exercise recovery following 30-day
1480 supplementation of trans-resveratrol and polyphenol-enriched extracts. *Sports (Basel)* 7(10). pii:E226.

1481 Jones DA, Newham DJ, Clarkson PM (1987) Skeletal muscle stiffness and pain following eccentric exercise of the elbow
1482 flexors. *Pain.* 30:233-42.

1483 Jouris KB, McDaniel JL, Weiss EP (2011) The effect of omega-3 fatty acid supplementation on the inflammatory
1484 response to eccentric strength exercise. *J Sports Sci Med.* 10:432-8.

1485 Jówko E, Sacharuk J, Balasinska B, Wilczak J, Charmas M, Ostaszewski P, Charmas R (2012) Effect of a single dose of
1486 green tea polyphenols on the blood markers of exercise-induced oxidative stress in soccer players. *Int J Sport Nutr Exerc*
1487 *Metab.* 22:486-96.

1488 Jude S, Amalraj A, Kunnumakkara AB, Divya C, Löffler M, Gopi S (2018) Development of validated methods and
1489 quantification of curcuminoids and curcumin metabolites and their pharmacokinetic study of oral administration of
1490 complete natural turmeric formulation (Cureit™) in human plasma via UPLC/ESI-Q-TOF-MS spectrometry. *Molecules.*
1491 23:2415.

1492 Jung HL, Kwak HE, Kim SS, Kim YC, Lee CD, Byurn HK, Kang HY (2011) Effects of Panax ginseng supplementation
1493 on muscle damage and inflammation after uphill treadmill running in humans. *Am J Chin Med.* 39:441-50.

1494 Kandylis P, Kokkinomagoulos E (2020) Food applications and potential health benefits of pomegranate and its
1495 derivatives. *Foods.* 9(2).

1496 Karabalaefar S, Hefzollasan M, Behpoor N, Ghalehgir S (2013) Effect of caffeine on the amount of perceived pain, joint
1497 range of motion and edema after delayed muscle soreness. *Pedagogics Psychology, Medical-Biological Problems of*
1498 *Physical Training and Sports.* 1:96-100.

1499 Kastello GM, Bretl MC, Clark ER, Delvaux CL, Hoepfner JD, McNea LP, Mesmer JA, Meyer CG, Mitchener CL, Rivera
1500 DA, Ruhl SN, Schuller KM, Steffen JL, Strauss JD (2014) The effect of cherry supplementation on exercise-induced
1501 oxidative stress. *J Food Nutr Sci* 1:1-6.

1502 Kim JM, Araki S, Kim DJ, Park CB, Takasuka N, Baba-Toriyama H, Ota T, Nir Z, Khachik F, Shimidzu N, Tanaka Y,
1503 Osawa T, Urajii T, Murakoshi M, Nishino H, Tsuda H (1998) Chemopreventive effects of carotenoids and curcumins on
1504 mouse colon carcinogenesis after 1,2-dimethylhydrazine initiation. *Carcinogenesis.* 19:81-5.

1505 Kim J, Lee J (2014) A review of nutritional intervention on delayed onset muscle soreness. Part I. *J exerc rehabil.* 10:349-
1506 56.

1507 Kim J, Lee J, Kim S, Yoon D, Kim J, Sung DJ (2015) Role of creatine supplementation in exercise-induced muscle
1508 damage: a mini review. *J Exerc Rehabil.* 11:244-50.

1509 Kim J, Lee KP, Beak S, Kang HR, Kim YK, Lim K (2019) Effect of black chokeberry on skeletal muscle damage and
1510 neuronal cell death. *J Exerc Nutrition Biochem.* 23:26-31.

1511 Kim J, So WY (2019) Effects of acute grape seed extract supplementation on muscle damage after eccentric exercise: A
1512 randomized, controlled clinical trial. *J Exerc Sci Fit.* 17:77-79.

1513 Kirk B, Mitchell J, Jackson M, Amirabdollahian F, Alizadehkhayat O, Clifford T (2017) A2 milk enhances dynamic
1514 muscle function following repeated sprint exercise, a possible ergogenic aid for A1-protein intolerant athletes? *Nutrients.*
1515 9(2).

1516 Klinkenberg LJ, Res PT, Haenen GR, Bast A, van Loon LJ, van Dieijen-Visser MP, Meex SJ (2013) Effect of antioxidant
1517 supplementation on exercise-induced cardiac troponin release in cyclists: a randomized trial. *PLoS One.* 8:e79280.

1518 Knitter AE, Panton L, Rathmacher JA, Petersen A, Sharp R (2000) Effect of β -hydroxy β -methylbutyrate on muscle
1519 damage after a prolonged run. *J Appl Physio.* 89:1340-4.

1520 Kocaadam B, Şanlıer N (2017) Curcumin, an active component of turmeric (*Curcuma longa*), and its effect on health.
1521 *Crit Rev Food Sci Nutr.* 57:2889-2895.

1522 Köhne JL, Ormsbee MJ, McKune AJ (2016) Supplementation strategies to reduce muscle damage and improve recovery
1523 following exercise in females: a systematic review. *Sports (Basel)* 4(4). pii: E51.

1524 Konrad M, Nieman DC, Henson DA, Kennerly KM, Jin, Wallner-Liebmann SJ (2011) The acute effect of ingesting a
1525 quercetin-based supplement on exercise-induced inflammation and immune changes in runners. *Int J Sport Nutr Exerc*
1526 *Metab.* 21:338-46.

1527 Korge P, Byrd SK, Campbell KB (1993) Functional coupling between sarcoplasmic-reticulum-bound creatine kinase and
1528 Ca^{2+} -ATPase. *Eur J Biochem.* 213:973-980.

1529 Kraemer WJ, Spiering BA, Volek JS, Ratamess NA, Sharman MJ, Rubin MR, French DN, Silvestre R, Hatfield DL, Van
1530 Heest JL, Vingren JL, Judelson DA, Deshenes MR, Maresh CM (2006) Androgenic responses to resistance exercise:
1531 effects of feeding and L-carnitine. *Med Sci Sports Exerc.* 38:1288-96.

1532 Kraemer WJ, Volek JS, French DN, Rubin MR, Sharman MJ, Gómez AL, Ratamess NA, Newton RU, Jemiolo B, Craig
1533 BW, Häkkinen K (2003) The effects of L-carnitine L-tartrate supplementation on hormonal responses to resistance
1534 exercise and recovery. *J Strength Cond Res.* 17:455-62.

1535 Kreider RB, Kalman DS, Antonio J, Ziegenfuss TN, Wildman R, Collins R, Candow DG, Kleiner SM, Almada AL,
1536 Lopez HL (2017) International Society of Sports Nutrition position stand: safety and efficacy of creatine supplementation
1537 in exercise, sport, and medicine. *J Int Soc Sports Nutr.* 14:18.

1538 Kuehl KS, Perrier ET, Elliot DL, Chesnutt JC (2010) Efficacy of tart cherry juice in reducing muscle pain during
1539 running: a randomized controlled trial. *J Int Soc Sports Nutr.* 7:17.

1540 Lamb KL, Ranchordas MK, Johnson E, Denning J, Downin F, Lynn A (2019) No effect of tart cherry juice or pomegranate
1541 juice on recovery from exercise-induced muscle damage in non-resistance trained men. *Nutrients.* 11(7):1593.

1542 Lamb KL, Ranchordas MK, Johnson E, Denning J, Downing F, Lynn A (2019) No effect of tart cherry juice or
1543 pomegranate juice on recovery from exercise-induced muscle damage in non-resistance trained men. *Nutrients.* 11(7).

1544 Lamprecht M, Bogner S, Schippinger G, Steinbauer K, Fankhauser F, Hallstroem S, Schuetz B, Greilberg JF (2012)
1545 Probiotic supplementation affects markers of intestinal barrier, oxidation, and inflammation in trained men: a randomized,
1546 double-blinded, placebo-controlled trial. *J Int Soc Sports Nutr.* 9:45.

1547 Larson-Meyer DE, Douglas CS, Thomas JJ, Johnson EC, Barcal JN, Heller JE, Hollis BW, Halliday TM (2019)
1548 Validation of a vitamin D specific questionnaire to determine vitamin D status in athletes. *Nutrients* 11:E2732.

1549 LaStayo PC, Woolf JM, Lewek MD, Snyder-Mackler L, Reich T, Lindstedt SL (2003) Eccentric muscle contractions:
1550 their contribution to injury, prevention, rehabilitation, and sport. *J Orthop Sports Phys Ther.* 33:557-71.

1551 Lawler JM, Barnes WS, Wu G, Song W, Demaree S (2002) Direct antioxidant properties of creatine. *Biochem Biophys*
1552 *Res Commun.* 290:47-52.

1553 Lee J, Goldfarb AH, Rescino MH, Hegde S, Patrick S, Apperson K (2002) Eccentric exercise effect on blood oxidative-
1554 stress markers and delayed onset of muscle soreness. *Med Sci Sports Exerc.* 34:443-448.

1555 Leeder J, Gissane C, van Someren K, Gregson W, Howatson G (2012) Cold water immersion and recovery from strenuous
1556 exercise: a meta-analysis. *Br J Sports Med.* 46:233-40.

1557 Leeder JDC, van Someren KA, Gaze D, Jewell A, Deshmukh NI, Shah I, Barker J, Howatson G (2014) Recovery and
1558 adaptation from repeated intermittent-sprint exercise. *Int J Sports Physiol Perform.* 9:489-96.

1559 Legault Z, Bagnall N, Kimmerly DS (2015) The influence of oral l-glutamine supplementation on muscle strength
1560 recovery and soreness following unilateral knee extension eccentric exercise. *Int J Sport Nutr Exerc Metab.* 25:417-26.

1561 Lembke P, Capodice J, Hebert K, Swenson T (2014) Influence of omega-3 (n3) index on performance and wellbeing in
1562 young adults after heavy eccentric exercise. *J Sports Sci Med.* 13:151-6.

1563 Lenn J, Uhl T, Mattacola C, Boissonneault G, Yates J, Ibrahim W, Bruckner G (2002) The effects of fish oil and
1564 isoflavones on delayed onset muscle soreness. *Med Sci Sports Exerc.* 34:1605-13.

1565 Levers K, Dalton R, Galvan E, Goodenough C, O'Connor A, Simbo S, Barringer N, Mertens-Talcott SU, Rasmussen C,
1566 Greenwood M, Riechman S, Crouse S, Kreider RB (2015) Effects of powdered Montmorency tart cherry supplementation
1567 on an acute bout of intense lower body strength exercise in resistance trained males. *J Int Soc Sports Nutr.* 12:41.

1568 Levers K, Dalton R, Galvan E, O'Connor A, Goodenough C, Simbo S, Mertens-Talcott SU, Rasmussen C, Greenwood
1569 M, Riechman S, Crouse S, Kreider RB (2016) Effects of powdered Montmorency tart cherry supplementation on acute
1570 endurance exercise performance in aerobically trained individuals. *J Int Soc Sports Nutr.* 26:13-22.

1571 Li P, Wu G (2018) Roles of dietary glycine, proline, and hydroxyproline in collagen synthesis and animal growth. *Amino
1572 acids.* 50:29-38.

1573 Lieber RL, Fridén J (1999) Mechanisms of muscle injury after eccentric contraction. *J Sci Med Sport.* 2:253-65.

1574 Lopez HL, Habowski SM, Sandrock JE, Kedia AW, Ziegenfuss TN (2014) Effects of BioCell Collagen on connective
1575 tissue protection and functional recovery from exercise in healthy adults: a pilot study. *Journal of the International Society
1576 of Sports Nutrition.* 11(Suppl 1):P48.

1577 Lyall KA, Hurst SM, Cooney J, Jensen D, Lo K, Hurst RD, Stevenson LM (2009) Short-term blackcurrant extract
1578 consumption modulates exercise-induced oxidative stress and lipopolysaccharide-stimulated inflammatory responses.
1579 *Am J Physiol Regul Integr Comp Physiol* 297:R70-81.

1580 Ma S, Tominaga T, Kanda K, Sugama K, Omae C, Hashimoto S, aoyama K, Yoshikai Y, Suzuki K (2020) Effects of an
1581 8-week protein supplementation regimen with hyperimmunized cow milk on exercise-induced organ damage and
1582 inflammation in male runners: a randomized, placebo controlled, cross-over study. *Biomedicines.* 8(3). pii: E51.

1583 Mach N, Fuster-Botella D (2017) Endurance exercise and gut microbiota: a review. *J Sport Health Sci.* 6:179-197.

1584 Machado AS, da Silva W, Souza MA, Carpes FP (2018) Green tea extract preserves neuromuscular activation and muscle
1585 damage markers in athletes under cumulative fatigue. *Front Physiol.* 9:1137.

1586 Machado M, Antunes WD, Tamy ALM, Azevedo PG, Barreto JG, Hackney AC (2009a) Effect of a single dose of caffeine
1587 supplementation and intermittent-interval exercise on muscle damage markers in soccer players. *J Exerc Sci Fit.* 2:91-97

1588 Machado M, Koch AJ, Willardson JM, dos Santos FC, Curty VM, Pereira LN (2010) Caffeine does not augment markers
1589 of muscle damage or leucocytosis following resistance. *Int J Sports Physiol Perform.* 5:18-26.

1590 Machado M, Pereira R, Sampaio-Jorge F, Knifis F, Hackney A (2009) Creatine supplementation: effects on blood creatine
1591 kinase activity responses to resistance exercise and creatine kinase activity measurement. *Brazilian Journal of
1592 Pharmaceutical Sciences.* 45:4.

1593 Machado M, Vigo JFF, Breder AC, Simoes JR, Ximenes MC, Hackney AC (2009b) Effect of short term caffeine
1594 supplementation and intermittent exercise on muscle damage markers. *Biol. Sport* 26:3-11

1595 Machin DR, Christmas KM, Chou TH, Hill SC, Van Pelt DW, Trombold JR, Coyle EF (2014) Effects of differing dosages
1596 of pomegranate juice supplementation after eccentric exercise. *Physiology Journal*.

1597 Mackey AL, Brandstetter S, Schjerling P, Bojsen-Moller J, Qvortrup K, Pedersen MM, Doessing S, Kjaer M, Magnusson
1598 SP, Langberg H. Sequenced response of extracellular matrix deadhesion and fibrotic regulators after muscle damage is
1599 involved in protection against future injury in human skeletal muscle. *FASEB J*. 25:1943-59.

1600 Malaguti M, Angeloni C, Hrelia S (2013) Polyphenols in exercise performance and prevention of exercise-induced muscle
1601 damage. *Oxid Med Cell Longev*. 825928.

1602 Maridakis V, O'Connor PJ, Dudley GA, McCully KK (2007) Caffeine attenuates delayed-onset muscle pain and force
1603 loss following eccentric exercise. *J Pain*. 8:237-43.

1604 Marques CG, Santos VC, Levada-Pires AC, Jacintho TM, Gorjão R, Pithon-Curi TC, Cury-Bonaventura MF (2015)
1605 Effects of DHA-rich fish oil supplementation on the lipid profile, markers of muscle damage, and neutrophil function in
1606 wheelchair basketball athletes before and after acute exercise. *Appl Physiol Nutr Metab*. 40:596-604.

1607 Martarelli D, Verdenelli MC, Scuri S, Cocchioni M, Silvi S, Cecchini C, Pompei P (2011) Effect of a probiotic intake on
1608 oxidant and antioxidant parameters in plasma of athletes during intense exercise training. *Curr Microbiol*. 62:1689-96.

1609 Martin-Rincon M, Gelabert-Rebato M, Galvan-Alvarez V, Gallego-Selles A, Marinez-Canton M, Lopez-Rios L, Wiebe
1610 JC, Martin-Rodriguez S, Arteaga-Ortiz R, Dorado C, Perez-Regalado S, Santana A, Morales-Alamo D, Calbet JAL (2020)
1611 Supplementation with a mango leaf extract (Zynamite®) in combination with quercetin attenuates muscle damage and
1612 pain and accelerates recovery after strenuous damaging exercise. *Nutrients*. 12(3). pii: E614.

1613 Martínez-Sánchez A, Alacid F, Rubio-Arias JA, Fernández-Lobato B, Ramos-Campo DJ, Aguayo E (2017b)
1614 Consumption of watermelon juice enriched in l-citrulline and pomegranate ellagitannins enhanced metabolism during
1615 physical exercise. *J Agric Food Chem*. 65:4395-4404.

1616 Martínez-Sánchez A, Ramos-Campo DJ, Fernández-Lobato B, Rubio-Arias JA, Alacid F, Aguayo E (2017a)
1617 Biochemical, physiological, and performance response of a functional watermelon juice enriched in L-citrulline during a
1618 half-marathon race. *Food Nutr Res*. 61:1330098.

1619 Matsumura MD, Zavorsky GS, Smoliga JM (2015) The effects of pre-exercise ginger supplementation on muscle damage
1620 and delayed onset muscle soreness. *Phytother Res*. 29:887-93.

1621 McAnulty SR, McAnulty LS, Nieman DC, Dumke CL, Morrow JD, Utter AC, Henson DA, Proulx WR, George GL
1622 (2004) Consumption of blueberry polyphenols reduces exercise-induced oxidative stress compared to vitamin C.
1623 *Nutrition Res*. 24:209-221.

1624 McAnulty SR, McAnulty LS, Nieman DC, Quindry JC, Hosick PA, Hudson MH, Still L, Henson DA, Milne GL, Morrow
1625 JD, Dumke CL, Utter AC, Triplett NT, Dibanardi A (2008) Chronic quercetin ingestion and exercise-induced oxidative
1626 damage and inflammation. *Appl Physiol Nutr. Metab.* 33:254-62.

1627 McCormick R, Peeling P, Binnie M, Dawson B, Sim M (2016) Effect of tart cherry juice on recovery and next day
1628 performance in well-trained water polo players. *J Int Soc Sports Nutr.* 13:41.

1629 McFarlin BK, Venable AS, Henning AL, Sampson JN, Pennell K, Vingren JL, Hill DW (2016) Reduced inflammatory
1630 and muscle damage biomarkers following oral supplementation with bioavailable curcumin. *BBA Clin.* 5:72-8.

1631 McGinley C, Shafat A, Donnelly AE (2009) Does antioxidant vitamin supplementation protect against muscle damage?
1632 *Sports Med.* 39:1011-32.

1633 McHugh MP (2003) Recent advances in the understanding of the repeated bout effect: the protective effect against muscle
1634 damage from a single bout of eccentric exercise. *Scand J Med Sci Sports.* 13:88-97.

1635 McHugh MP, Connolly DA, Eston RG, Gleim GW (1999) Exercise-induced muscle damage and potential mechanisms
1636 for the bout effect. *Sports Med* 27:157-70.

1637 McKinnon NB, Graham MT, Tiidus PM (2012) Effect of creatine supplementation on muscle damage and repair
1638 following eccentrically-induced damage to the elbow flexors muscles. *J Sports Sci Med.* 11:653-9.

1639 McLeay Y, Barnes MJ, Mundel T, Hurst SM, Hurst RD, Stannard SR (2012) Effect of New Zealand blueberry consumption
1640 on recovery from eccentric exercise-induced muscle damage. *J Int Soc Sports Nutr.* 9:19.

1641 McLeay Y, Stannard S, Barnes M (2017) The effect of taurine on the recovery from eccentric exercise-induced muscle
1642 damage in males. *Antioxidants (Basel)* 6(4).

1643 Meamarbashi A, Rajabi A (2015) Preventive effects of 10-day supplementation with saffron and indomethacin on the
1644 delayed-onset muscle soreness. *Clin J Sport Med.* 25:105-12.

1645 Mero AA, Ojala T, Hulmi JJ, Puurtinen R, Karila TA, Seppälä T (2010) Effects of alfa-hydroxy-isocaproic acid on body
1646 composition, DOMS and performance in athletes. *J Int Soc Sports Nutr.* 7:1.

1647 Miliadis GA, Nomikos T, Fragopoulou E, Athanasopoulos S, Antonopoulou S (2006) Effect of baseline serum levels of Se
1648 on markers of eccentric exercise-induced muscle injury. *Biofactors.* 26:161-70.

1649 Miller PC, Bailey SP, Barnes ME, Derr SJ, Hall EE (2004) The effects of protease supplementation on skeletal muscle
1650 function and DOMS following downhill running. *J Sports Sci.* 22:365-72.

1651 Minajeva A, Ventura-Clapier R, Veksler V (1996) Ca²⁺ uptake by cardiac sarcoplasmic reticulum ATPase in situ strongly
1652 depends on bound creatine kinase. *Pflugers Arch.* 432:904-912.

1653 Mizumura K, Taguchi T (2016) Delayed onset muscle soreness: involvement of neurotrophic factors. *J Physiol Sci.* 66:43-
1654 52.

1655 Myburgh KH (2014) Polyphenol supplementation: Benefits for exercise performance or oxidative stress? *Sports Med.* 44
1656 Suppl 1:S57-70.

1657 Montenegro CF, Kwong DA, Minow ZA, Davis BA, Lozada CF, Casazza GA (2017) Betalain-rich concentrate
1658 supplementation improves exercise performance and recovery in competitive triathletes. *Appl Physiol Nutr Metab.*
1659 42:166-172.

1660 Morgan DL (1990) New insights into the behavior of muscle during lengthening. *Biophys J.* 57:209-21.

1661 Morgan PT, Wollman PM, Jackman SR, Bowtell JL (2018) Flavanol-rich cacao mucilage juice enhances recovery of
1662 power but not strength from intensive exercise in healthy, young men. *Sports (Basel).* 6(4). pii: E159.

1663 Nakhostin-Roohi B, Javanamani R, Zardoost N, Ramazanzadeh R (2017) Influence of glutamine supplementation on
1664 muscle damage and oxidative stress indices following 14Km running. *Hormozgan Medical Journal.* 20:323-331.

1665 Nakhostin-Roohi B, Moradlou AN, Hamidabad SM, Ghanivand B (2016) The effect of curcumin supplementation on
1666 selected markers of delayed onset muscle soreness (DOMS). *Annals of Applied Sport Science.* 2:25-31.

1667 Naughton M, Miller J, Slater GJ (2018) Impact-induced muscle damage: performance implications in response to a novel
1668 collision simulator and associated timeline of recovery. *J Sports Sci Med.* 17:417-425.

1669 Nemati A, Alipanah-Moghadam R, Molazadeh L, Naghizadeh Baghi A (2019) The effect of glutamine supplementation
1670 on oxidative stress and matrix metalloproteinase 2 and 9 after exhaustive exercise. *Drug Des Devel Ther.* 13:4215-4223.

1671 Neubauer O, König D, Wagner KH (2008) Recovery after an ironman triathlon:sustained inflammatory responses and
1672 muscular stress. *Eur J Appl Physiol.* 104:417-26.

1673 Neubauer O, Sabapathy S, Lazarus R, Jowett JB, Desbrow B, Peake JM, Cameron-Smith D, Haseler LJ, Wagner KH,
1674 Bulmer AC (2013) Transcriptome analysis of neutrophils after endurance exercise reveals novel signaling mechanisms
1675 in the immune response to physiological stress. *J Appl Physiol.* 114:1677-88.

1676 Nguyen HX, Tidball JG (2003) Interactions between neutrophils and macrophages promote macrophage killing of rat
1677 muscle cells in vitro. *J Physiol.* 547:125-32.

1678 Nicol LM, Rowlands DS, Fazakerly R, Kellett J (2015) Curcumin supplementation likely attenuates delayed onset muscle
1679 soreness (DOMS). *Eur J Appl Physiol.* 115:1769-77.

1680 Nieman DC, Gillitt ND, Shanelly RA, Dew D, Meaney MP, Luo B (2013) Vitamin D2 supplementation amplifies
1681 eccentric exercise-induced muscle damage in NASCAR pit crew athletes. *Nutrients.* 6:63-75.

1682 Nieman DC, Henson DA, Davis JM, Angela Murphy E, Jenkins DP, Gross SJ, Carmichael MD, Quindry JC, Dumke CL,
1683 Utter AC, McAnulty SR, McAnulty LS (2007a) Quercetin's influence on exercise-induced changes in plasma cytokines
1684 and muscle and leukocyte cytokine mRNA. *J Appl Physiol.* 103:1728-35.

1685 Nieman DC, Henson DA, Davis JM, Dumke CL, Gross SJ, Jenkins DP, Murphy EA, Carmichael MD, Quindry JC,
1686 McAnulty SR, McAnulty LS, Utter AC, Mayer EP (2007b) Quercetin ingestion does not alter cytokine changes in athletes
1687 competing in the western states endurance run. *J Interferon Cytokine Res.* 27:1003-11.

1688 Nieman DC, Henson DA, McAnulty SR, Jin F, Maxwell KR (2009) n-3 polyunsaturated fatty acids do not alter immune
1689 and inflammation measures in endurance athletes. *Int J Sport Nutr Exerc Metab.* 19:536-46.

1690 Nishizawa M, Hara T, Miura T, Fujita S, Yoshigai E, Ue H, Hayashi Y, Kwon AH, Okumura T, Isaka T (2011)
1691 Supplementation with a flavanol-rich lychee fruit extract influences the inflammatory status of young athletes. *Phytother*
1692 *Res.* 25:1486-93.

1693 Nosaka K, Sacco P, Mawatari K (2006) Effects of amino acid supplementation on muscle soreness and damage. *Int J*
1694 *Sport Nutr Exerc Metab.* 16:620-35.

1695 Nunan D, Howatson G, van Someren KA (2010) Exercise-induced muscle damage is not attenuated by β -hydroxy β -
1696 methylbutyrate and alpha-ketoisocaproic acid supplementation. *J Strength Cond Res.* 24:531-7.

1697 O'Fallon KS, Kaushik D, Michniak-Kohn B, Dunne CP, Zambraski EJ, Clarkson PM (2012) Effects of quercetin
1698 supplementation on markers of muscle damage and inflammation after eccentric exercise. *Int J Sport Nutr Exrc Metab.*
1699 22:430-7.

1700 Ochi E, Tsuchiya Y (2018) Eicosapentaenoic acid (EPA) and Docosahexaenoic acid (DHA) in muscle damage and
1701 function. *Nutrients.* 10(5).

1702 Olsen S, Aagaard P, Kadi F, Tufekovic G, Verney J, Olesen JL, Suetta C, Kjaer M (2006) Creatine supplementation
1703 augments the increase in satellite cell and myonuclei number in human skeletal muscle induced by strength training. *J*
1704 *Physiol* 573:525-34.

1705 Osmond AD, Directo DJ, Elam ML, Juache G, Kreipke VC, Saralegui DE, Wildman R, Wong M, Jo E (2019) The effects
1706 of leucine-enriched branched-chain amino acid supplementation on recovery after high-intensity resistance exercise. *Int*
1707 *J Sports Physiol Perform.* 14(8):1081-1088.

1708 Owens DJ, Fraser WD, Close GL (2015) Vitamin D and the athlete: emerging insights. *Eur J Sport Sci.* 15:73-84.

1709 Owens DJ, Sharples AP, Polydorou I, Alwan N, Donovan T, Tang J, fraser WD, Cooper RG, Morton JP, Stewart C,
1710 Close GL (2015) A systems-based investigation into vitamin D and skeletal muscle repair, regeneration, and
1711 hypertrophy. *Am J Physiol Endocrinol Metab.* 309:E1019-31.

1712 Owens DJ, Twist C, Cobbley JN, Howatson G, Close GL (2019) Exercise-induced muscle damage: What is it, what
1713 causes it and what are the nutritional solutions?. *Eur J Sport Sci.* 19:71-85.

1714 Paddon-Jones D, Keech A, Jenkins D (2001) Short-term β -hydroxy β -methylbutyrate supplementation does not reduce
1715 symptoms of eccentric muscle damage. *Int J Sport Nutr Exerc Metab.* 11:442-50.

1716 Panza VP, Diefenthaler F, Tamborindeguy AC, Camargo C de Q, de Moura BM, Brunetta HS, Sakugawa RL, de Oliveira
1717 MV, Puel Ede O, Nunes EA, da Silva EL (2016) Effects of mate tea consumption on muscle strength and oxidative stress
1718 markers after eccentric exercise. *Br J Nutr.* 115:1370-8.

1719 Parandak K, Arazi H, Khoshkharesh F, Nakhostin-Roohi B (2014) The effect of two-week L-carnitine supplementation
1720 on exercise-induced oxidative stress and muscle damage. *Asian J Sports Med.* 5:123-8.

1721 Pari L, Murugan P (2006) Tetrahydrocurcumin: effect on chloroquine-mediated oxidative damage in rat kidney. *Basic
1722 Clin Pharmacol Toxicol.* 99:329-34.

1723 Parthimos T, Schulpis KH, Angelogianni P, Tsopanakis C, Parthimos N, Tsakiris S (2008) The in vivo and in vitro effects
1724 of L-carnitine supplementation on the erythrocyte membrane acetylcholinesterase, Na⁺, K⁺-ATPase activities in basketball
1725 players. *Clin Chem Lab.* 46:137-42.

1726 Pasiakos SM, Lieberman HR, McLellan TM (2014) Effects of protein supplements on muscle damage, soreness and
1727 recovery of muscle function and physical performance: a systematic review. *Sports Med* 44:655-70.

1728 Patrizio F, Ditroilo M, Felici F, Duranti G, De Vito G, Sabatini S, Sacchetti M, Bazzucchi I (2018) The acute effect of
1729 quercetin on muscle performance following a single resistance training session. *Eur J Appl Physiol.* 118:1021-1031.

1730 Peake JM, Neubauer O, Della Gatta PA, Nosaka K (2017) Muscle damage and inflammation during recovery from
1731 exercise. *J Appl Physiol.* 22:559-570.

1732 Peake JM, Suzuki K, Wilson G, Hordern M, Nosaka K, Mackinnon L, Coombes JS (2005) Exercise-induced muscle
1733 damage, plasma cytokines, and markers of neutrophil activation. *Med Sci Sports Exerc.* 37:737-45.

1734 Pérez-Guisado J, Jakerman PM (2010) Citrulline malate enhances athletic anaerobic performance and relieves muscle
1735 soreness. *J Strength Cond Res.* 24:1215-22.

1736 Peschek K, Pritchett R, Bergman E, Pritchett K (2013) The effects of acute post exercise consumption of two cocoa-based
1737 beverages with varying flavanol content on indices of muscle recovery following downhill treadmill running. *Nutrients.*
1738 6:50-62.

1739 Peternelj TT, Coombes JS (2011) Antioxidant supplementation during exercise training: beneficial or detrimental? *Sports
1740 Med.* 41:1043-69.

1741 Pfeiffer E, Hoehle SI, Walch SG, Riess A, Sólyom AM, Metzler M (2007) Curcuminoids form reactive glucuronides in
1742 vitro. *J Agric Food Chem.* 55:538-44.

1743 Phillips SM, Van Loon LJ (2011) Dietary protein for athletes: from requirements to optimum adaptation. *J Sports Sci.*
1744 1:S29-38.

1745 Phillips T, Childs AC, Dreon DM, Phinney S, Leeuwenburgh C (2003) A dietary supplement attenuates IL-6 and CRP
1746 after eccentric exercise in untrained males. *Med Sci Sports Exerc.* 35:2032-7.

1747 Philpott JD, Donnelly C, Walshe IH, MacKinley EE, Dick J, Galloway SDR, Tipton KD, Witard OC (2018) Adding
1748 fish oil to whey protein, leucine, and carbohydrate over a six-week supplementation period attenuates muscle soreness
1749 following eccentric exercise in competitive soccer players. *Int J Sport Nutr Exerc Metab.* 28:26-36.

1750 Philpott JD, Witard OC, Galloway SDR (2019) Applications of omega-3 polyunsaturated fatty acid supplementation for
1751 sport performance. *Res Sports Med.* 27:219-237.

1752 Pincheira PA, Hoffman BW, Cresswell AG, Carroll TJ, Brown NAT, Lichtwark GA (2018). The repeated bout effect can
1753 occur without mechanical and neuromuscular changes after a bout of eccentric exercise. *Scand J Med Sci Sports.* 28:2123-
1754 2134.

1755 Pingitore A, Lima GPP, Mastorci F, Quinones A, Lervasi G, Vassalle C (2015) Exercise and oxidative stress: potential
1756 effects of antioxidant dietary strategies in sports. *Nutrition [internet]. Elsevier Inc.* 31:916-22.

1757 Plowman SA, Smith DL (2013) *Exercise Physiology: For Health, Fitness and Performance.* Lippincott & Wilkins,
1758 Philadelphia.

1759 Pons V, Riera J, Capó X, Martorell M, Sureda A, Tur JA, Drobnic F, Pons A (2018) Calorie restriction regime enhances
1760 physical performance of trained athletes. *J Int Soc Sports Nutr.* 15:12.

1761 Portal S, Zadik Z, Rabinowitz J, Pilz-Burstein R, Adler-Portal D, Meckel Y, Cooper DM, Eliakim A, Nemet D (2011)
1762 The effect of HMB supplementation on body composition, fitness, hormonal and inflammatory mediators in elite
1763 adolescent volleyball players: a prospective randomized, double-blind, placebo-controlled study. *Eur J Appl Physiol.*
1764 111:2261-9.

1765 Poulos A, Georgakouli K, Draganidis D, Deli CK, Tsimeas PD, Chatzinikolaou A, Papanikolaou K, Batrakoulis A, Mohr
1766 M, Jamurtas AZ, Fatouros IG (2019) Protein-based supplementation to enhance recovery in team sports: what is the
1767 evidence? *J Sports Sci Med.* 18:523-536.

1768 Poulous A, Georgakouli K, Draganidis D, Deli CK, Tsimeas PD, Chatzinikolaou A, Papanikolaou K, Batrakoulis A,
1769 Mohr M, Jamurtas AZ, Fatouros IG (2019) Protein-based supplementation to enhance recovery in team sports: what is
1770 the evidence? *J Sports Sci Med.* 18:523-536.

1771 Proske U, Morgan DL (2001) Muscle damage from eccentric exercise: mechanism, mechanical signs, adaptation and
1772 clinical applications. *J Physiol.* 537:333-45.

1773 Proske U, Morgan DL (2005) Damage to skeletal muscle from eccentric exercise. *Exerc Sport Sci Rev.* 33:98-104.

1774 Pumpa KL, Fallon KE, Bensoussan A, Papalia S (2013) The effects of panax notoginseng on delayed onset muscle
1775 soreness and muscle damage in well-trained males: a double blind randomised controlled trial. *Complement Ther Med.*
1776 21:131-40.

1777 Qamar MM, Javed MS, Dogar MZUH, Basharat A (2019) Beat the exercise-induced damage. *J Pak Med Assoc.* 69:1682-
1778 1686.

1779 Quinlan R, Hill JA (2019) The efficacy of tart cherry juice in aiding recovery after intermittent exercise. *Int J Sports*
1780 *Physiol Perform.* 15:1-7.

1781 Ra SG, Akazawa N, Choi Y, Matsubara T, oikawa S, Kumagai H, Tanahashi K, Ohmori H, Maeda S (2015) Taurine
1782 supplementation reduces eccentric exercise-induced delayed onset muscle soreness in young men. *Adv Exp Med Biol.*
1783 803:765-72.

1784 Ra SG, Miyazaki T, Ishikura K, Nagayama H, Komine S, Nakata Y, Maeda S, Matsuzaki Y, Ohmori H (2013) Combined
1785 effect of branched-chain amino acids and taurine supplementation on delayed onset muscle soreness and muscle damage
1786 in high-intensity eccentric exercise. *J Int Soc Sports Nutr.* 10:51.

1787 Rahimi MH, Shab-Bidar S, Mollahosseini M, Djafarian K (2017) Branched-chain amino acid supplementation and
1788 exercise-induced muscle damage in exercise recovery: a meta-analysis of randomized clinical trials. *Nutrients.* 42:30-36.

1789 Rahimi R (2011) Creatine supplementation decreases oxidative DNA damage and lipid peroxidation induced by a single
1790 bout of resistance exercise. *J Strength Cond res.* 25:3448-55.

1791 Rahmani NF, Farzaneh E, Damirchi A, Shamsi Majlan A (2013) Effect of l-glutamine supplementation on
1792 electromyographic activity of the quadriceps muscle injured by eccentric exercise. *Iran J Basic Med Sci.* 16:808-812.

1793 Ramos-Campo DJ, Ávila-Gandía V, López-Román FJ, Minárrro J, Contreras C, Soto-Méndez F, Domingo Pedrol JC,
1794 Luque-Rubia AJ (2020) Supplementation of re-esterified docosahexaenoic and eicosapentaenoic acids reduce
1795 inflammatory and muscle damage markers after exercise in endurance athletes: a randomized, controlled crossover trial.
1796 *Nutrients.* 12(2). pii:E719.

1797 Ranchordas MK, Rogerson D, Soltani H, Costello JT (2017) Antioxidants for preventing and reducing muscle soreness
1798 after exercise. *Cochrane Database syst Rev.* 12:CD009789.

1799 Rankin P, Stevenson E, Cockburn E (2015) The effect of milk on the attenuation of exercise-induced muscle damage in
1800 males and females. *Eur J Appl Physiol* 115:1245-61.

1801 Rawson ES, Conti MP, Miles MP (2007) Creatine supplementation does not reduce muscle damage or enhance recovery
1802 from resistance exercise. *J Strength Cond Res.* 21:1208-13.

1803 Rawson ES, Gunn B, Clarkson PM (2001) The effects of creatine supplementation on exercise-induced muscle damage.
1804 *J Strength Cond Res.* 15:178-84.

1805 Rawson ES, Miles MP, Larson-Meyer DE (2018) Dietary supplements for health, adaptation, and recovery in athletes.
1806 *Int J Sport Nutr Exerc Metab* 28:188-199.

1807 Righi NC, Schuch FB, De Nardi AT, Pippi CM, de Almeida Righi G, Puntel GO, da Silva AMV, Signori LU (2020)
1808 Effects of vitamin C on oxidative stress, inflammation, muscle soreness, and strength following acute exercise: meta-
1809 analyses of randomized clinical trials. *Eur J Nutr.* [Epub ahead of print].

1810 Ring SM, Dannecker EA, Peterson CA (2010) Vitamin D status is not associated with outcomes of experimentally-
1811 induced muscle weakness and pain in young, healthy volunteers. *J Nutr Metab.* 2010:674240.

1812 Robinson MJ, Burd NA, Breen L, Rerecich T, Yang Y, Hector AJ, Baker SK, Phillips SM (2013) Dose-dependent
1813 responses of myofibrillar protein synthesis with beef ingestion are enhanced with resistance exercise in middle-aged men.
1814 *Appl Physiol Nutr Metab.* 38:120-5.

1815 Rosene J, Matthews T, Ryan C, Belmore K, Bergsten A, Blaisdell J, Gaylord J, Love R, Marrone M, Ward M, Wilson E
1816 (2009) Short and longer-term effects of creatine supplementation on exercise induced muscle damage. *J Sports Sci Med*
1817 8:89-96.

1818 Sacheck JM, Milbury PE, Cannon JG, Roubenoff R, Blumberg JB (2003) Effect of vitamin E and eccentric exercise on
1819 selected biomarkers of oxidative stress in young and elderly men. *Free Radic Biol Med.* 34:1575-1588.

1820 Sadowska-Krępa E, Kłapcińska B, Podgórski T, Szade B, Tyl K, Hadzik A (2015) effects of supplementation with acai
1821 (*Euterpe oleracea* Mart.) berry-based juice blend on the blood antioxidant defence capacity and lipid profile in junior
1822 hurdlers. A pilot study. *Biol Sport.* 32:161-8.

1823 Samaras A, Tsarouhas K, Paschalidis E, Giamouzis G, Triposkiadis F, Tsitsimpikou C, Becker AT, Goutzourelas N,
1824 Kouretas D (2014) Effect of a special carbohydrate-protein bar and tomato juice supplementation on oxidative stress
1825 markers and vascular endothelial dynamics in ultra-marathon runners. *Food Chem Toxicol.* 69:231-6.

1826 Sandur SK, Pandey MK, Sung B, Ahn KS, Murakami A, Sethi G, Limtrakul P, Badmaev V, Aggarwal BB (2007)
1827 Curcumin, demethoxycurcumin, bisdemethoxycurcumin, tetrahydrocurcumin and turmerones differentially regulate anti-
1828 inflammatory and anti-proliferative responses through a ROS-independent mechanism. *Carcinogenesis.* 28:1765-73.

1829 Santos RV, Bassit RA, Caperuto EC, Costa Rosa LF (2004) The effect of creatine supplementation upon inflammatory
1830 and muscle soreness markers after a 30Km race. *Life Sci.* 75:1917-24.

1831 Schaffer S and Kim HW (2018) Effects and mechanisms of taurine as a therapeutic agent. *Biomol Ther (Seoul)* 26:225-
1832 241.

1833 Sciberras JN, Galloway SD, Fenech A, Grech G, Farrugia C, Duca D, Mifsud J (2015) The effect of turmeric (Curcumin)
1834 supplementation on cytokine and inflammatory marker responses following 2 hours of endurance cycling. *J Int Soc Sports*
1835 *Nutr.* 12:5.

1836 Seeram NP, Momin RA, Nair MG and Bourquin LD (2001) Cyclooxygenase inhibitory and antioxidant cyaniding
1837 glycosides in cherries and berries. *Phytomedicine.* 8:362-369.

1838 Sellami M, Slimeni O, Porkrywka A, Kuvačić G, Hayes DL, Milic M, Padulo J (2018) Herbal medicine for sports: a
1839 review. *J Int Soc Sports Nutr.* 15:14.

1840 Shafat A, Butler P, Jensen RL, Donnelly AE (2004) Effects of dietary supplementation with vitamin C and E on muscle
1841 function during and after eccentric contractions in humans. *Eur J Appl Physiol.* 93:196-202.

1842 Shanely RA, Nieman DC, Knab AM, Gillitt ND, Meaney MP, Jin F, Sha W, Cialdella-Kam L (2014) Influence of
1843 vitamin D mushroom powder supplementation on exercise-induced muscle damage in vitamin D insufficient high
1844 school athletes. *J Sports Sci.* 32:670-9.

1845 Shanely RA, Nieman DC, Perkins-Veazie P, Henson DA, Meaney MP, Knab AM, Cialdell-Kam L (2016) Comparison
1846 of watermelon and carbohydrate beverage on exercise-induced alterations in systemic inflammation, immune
1847 dysfunction, and plasma antioxidant capacity. *Nutrients.* 8(8).

1848 Shaw G, Lee-Barthel A, Ross ML, Wang B, Baar K (2017) Vitamin C-enriched gelatin supplementation before
1849 intermittent activity augments collagen synthesis. *Am J Clin Nutr.* 105:136-143.

1850 Shenoy S, Dhawan M and Sandhu JS (2016) Four weeks of supplementation with isolated soy protein attenuates exercise-
1851 induced muscle damage and enhances muscle recovery in well trained athletes: a randomized trial. *Asian J Sports Med.*
1852 7:e33528.

1853 Shing CM, Chong S, Driller MW, Fell JW (2016) Acute protease supplementation effects on muscle damage and recovery
1854 across consecutive days of cycle racing. *Eur J Sport Sci.* 16:206-12.

1855 Shing CM, Peake JM, Lim CL, Briskey D, Walsh NP, Fortes MB, Ahuja KD, Vitetta L (2014) Effects of probiotics
1856 supplementation on gastrointestinal permeability, inflammation and exercise performance in the heat. *Eur J Appl Physiol.*
1857 114:93-103.

1858 Shoba G, Joy D, Joseph T, Majeed M, Rajendran R, Srinivas PS (1998) Influence of piperine on the pharmacokinetics of
1859 curcumin in animals and human volunteers. *Planta Med.* 64:353-6.

1860 Shuler FD, Wingate MK, Moore GH, Giangarra C (2012) Sports health benefits of vitamin D. *Sports Health.* 4:496-501.

1861 Siervo M, Scialò F, Shannon OM, Stephan BCM, Ashor AW (2018) Does dietary nitrate say NO to cardiovascular ageing?
1862 Current evidence and implications for research. *Proc Nutr Soc.* 77:112-123.

1863 Sousa M, Teixeira VH, Soares J (2014) Dietary strategies to recover from exercise-induced muscle damage. *Int J Food*
1864 *Sci Nutr.* 65:151-63.

1865 Spiering BA, Kraemer WJ, Hatfield DL, Vingren JL, Fragala MS, Ho JY, Thomas GA, Häkkinen K, Volek JS (2008)
1866 Effects of l-carnitine l-tartrate supplementation on muscle oxygenation responses to resistance exercise. *J Strength Cond.*
1867 *Res.* 22:1130-5.

1868 Spiering BA, Kraemer WJ, Vingren JL, Hatfield DL, Fragala MS, Ho JY, Maresh CM, Anderson JM, Volek JS (2007)
1869 Responses of criterion variables to different supplemental doses of L-carnitine L-tartrate. *J Strength Cond Res.* 21:259-
1870 64.

1871 Street B, Byrne C, Eston R (2011) Glutamine supplementation in recovery from eccentric exercise attenuates strength
1872 loss and muscle soreness. *J Exerc Sci Fit.* 2:116-122.

1873 Su QS, Tian Y, Zhang JG, Zhang H (2008) Effects of allicin supplementation on plasma markers of exercise-induced
1874 muscle damage, IL-6 and antioxidant capacity. *Eur J Appl Physiol.* 103:275-83.

1875 Sugita M, Kapoor MP, Nishimura A, Okubo T (2016) Influence of green tea catechins on oxidative stress metabolites at
1876 rest and during exercise in healthy humans. *Nutrition.* 32:321-331.

1877 Takagi R, Ogasawara R, Tsutaki A, Nakazato K, Ishii N (2016) Regional adaptation of collagen in skeletal muscle to
1878 repeated of strenuous eccentric exercise. *Pflugers Arch.* 468:1565-72.

1879 Tanabe Y, Chino K, Ohnishi T, ozawa H, Sagayama H, Maeda S, Takahashi H (2019b) Effects of oral curcumin ingested
1880 before or after eccentric exercise on markers of muscle damage. *Scand J Med Sci Sports.* 29:524-534.

1881 Tanabe Y, Chino K, Sagayama H, Lee HJ, Ozawa H, Maeda S, Takahashi H (2019a) Effective timing of curcumin
1882 ingestion to attenuate eccentric exercise-induced muscle soreness in men. *J Nutr sci Vitaminol* 65:82-89.

1883 Tanabe Y, Maeda S, Akazawa N, Zempo-Miyaki A, Choi Y, Ra SG, Imaizumi A, Otsuka Y, Nosaka K (2015) Attenuation
1884 of indirect markers of eccentric exercise-induced muscle damage by curcumin. *Eur J Appl Physiol.* 115:1949-57.

1885 Tarazona-Diaz MP, Alacid F, Carrasco M, Martinez I, Aguayo E (2013) Watermelon juice: potential functional drink for
1886 sore muscle relief in athletes. *J Agric Food Chem* 61:7522-8.

1887 Tartibian B, Maleki BH, Abbasi A (2009) The effects of ingestion of omega-3 fatty acids on perceived pain and
1888 external symptoms of delayed onset muscle soreness in untrained men. *Clin J Sport Med.* 19:115-9.

1889 Tartibian B, Maleki BH, Abbasi A (2011) omega-3 fatty acids supplementation attenuates inflammatory markers after
1890 eccentric exercise in untrained men. *Clin J Sport Med.* 21:131-7.

1891 Teixeira FJ, Matias CN, Monteiro CP, Valamatos MJ, Reis JF, Morton RW, Alves F, Sardinha LB, Phillips SM (2019)
1892 Leucine metabolites do not attenuate training-induced inflammation in young resistance trained men. *J Sports Sci.*
1893 37:2037-2044.

1894 Thompson D, Bailey DM, Hill J, Hurst T, Powell JR, Williams C (2004) Prolonged vitamin C supplementation and
1895 recovery from eccentric exercise. *Eur J Appl Physiol.* 92:133-8.

1896 Thompson D, Williams C, Kingsley M, Nicholas CW, Lakomy HK, McArdle F, Jackson MJ (2001) Muscle soreness and
1897 damage parameters after prolonged intermittent shuttle-running following acute vitamin C supplementation. *Int J Sports*
1898 *Med.* 22:68-75.

1899 Tidball JG, Dorshkind K, Wehling-Henricks M (2014) Shared signaling systems in myeloid cell-mediated muscle
1900 regeneration. *Development*. 141:1184-96.

1901 Tinsley GM, Gann JJ, Huber SR, Andre TL, La Bounty PM, Bowden RG, Gordon PM, Grandjean PW (2017) Effects
1902 of fish oil supplementation on postresistance exercise muscle soreness. *J Diet Suppl*. 14:89-100.

1903 Tipton KD, Borsheim E, Wolf SE, Sanford AP, Wolfe RR (2003) Acute response of net muscle protein balance reflects
1904 24-h balance after exercise and amino acid ingestion. *Am J Physiol Endocrinol Metab*. 284:E76-89.

1905 Tipton KD (2015) Nutritional support for exercise-induced injuries. *Sports Med*. 1:S93-104.

1906 Torregrosa-García A, Ávila-Gandía V, Luque-Rubia AJ, Abellán-Ruiz MS, Querol-Calderón M and López-Román J
1907 (2019) Pomegranate extract improves maximal performance of trained cyclists after an exhausting endurance trial:a
1908 randomised controlled trial. *Nutrients*. 11:721.

1909 Trombold JR, Barnes JN, Critchley L, Coyle EF (2010) Ellagitannin consumption improves strength recovery 2-3 d after
1910 eccentric exercise. *Med Sci Sports Exerc*. 42:493-8.

1911 Trombold JR, Reinfeld AS, Casler JR, Coyle EF (2011) The effect of pomegranate juice supplementation on strength and
1912 soreness after eccentric exercise. *J Strength Cond. Res*. 25:1782-8.

1913 Trommelen J, Betz MW, van Loon LJC (2019) The muscle protein synthetic response to meal ingestion following
1914 resistance-type exercise. *Sports Med*. 49:185-197.

1915 Tsitsimpikou C, Kioukia-Fougia N, Tsarouhas K, Stamatopoulos P, Rentoukas E, Koudounakos A, Papalexis P,
1916 Liesivuori J, Jamurtas A (2013) Administration of tomato juice ameliorates lactate dehydrogenase and creatinine kinase
1917 responses to anaerobic training. *Food Chem Toxicol*. 61:9-13.

1918 Tsuchiya Y, Yanagimoto K, Nakazato K, Hayamizu K, Ochi E (2016) Eicosapentaenoic and docosahexaenoic acids-
1919 rich fish oil supplementation attenuates strength loss and limited joint range of motion after eccentric contractions: a
1920 randomized, double-blind, placebo-controlled, parallel-group trial. *Eur J Appl Physiol*. 116:1179-88.

1921 Tsuchiya Y, Yanagimoto K, Ueda H, Ochi E (2019) Supplementation of eicosapentaenoic acid-rich fish oil attenuates
1922 muscle stiffness after eccentric contractions of human elbow flexors. *J Int Soc Sports Nutr*. 16:19.

1923 Udani JK, Singh BB, Singh VJ, Sandoval E (2009) BounceBack capsules for reduction of DOMS after eccentric exercise:
1924 a randomized, double-blind, placebo-controlled, crossover pilot study. *J int Soc Sports Nutr*. 6:14.

1925 Vaile JM, Gill ND, Blazeovich AJ (2007) The effect of contrast water therapy on symptoms of delayed onset muscle
1926 soreness. *J Strength Cond Res*. 21:697-702.

1927 van der Merwe M, Bloomer RJ (2016) The influence of methylsulfonylmethane on inflammation-associated cytokine
1928 release before and following strenuous exercise. *J Sports Med (Hindawi Publ Corp)*. 7498359.

1929 van Someren KA, Edwards AJ, Howatson G (2005) Supplementation with β -hydroxy β -methylbutyrate (HMB) and alpha-
1930 ketoisocaproic acid (KIC) reduces signs and symptoms of exercise-induced muscle damage in man, *Int J Sport Nutr.*
1931 *Exerc Metab.* 15:413-24.

1932 Veggi KFT, Machado M, Koch AJ, Santana SC, Oliveira SS, Stec MJ (2013) Oral creatine supplementation augments
1933 the repeated bout effect. *Int J Sport Nutr Exerc Metab* 23:378-87.

1934 Verburg E, Murphy RM, Stephenson DG, Lamb GD (2005) Disruption of excitation-contraction coupling and titin by
1935 endogenous Ca^{2+} -activated proteases in toad muscle fibres. *J Physiol* 564:775-90.

1936 Vierck JL, Icenogge DL, Bucci L, Dodson MV (2003) The effects of ergogenic compounds on myogenic satellite cells.
1937 *Med Sci Sports exerc.* 35:769-76.

1938 Vimercatti NS, Zovico PVC, Carvalho AS, Barreto JG, Machado M (2008) Two doses of caffeine do not increase the
1939 risk of exercise-induced muscle damage or leucocytosis. *Physical Education and Sport* 52:96-99

1940 Vitale KC, Hueglin S, Broad E (2017) Tart cherry juice in athletes: a literature review and commentary. *Curr Sports*
1941 *Med Rep.* 16:230-239.

1942 Volek JS, Kraemer WJ, Rubin MR, Gómez AL, Ratamess NA, Gaynor P (2002) L-Carnitine L-tartrate supplementation
1943 favorably affects markers of recovery from exercise stress. *Am J Physiol Endocrinol Metab.* 282:E474-82.

1944 Waldron M, Whelan K, Jeffries O, Burt D, Howe L, Patterson SD (2017) The effects of acute branched-chain amino acid
1945 supplementation on recovery from a single bout of hypertrophy exercise in resistance-trained athletes. *Appl physiol Nutr*
1946 *Metab.* 42:630-636.

1947 Wang H, Nair MG, Strasburg GM, Chang YC, Booren AM, Gray JI and DeWitt DI (1999). Antioxidant and anti-
1948 inflammatory activities of anthocyanins and their aglycon, cyanidin, from tart cherries. *J Nat Prod.* 62:294-296.

1949 Warren GL, Ingalls CP, Lowe DA, Armstrong RB (2001) Excitation-contraction uncoupling: major role in contraction-
1950 induced muscle injury. *Exerc Sport Sci Rev.* 29:82-7.

1951 Warren GL, Lowe DA, Armstrong RB (1999) Measurement tools used in the study of eccentric contraction-induced
1952 injury. *Sports Med.* 27:43-59.

1953 Willis KS, Peterson NJ, Larson-Meyer DE (2008). Should we be concerned about the vitamin D status of athletes? *Int J*
1954 *Sport Nutr Exerc Metab* 18:204-24.

1955 Willoughby DS, Rosene JM (2003) Effects of oral creatine and resistance training on myogenic regulatory factor
1956 expression. *Med sci Sports Exerc.* 35:923-9.

1957 Wilson JM, Kim JS, Lee SR, Rathmacher JA, Dalmau B, Kingsley JD, Koch H, Manninen AH, Saadat R, Panton LB
1958 (2009) Acute and timing effects of β -hydroxy β -methylbutyrate (HMB) on indirect markers of skeletal muscle damage.
1959 *Nutr. Metab (Lond).* 6:6.

1960 Wilson JM, Lowery RP, Joy JM, Andersen JC, Wilson SM, Stout JR, Duncan N, Fuller JC, Baier SM, Naimo MA,
1961 Rathmacher J (2014) The effects of 12 weeks of β -hydroxy β -methylbutyrate free acid supplementation on muscle mass,
1962 strength, and power in resistance-trained individuals: a randomized, double-blind, placebo-controlled study. *Eur J Appl*
1963 *Physiol.* 114:1217-27.

1964 Wilson JM, Lowery RP, Joy JM, Walters JA, Baier SM, Fuller JC Jr, Stout JR, Norton LE, Sikorski EM, Wilson SM,
1965 Duncan NM, Zanchi NE, Rathmacher J (2013) β -hydroxy β -methylbutyrate free acid reduces markers of exercise-induced
1966 muscle damage and improves recovery in resistance-trained men. *Br J Nutr.* 110:538-44.

1967 Wilson LR, Tripkovic L, Hart KH, Lanham-New SA (2017) Vitamin D deficiency as a public health issue: using
1968 vitamin D2 or vitamin D3 in future fortification strategies. *Proc Nutr Soc.* 76:392-399.

1969 Wilson PB (2015) Ginger (*Zingiber officinale*) as an analgesic and ergogenic aid in sport: a systemic review. *J Strength*
1970 *Cond Res.* 29:2980-95.

1971 Wilson PB, Fitzgerald JS, Rhodes S, Lundstrom CJ, Stacy J (2015) Effectiveness of ginger root (*Zingiber officinale*) on
1972 running-induced muscle soreness and function: a pilot study. *Int J of Athl Ther Train.* 20:44-50.

1973 Wojcik JR, Walber-Rankin J, Smith LL, Gwazdauskas FC (2001) Comparison of carbohydrate and milk-based beverages
1974 on muscle damage and glycogen following exercise. *Int J Sport Nutr exerc Metab.* 11:406-19.

1975 Wu X, Beecher GR, Holden JM, Haytowitz DB, Gebhardt SE, Prior RL (2004) Lipophilic and hydrophilic antioxidant
1976 capacities of common foods in the United States. *J Agric Food Chem.* 52:4026-37.

1977 Yang HH, Li XL, Zhang WG, Figueroa A, Chen LH, Qin LQ (2019) Effect of oral L-citrulline on brachial and aortic
1978 blood pressure defined by resting status: evidence from randomized controlled trials. *Nutr Metab (Lond).* 16:89.

1979 Yarizadh H, Shab-Bidar S, Zamani B, Vanani AN, Baharloo H, Djafarian K (2020) The effect of L-Carnitine
1980 supplementation on exercise-induced muscle damage: a systematic review and meta-analysis of randomized clinical trials.
1981 *J Am Coll Nutr.* 10:1-12.

1982 Yu JG, Liu JX, Carlsson L, Thornell LE, Stål PS (2013) Re-evaluation of sarcolemma injury and muscle swelling in
1983 human skeletal muscles after eccentric exercise. *PloS One.* 8:e62056