

Northumbria Research Link

Citation: Wang, Zhao, Mao, Yaping, Das, Kinkar Chandra and Shang, Yilun (2020) Nordhaus-Gaddum-Type Results for the Steiner Gutman Index of Graphs. *Symmetry*, 12 (10). p. 1711. ISSN 2073-8994

Published by: MDPI

URL: <https://doi.org/10.3390/sym12101711> <<https://doi.org/10.3390/sym12101711>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/44695/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

Article

Nordhaus–Gaddum-Type Results for the Steiner Gutman Index of Graphs

Zhao Wang ¹, Yaping Mao ^{2,3} , Kinkar Chandra Das ^{4,*}  and Yilun Shang ^{5,*} ¹ College of Science, China Jiliang University, Hangzhou 310018, Zhejiang, China; wangzhao@mail.bnu.edu.cn² Department of Mathematics, Qinghai Normal University, Xining 810008, Qinghai, China; maoyaping@ymail.com³ Center for Mathematics and Interdisciplinary Sciences of Qinghai Province, Xining 810008, Qinghai, China⁴ Department of Mathematics, Sungkyunkwan University, Suwon 16419, Korea⁵ Department of Computer and Information Sciences, Northumbria University, Newcastle NE1 8ST, UK

* Correspondence: kinkardas2003@googlemail.com (K.C.D.); yilun.shang@northumbria.ac.uk (Y.S.)

Received: 18 September 2020; Accepted: 12 October 2020; Published: 16 October 2020



Abstract: Building upon the notion of the Gutman index $SGut(G)$, Mao and Das recently introduced the Steiner Gutman index by incorporating Steiner distance for a connected graph G . The Steiner Gutman k -index $SGut_k(G)$ of G is defined by $SGut_k(G) = \sum_{S \subseteq V(G), |S|=k} (\prod_{v \in S} deg_G(v)) d_G(S)$, in which $d_G(S)$ is the Steiner distance of S and $deg_G(v)$ is the degree of v in G . In this paper, we derive new sharp upper and lower bounds on $SGut_k$, and then investigate the Nordhaus–Gaddum-type results for the parameter $SGut_k$. We obtain sharp upper and lower bounds of $SGut_k(G) + SGut_k(\overline{G})$ and $SGut_k(G) \cdot SGut_k(\overline{G})$ for a connected graph G of order n , m edges, maximum degree Δ and minimum degree δ .

Keywords: distance; Steiner distance; Gutman index; Steiner Gutman k -index**MSC:** 05C05; 05C12; 05C35

1. Introduction

We consider simple, undirected graphs in this paper. For the standard theoretical graph terminology and notation not defined here, follow [1]. For a graph G , let $V(G)$ and $E(G)$ represent its sets of vertices and edges, respectively. Let $|E(G)| = m$ be the size of G . The complement of G is conventionally denoted by \overline{G} . For a vertex $v \in V(G)$, $deg_G(v)$ is the degree of v . The maximum and minimum degrees are, respectively, denoted by Δ and δ . Like degrees, distance is a fundamental concept of graph theory [2]. For two vertices $u, v \in V(G)$ with connected G , the distance $d(u, v) = d_G(u, v)$ between these two vertices is defined as the length of a shortest path connecting them. An excellent survey paper on this subject can be found in [3].

The above classical graph distance was extended by Chartrand et al. in 1989 to the Steiner distance, which since then has become an essential concept of graph theory. Given a graph $G(V, E)$ and a vertex set $S \subseteq V(G)$ containing no less than two vertices, an S -Steiner tree (or an S -tree, a Steiner tree connecting S) is defined as a subgraph $T(V', E')$ of G , which is a subtree satisfying $S \subseteq V'$. If G is connected with order no less than 2 and $S \subseteq V$ is nonempty, the Steiner distance $d(S)$ among the vertices of S (sometimes simply put as the distance of S) is the minimum size of connected subgraph whose vertex sets contain the set S . Clearly, for a connected subgraph $H \subseteq G$ with $S \subseteq V(H)$ and $|E(H)| = d(S)$, H is a tree. When T is subtree of G , we have $d(S) = \min\{|E(T)|, S \subseteq V(T)\}$. For $S = \{u, v\}$, $d(S) = d(u, v)$ reduces to the classical distance between the two vertices u and v . Another basic observation is that if $|S| = k$, $d(S) \geq k - 1$. For more results regarding varied properties of the Steiner distance, we refer to the reader to [3–8].

In [9], Li et al. generalized the concept of Wiener index through incorporating the Steiner distance. The Steiner k -Wiener index $SW_k(G)$ of G is defined by

$$SW_k(G) = \sum_{\substack{S \subseteq V(G) \\ |S|=k}} d(S).$$

For $k = 2$, it is easy to see the Steiner Wiener index coincides with the ordinary Wiener index. The interesting range of the Steiner k -Wiener index SW_k resides in $2 \leq k \leq n - 1$, and the two trivial cases give $SW_1(G) = 0$ and $SW_n(G) = n - 1$.

Gutman [10] studied the Steiner degree distance, which is a generalization of ordinary degree distance. Formally, the k -center Steiner degree distance $SDD_k(G)$ of G is given as

$$SDD_k(G) = \sum_{\substack{S \subseteq V(G) \\ |S|=k}} \left(\sum_{v \in S} \deg_G(v) \right) d_G(S).$$

The Gutman index of a connected graph G is defined as

$$Gut(G) = \sum_{u, v \in V(G)} \deg_G(u) \deg_G(v) d_G(u, v).$$

The Gutman index of graphs attracted attention very recently. For its basic properties and applications, including various lower and upper bounds, see [11–13] and the references cited therein. Recently, Mao and Das [14] further extended the concept of the Gutman index by incorporating Steiner distance and considering the weights as multiplications of degrees. The Steiner k -Gutman index $SGut_k(G)$ of G is defined by

$$SGut_k(G) = \sum_{\substack{S \subseteq V(G) \\ |S|=k}} \left(\prod_{v \in S} \deg_G(v) \right) d_G(S).$$

Note that this index is a natural generalization of the classical Gutman index—in particular, for $k = 2$, $SGut_k(G) = Gut(G)$. This is the reason the product of the degrees comes to the definition of Steiner k -Gutman index. The weighting of multiplication of degree or expected degree has also been extensively explored in, for example, the field of random graphs [15,16] and proves to be very prolific. For more results on Steiner Wiener index, Steiner degree distance and Steiner Gutman index, we refer to the reader to [9,10,14,17–19].

For a given a graph parameter $f(G)$ and a positive integer n , the well-known Nordhaus–Gaddum problem is to determine sharp bounds for: (1) $f(G) + f(\overline{G})$ and (2) $f(G) \cdot f(\overline{G})$ over the class of connected graph G , with order n , m edges, maximum degree Δ and minimum degree δ characterizing the extremal graphs. Many Nordhaus–Gaddum type relations have attracted considerable attention in graph theory. Comprehensive results regarding this topic can be found in e.g., [20–24].

In Section 2, we obtain sharp upper and lower bounds on $SGut_k$ of graph G . In Section 3, we obtain sharp upper and lower bounds of $SGut_k(G) + SGut_k(\overline{G})$ and $SGut_k(G) \cdot SGut_k(\overline{G})$ for a connected graph G in terms of n , m , maximum degree Δ and minimum degree δ .

2. Sharp Bounds for the Steiner Gutman Index

In [14], the following results have been obtained:

Lemma 1 ([14]). *Let K_n , S_n and P_n be the complete graph, star graph and path graph of order n , respectively, and let k be an integer such that $2 \leq k \leq n$. Then*

- (1) $SGut_k(K_n) = \binom{n}{k}(n-1)^n(k-1);$
- (2) $SGut_k(S_n) = (kn-2k+1)\binom{n-1}{k-1};$
- (3) $SGut_k(P_n) = 2^k(k-1)\binom{n}{k+1}.$

For connected graph G of order n with m edges, the authors in [14] derived the following upper and lower bounds on $SGut_k(G)$.

Lemma 2 ([14]). *Let G be a connected graph of order n with m edges, and let k be an integer with $2 \leq k \leq n$. Then*

$$(n-1) \left(\frac{2m}{k}\right)^k \binom{n-1}{k-1} \geq SGut_k(G) \geq \begin{cases} 2m(k-1)\binom{n-1}{k-1} & \text{if } \delta \geq 2 \\ (k-1)\binom{n}{k} & \text{if } \delta = 1. \end{cases}$$

We now give lower and upper bounds for $SGut_k(G)$ in terms of n , m , maximum degree Δ and minimum degree δ :

Proposition 1. *Let G be a connected graph of order $n \geq 3$ with m edges and maximum degree Δ , minimum degree δ . Additionally, let k be an integer with $2 \leq k \leq n$. Then*

$$2m(n-1) \binom{n-1}{k-1} \frac{\Delta^{k-1}}{k} \geq SGut_k(G) \geq \begin{cases} 2m(k-1)\binom{n-1}{k-1} \frac{\delta^{k-1}}{k} & \text{if } \delta \geq 2 \\ k\binom{p}{k} + 2^q(k-1) \left[\binom{n}{k} - \binom{p}{k}\right] & \text{if } \delta = 1, \end{cases}$$

where p is the number of pendant vertices in G , and $q = \max\{k-p, 1\}$. The equality of upper bound holds if and only if G is a regular graph with $k = n$. The equality of lower bound holds if and only if G is a regular $(n-k+1)$ -connected graph of order n ($\delta \geq 2$), or $G \cong P_n$ and $k = n > 3$ ($\delta = 1$), or $G \cong P_3$ and $k = 2$ ($\delta = 1$).

Proof. Upper bound: For any $S \subseteq V(G)$ and $|S| = k$, we have $k-1 \leq d_G(S) \leq n-1$, and hence

$$(k-1) \sum_{\substack{S \subseteq V(G) \\ |S|=k}} \left(\prod_{v \in S} deg_G(v) \right) \leq SGut_k(G) \leq (n-1) \sum_{\substack{S \subseteq V(G) \\ |S|=k}} \left(\prod_{v \in S} deg_G(v) \right). \tag{1}$$

Let

$$M = \sum_{\substack{S \subseteq V(G) \\ |S|=k}} \left(\prod_{v \in S} deg_G(v) \right) = \sum_{\{v_1, v_2, \dots, v_k\} \subseteq V(G)} deg_G(v_1) deg_G(v_2) \cdots deg_G(v_k).$$

and

$$N = \sum_{\{v_1, v_2, \dots, v_k\} \subseteq V(G)} [deg_G(v_1) + deg_G(v_2) + \cdots + deg_G(v_k)].$$

We first prove the upper bound. Without loss of generality, we can assume that $deg_G(v_1) \leq deg_G(v_2) \leq \dots \leq deg_G(v_k)$. Since

$$deg_G(v_1) deg_G(v_2) \cdots deg_G(v_k) \leq \Delta^{k-1} deg_G(v_k) \tag{2}$$

$$\leq \frac{\Delta^{k-1}}{k} (deg_G(v_1) + deg_G(v_2) + \cdots + deg_G(v_k)), \tag{3}$$

it follows that

$$\begin{aligned}
 M &= \sum_{\{v_1, v_2, \dots, v_k\} \subseteq V(G)} \deg_G(v_1) \deg_G(v_2) \dots \deg_G(v_k) \\
 &\leq \frac{\Delta^{k-1}}{k} \sum_{\{v_1, v_2, \dots, v_k\} \subseteq V(G)} [\deg_G(v_1) + \deg_G(v_2) + \dots + \deg_G(v_k)] \\
 &\leq \frac{\Delta^{k-1}}{k} N.
 \end{aligned}$$

For each $v \in V(G)$, there are $\binom{n-1}{k-1}$ k -subsets in G such that each of them contains v . The contribution of vertex v is exactly $\binom{n-1}{k-1} \deg_G(v)$. From the arbitrariness of v , we have

$$N = \binom{n-1}{k-1} \sum_{v \in V(G)} \deg_G(v) = 2m \binom{n-1}{k-1},$$

and hence

$$\text{SGut}_k(G) \leq (n-1)M \leq (n-1) \frac{\Delta^{k-1}}{k} N = 2m(n-1) \binom{n-1}{k-1} \frac{\Delta^{k-1}}{k}. \tag{4}$$

Suppose that the left equality holds. Then all the inequalities in the above must be equalities. From the equality in (3), one can easily see that G is a regular graph. From the equality in (4), we have $d(S) = n - 1$ for any $S \subseteq V(G)$, $|S| = k$. Since G is connected, then there exists an $S \subseteq V(G)$ such that $|d_G(S)| = k - 1$. If $k \leq n - 1$, then one can easily see that the upper bound is strict as $|d_G(S)| = k - 1 \leq n - 2$ for some S . Otherwise, $k = n$. Since G is connected, we have $|d_G(S)| = n - 1$ for any $S \subseteq V(G)$. Hence G is a regular graph with $k = n$.

Conversely, one can see easily that the left equality holds for regular graph with $k = n$.

Lower bound: Without loss of generality, we can assume that $\deg_G(v_1) \leq \deg_G(v_2) \leq \dots \leq \deg_G(v_k)$. First we assume that $\delta \geq 2$. Then

$$\begin{aligned}
 \deg_G(v_1) \deg_G(v_2) \dots \deg_G(v_k) &\geq \delta^{k-1} \deg_G(v_k) \\
 &\geq \frac{\delta^{k-1}}{k} (\deg_G(v_1) + \deg_G(v_2) + \dots + \deg_G(v_k)),
 \end{aligned} \tag{5}$$

since $\deg_G(v_1) \leq \deg_G(v_2) \leq \dots \leq \deg_G(v_k)$. Furthermore, we have

$$\text{SGut}_k(G) \geq (k-1) \sum_{\{v_1, v_2, \dots, v_k\} \subseteq V(G)} \deg_G(v_1) \deg_G(v_2) \dots \deg_G(v_k) \tag{6}$$

$$\begin{aligned}
 &\geq (k-1) \frac{\delta^{k-1}}{k} \sum_{\{v_1, v_2, \dots, v_k\} \subseteq V(G)} [\deg_G(v_1) + \deg_G(v_2) + \dots + \deg_G(v_k)] \tag{7} \\
 &= (k-1) \frac{\delta^{k-1}}{k} N \\
 &= 2m(k-1) \binom{n-1}{k-1} \frac{\delta^{k-1}}{k}.
 \end{aligned}$$

Next we assume that $\delta = 1$. If $\deg_G(v_1) = \deg_G(v_2) = \dots = \deg_G(v_k) = 1$, then $d_G(S) \geq k$ and $\deg_G(v_1) \deg_G(v_2) \dots \deg_G(v_k) = 1$. If there exists some v_i such that $\deg_G(v_i) \geq 2$, then $d_G(S) \geq k - 1$ and $\deg_G(v_1) \deg_G(v_2) \dots \deg_G(v_k) \geq 2^{\max\{k-p, 1\}} = 2^q$, where $1 \leq i \leq k$. Therefore, we have

$$\text{SGut}_k(G) \geq k \sum_{\substack{\{v_1, v_2, \dots, v_k\} \subseteq V(G), \\ \text{deg}_G(v_1) = \text{deg}_G(v_2) = \dots = \text{deg}_G(v_k) = 1}} \text{deg}_G(v_1) \text{deg}_G(v_2) \dots \text{deg}_G(v_k) + (k-1) \sum_{\substack{\{v_1, v_2, \dots, v_k\} \subseteq V(G), \\ \text{some } \text{deg}_G(v_i) \geq 2}} \text{deg}_G(v_1) \text{deg}_G(v_2) \dots \text{deg}_G(v_k) \tag{8}$$

$$\geq k \binom{p}{k} + 2^q (k-1) \left[\binom{n}{k} - \binom{p}{k} \right]. \tag{9}$$

Suppose that the right equality holds. Then all the inequalities in the above must be equalities. Suppose that $\delta \geq 2$. From the equality in (6), $d_G(S) = k - 1$ for any $S \subseteq V(G)$ and $|S| = k$, that is, $G[S]$ is connected for any $S \subseteq V(G)$ and $|S| = k$, and hence G is $(n - k + 1)$ -connected. From the equality in (7), we have $\text{deg}_G(v_1) = \text{deg}_G(v_2) = \dots = \text{deg}_G(v_k)$ for any $S = \{v_1, v_2, \dots, v_k\} \subseteq V(G)$, and hence G is a regular graph. Thus, G is a regular $(n - k + 1)$ -connected graph of order n .

Next suppose that $\delta = 1$. From the equality in (9), we obtain $\text{deg}_G(v_i) = 1$ or $\text{deg}_G(v_i) = 2$ for any vertex $v_i \in V(G)$. Since G is connected, $G \cong P_n$ and $p = 2$. If $k \geq 3$, then $q = k - p \geq 1$. In this case $d_G(S) = k - 1$ for any $S \subseteq V(G)$ and $|S| = k$. One can easily see that $G \cong P_n$ and $k = n > 3$ (otherwise, $d_G(S) > k - 1$ for some $S \subseteq V(G)$ as $q = k - p$). Otherwise, $k = p = 2$ and hence $q = 1$. In this case $G \cong P_3$ and $k = 2$.

Conversely, one can see easily that the equality holds on lower bound for a regular $(n - k + 1)$ -connected graph of order n ($\delta \geq 2$), or $G \cong P_n$ and $k = n > 3$ ($\delta = 1$), or $G \cong P_3$ and $k = 2$ ($\delta = 1$). □

Example 1. Let $G \cong K_n$ with $k = n$. Then

$$\text{SGut}_k(G) = (n - 1)^{n+1} = 2m(n - 1) \binom{n - 1}{k - 1} \frac{\Delta^{k-1}}{k}.$$

Let $G \cong K_n \setminus sK_2$ ($n = 2s$) with $k = 3$. Then G is a $n - 2$ regular graph of order n . Then

$$\text{SGut}_k(G) = 2(n - 2)^3 \binom{n}{3} = 2m(k - 1) \binom{n - 1}{k - 1} \frac{\delta^{k-1}}{k}.$$

Let $G \cong P_n$ with $k = n > 3$. Then

$$\text{SGut}_k(G) = 2^{n-2}(n - 1) = k \binom{p}{k} + 2^q (k - 1) \left[\binom{n}{k} - \binom{p}{k} \right] \text{ as } p = 2.$$

Let $G \cong P_n$ with $k = 2$. Then

$$\text{SGut}_k(G) = 6 = k \binom{p}{k} + 2^q (k - 1) \left[\binom{n}{k} - \binom{p}{k} \right] \text{ as } p = 2.$$

3. Nordhaus–Gaddum-Type Results on $\text{SGut}_k(G)$

We are now in a position to give the Nordhaus–Gaddum-type results on $\text{SGut}_k(G)$.

Theorem 1. Let G be a connected graph of order n with m edges, maximum degree Δ , minimum degree δ and a connected \bar{G} . Additionally, let k be an integer with $2 \leq k \leq n$. Then

(1)

$$\text{SGut}_k(G) + \text{SGut}_k(\bar{G}) \leq (n - 1)^2 \binom{n}{k} s_1^{k-1}$$

and

$$\text{SGut}_k(G) \cdot \text{SGut}_k(\overline{G}) \leq 2m(n^2 - n - 2m)(n - 1)^2 \binom{n - 1}{k - 1}^2 \frac{\Delta^{k-1} (n - \delta - 1)^{k-1}}{k^2},$$

where $s_1 = \max\{\Delta, n - \delta - 1\}$. Moreover, the upper bounds are sharp.

(2)

$$\begin{aligned} & \text{SGut}_k(G) + \text{SGut}_k(\overline{G}) \\ \geq & \begin{cases} (n - 1)(k - 1) \binom{n}{k} t_1^{k-1} & \text{if } \delta \geq 2, \Delta \leq n - 3 \\ 2m(k - 1) \binom{n-1}{k-1} \frac{\delta^{k-1}}{k} + k \binom{n}{k} & \text{if } \delta \geq 2, \Delta = n - 2 \\ k \binom{n}{k} + [n(n - 1) - 2m](k - 1) \binom{n-1}{k-1} \frac{(n - \Delta - 1)^{k-1}}{k} & \text{if } \delta = 1, \Delta \leq n - 3 \\ 2k \binom{n}{k} & \text{if } \delta = 1, \Delta = n - 2, \end{cases} \end{aligned}$$

where $t_1 = \min\{\delta, n - \Delta - 1\}$.

(3)

$$\begin{aligned} & \text{SGut}_k(G) \cdot \text{SGut}_k(\overline{G}) \\ \geq & \begin{cases} 2m(n^2 - n - 2m)(k - 1)^2 \binom{n-1}{k-1}^2 \frac{\delta^{k-1} (n - \Delta - 1)^{k-1}}{k^2} & \text{if } \delta \geq 2, \Delta \leq n - 3 \\ 2m(k - 1) \binom{n}{k} \binom{n-1}{k-1} \delta^{k-1} & \text{if } \delta \geq 2, \Delta = n - 2 \\ [n(n - 1) - 2m](k - 1) \binom{n}{k} \binom{n-1}{k-1} (n - \Delta - 1)^{k-1} & \text{if } \delta = 1, \Delta \leq n - 3 \\ k^2 \binom{n}{k}^2 & \text{if } \delta = 1, \Delta = n - 2. \end{cases} \end{aligned}$$

Proof. (1) From Proposition 1, we have

$$\text{SGut}_k(G) \leq 2m(n - 1) \binom{n - 1}{k - 1} \frac{\Delta^{k-1}}{k}$$

and

$$\text{SGut}_k(\overline{G}) \leq [n(n - 1) - 2m](n - 1) \binom{n - 1}{k - 1} \frac{(n - \delta - 1)^{k-1}}{k},$$

and hence

$$\text{SGut}_k(G) + \text{SGut}_k(\overline{G}) \leq (n - 1)^2 \binom{n}{k} s_1^{k-1}$$

and

$$\text{SGut}_k(G) \cdot \text{SGut}_k(\overline{G}) \leq 2m(n^2 - n - 2m)(n - 1)^2 \binom{n - 1}{k - 1}^2 \frac{\Delta^{k-1} (n - \delta - 1)^{k-1}}{k^2}.$$

(2) From Proposition 1, if $\delta \geq 2$ and $\Delta \leq n - 3$, then

$$\begin{aligned} & \text{SGut}_k(G) + \text{SGut}_k(\overline{G}) \\ \geq & 2m(k - 1) \binom{n - 1}{k - 1} \frac{\delta^{k-1}}{k} + [n(n - 1) - 2m](k - 1) \binom{n - 1}{k - 1} \frac{(n - \Delta - 1)^{k-1}}{k} \\ \geq & (n - 1)(k - 1) \binom{n}{k} t_1^{k-1}. \end{aligned}$$

If $\delta(G) \geq 2$ and $\Delta = n - 2$, then

$$\begin{aligned} & \text{SGut}_k(G) + \text{SGut}_k(\overline{G}) \\ & \geq 2m(k-1) \binom{n-1}{k-1} \frac{\delta^{k-1}}{k} + k \binom{p'}{k} + 2^{q'}(k-1) \left[\binom{n}{k} - \binom{p'}{k} \right] \\ & \geq 2m(k-1) \binom{n-1}{k-1} \frac{\delta^{k-1}}{k} + k \binom{p'}{k} + 2(k-1) \left[\binom{n}{k} - \binom{p'}{k} \right] \\ & \geq 2m(k-1) \binom{n-1}{k-1} \frac{\delta^{k-1}}{k} + k \binom{p'}{k} + k \left[\binom{n}{k} - \binom{p'}{k} \right] \\ & = 2m(k-1) \binom{n-1}{k-1} \frac{\delta^{k-1}}{k} + k \binom{n}{k}, \end{aligned}$$

where p' is the number of pendant vertices in G , and $q' = \max\{k - p', 1\}$.

If $\delta = 1$ and $\Delta \leq n - 3$, then

$$\begin{aligned} & \text{SGut}_k(G) + \text{SGut}_k(\overline{G}) \\ & \geq k \binom{p}{k} + 2^q(k-1) \left[\binom{n}{k} - \binom{p}{k} \right] + [n(n-1) - 2m](k-1) \binom{n-1}{k-1} \frac{(n-\Delta-1)^{k-1}}{k} \\ & \geq k \binom{n}{k} + [n(n-1) - 2m](k-1) \binom{n-1}{k-1} \frac{(n-\Delta-1)^{k-1}}{k}, \end{aligned}$$

where p is the number of pendant vertices in \overline{G} , and $q = \max\{k - p, 1\}$.

If $\delta = 1$ and $\Delta = n - 2$, then

$$\begin{aligned} & \text{SGut}_k(G) + \text{SGut}_k(\overline{G}) \\ & \geq k \binom{p}{k} + 2^q(k-1) \left[\binom{n}{k} - \binom{p}{k} \right] + k \binom{p'}{k} + 2^{q'}(k-1) \left[\binom{n}{k} - \binom{p'}{k} \right] \\ & \geq k \binom{n}{k} + k \binom{n}{k} \geq 2k \binom{n}{k}, \end{aligned}$$

where p, p' are the number of pendant vertices in G, \overline{G} , respectively, and $q = \max\{k - p, 1\}$, $q' = \max\{k - p', 1\}$.

From the above argument, we have

$$\text{SGut}_k(G) + \text{SGut}_k(\overline{G}) \geq \begin{cases} (n-1)(k-1) \binom{n}{k} t_1^{k-1} & \text{if } \delta \geq 2, \Delta \leq n-3 \\ 2m(k-1) \binom{n-1}{k-1} \frac{\delta^{k-1}}{k} + k \binom{n}{k} & \text{if } \delta \geq 2, \Delta = n-2 \\ k \binom{n}{k} + [n(n-1) - 2m](k-1) \binom{n-1}{k-1} \frac{(n-\Delta-1)^{k-1}}{k} & \text{if } \delta = 1, \Delta \leq n-3 \\ 2k \binom{n}{k} & \text{if } \delta = 1, \Delta = n-2. \end{cases}$$

For (3), from Proposition 1, if $\delta \geq 2$ and $\Delta \leq n - 3$, then

$$\text{SGut}_k(G) \cdot \text{SGut}_k(\overline{G}) \geq 2m(n^2 - n - 2m)(k - 1)^2 \binom{n-1}{k-1}^2 \frac{\delta^{k-1} (n - \Delta - 1)^{k-1}}{k^2}.$$

If $\delta \geq 2$ and $\Delta = n - 2$, then

$$\begin{aligned} & \text{SGut}_k(G) \cdot \text{SGut}_k(\overline{G}) \\ & \geq \left[2m(k - 1) \binom{n-1}{k-1} \frac{\delta^{k-1}}{k} \right] \left[k \binom{p'}{k} + 2^{q'}(k - 1) \left[\binom{n}{k} - \binom{p'}{k} \right] \right] \\ & \geq 2m(k - 1) \binom{n}{k} \binom{n-1}{k-1} \delta^{k-1}, \end{aligned}$$

where p' is the number of pendant vertices in \overline{G} , and $q' = \max\{k - p', 1\}$.

If $\delta = 1$ and $\Delta \leq n - 3$, then

$$\begin{aligned} & \text{SGut}_k(G) \cdot \text{SGut}_k(\overline{G}) \\ & \geq \left[[n(n - 1) - 2m](k - 1) \binom{n-1}{k-1} \frac{(n - \Delta - 1)^{k-1}}{k} \right] \left[k \binom{p}{k} + 2^q(k - 1) \left[\binom{n}{k} - \binom{p}{k} \right] \right] \\ & \geq [n(n - 1) - 2m](k - 1) \binom{n}{k} \binom{n-1}{k-1} (n - \Delta - 1)^{k-1}, \end{aligned}$$

where p is the number of pendant vertices in G , and $q = \max\{k - p, 1\}$.

If $\delta(G) = 1$ and $\Delta = n - 2$, then

$$\begin{aligned} & \text{SGut}_k(G) \cdot \text{SGut}_k(\overline{G}) \\ & \geq \left[k \binom{p}{k} + 2^q(k - 1) \left[\binom{n}{k} - \binom{p}{k} \right] \right] \left[k \binom{p'}{k} + 2^{q'}(k - 1) \left[\binom{n}{k} - \binom{p'}{k} \right] \right] \\ & \geq k^2 \binom{n}{k}^2, \end{aligned}$$

where p, p' are the number of pendant vertices in G and \overline{G} , respectively, and $q = \max\{k - p, 1\}$, $q' = \max\{k - p', 1\}$.

From the above argument, we have

$$\text{SGut}_k(G) \cdot \text{SGut}_k(\overline{G}) \geq \begin{cases} 2m(n^2 - n - 2m)(k - 1)^2 \binom{n-1}{k-1}^2 \frac{\delta^{k-1} (n - \Delta - 1)^{k-1}}{k^2} & \text{if } \delta(G) \geq 2, \Delta \leq n - 3 \\ 2m(k - 1) \binom{n}{k} \binom{n-1}{k-1} \delta^{k-1} & \text{if } \delta(G) \geq 2, \Delta = n - 2 \\ [n(n - 1) - 2m](k - 1) \binom{n}{k} \binom{n-1}{k-1} (n - \Delta - 1)^{k-1} & \text{if } \delta(G) = 1, \Delta \leq n - 3 \\ k^2 \binom{n}{k}^2 & \text{if } \delta(G) = 1, \Delta = n - 2. \end{cases}$$

To show the sharpness of the upper bound and the lower bound for $\delta(G) \geq 2, \Delta \leq n - 3$, we let G and \overline{G} be two $\frac{n-1}{2}$ -regular graphs of order n , where n is odd. If $k = n$, then $\text{SGut}_k(G) = (n - 1) \binom{\frac{n-1}{2}}{n}^n$, $\text{SGut}_k(\overline{G}) = (n - 1) \binom{\frac{n-1}{2}}{n}^n$, $s_1 = \max\{\Delta, n - \delta - 1\} = \frac{n-1}{2}$, $\Delta(n - \delta - 1) = \left(\frac{n-1}{2}\right)^2$, $t_1 = \min\{\delta, n - \Delta - 1\} = \frac{n-1}{2}$ and $\delta(n - \Delta - 1) = \left(\frac{n-1}{2}\right)^2$. Furthermore, we have $\text{SGut}_k(G) + \text{SGut}_k(\overline{G}) = 2(n - 1) \binom{\frac{n-1}{2}}{n}^n = (n - 1)^2 \binom{n}{k} s_1^{k-1}$, $\text{SGut}_k(G) \cdot \text{SGut}_k(\overline{G}) = (n - 1)^2 \binom{\frac{n-1}{2}}{n}^{2n} =$

$$2m(n^2 - n - 2m)(n - 1)^2 \binom{n-1}{k-1}^2 \frac{\Delta^{k-1} (n-\delta-1)^{k-1}}{k^2}, \text{ SGut}_k(G) + \text{SGut}_k(\overline{G}) = 2(n - 1) \left(\frac{n-1}{2}\right)^n = (n - 1)(k - 1) \binom{n}{k} t_1^{k-1} \text{ and } \text{SGut}_k(G) \cdot \text{SGut}_k(\overline{G}) = (n - 1)^2 \left(\frac{n-1}{2}\right)^{2n} = 2m(n^2 - n - 2m)(k - 1)^2 \binom{n-1}{k-1}^2 \frac{\delta^{k-1} (n-\Delta-1)^{k-1}}{k^2}. \quad \square$$

The following corollary is immediate from the above theorem.

Corollary 1. Let G be a connected graph of order $n \geq 4$ with maximum degree Δ and minimum degree δ . Then (1)

$$(n - 1)^2 \binom{n}{k} s_1^{k-1} \geq \text{SGut}_k(G) + \text{SGut}_k(\overline{G}) \geq \begin{cases} (n - 1)(k - 1) \binom{n}{k} t_1^{k-1} & \text{if } \delta \geq 2, \Delta \leq n - 3 \\ n(k - 1) \binom{n-1}{k-1} \frac{\delta^k}{k} + k \binom{n}{k} & \text{if } \delta \geq 2, \Delta = n - 2 \\ k \binom{n}{k} + n(k - 1) \binom{n-1}{k-1} \frac{(n-\Delta-1)^k}{k} & \text{if } \delta = 1, \Delta \leq n - 3 \\ 2k \binom{n}{k} & \text{if } \delta = 1, \Delta = n - 2, \end{cases}$$

where $s_1 = \min\{\Delta, n - \delta - 1\}$, $t_1 = \min\{\delta, n - \Delta - 1\}$; (2)

$$n^2 \binom{n-1}{k-1}^2 \frac{\Delta^{k-1} (n - \delta - 1)^{k-1} (n - 1)^4}{4k^2} \geq \text{SGut}_k(G) \cdot \text{SGut}_k(\overline{G}) \geq \begin{cases} n^2(k - 1)^2 \binom{n-1}{k-1}^2 \frac{\delta^k (n-\Delta-1)^k}{k^2} & \text{if } \delta \geq 2, \Delta \leq n - 3 \\ n(k - 1) \binom{n}{k} \binom{n-1}{k-1} \delta^k & \text{if } \delta \geq 2, \Delta = n - 2 \\ n(k - 1) \binom{n}{k} \binom{n-1}{k-1} (n - \Delta - 1)^k & \text{if } \delta = 1, \Delta \leq n - 3 \\ k^2 \binom{n}{k}^2 & \text{if } \delta = 1, \Delta = n - 2. \end{cases}$$

The following is the famous inequality by Pólya and Szegő:

Lemma 3. (Pólya–Szegő inequality) [25] Let (a_1, a_2, \dots, a_r) and (b_1, b_2, \dots, b_r) be two positive r -tuples such that there exist positive numbers M_1, m_1, M_2, m_2 satisfying:

$$0 < m_1 \leq a_i \leq M_1, 0 < m_2 \leq b_i \leq M_2, 1 \leq i \leq r.$$

Then

$$\frac{\sum_{i=1}^r a_i^2 \sum_{i=1}^r b_i^2}{\left(\sum_{i=1}^r a_i b_i\right)^2} \leq \frac{1}{4} \left(\sqrt{\frac{M_1 M_2}{m_1 m_2}} + \sqrt{\frac{m_1 m_2}{M_1 M_2}} \right)^2. \tag{10}$$

We now give more lower and upper bounds for $\text{SGut}_k(G) \cdot \text{SGut}_k(\overline{G})$ in terms of n, Δ and δ .

Theorem 2. Let G be a connected graph of order n with maximum degree Δ , minimum degree δ and a connected \overline{G} . Additionally, let k be an integer with $2 \leq k \leq n$. Then

$$\text{SGut}_k(G) \cdot \text{SGut}_k(\overline{G}) \geq \begin{cases} (k-1)^2 \delta^k (n-\delta-1)^k \binom{n}{k}^2 & \text{if } \Delta + \delta \leq n-1, \\ (k-1)^2 \Delta^k (n-\Delta-1)^k \binom{n}{k}^2 & \text{if } \Delta + \delta \geq n-1 \end{cases} \tag{11}$$

with equality holding if and only if G is a regular graph with $d_G(S) = d_{\overline{G}}(S) = k-1$ for any $S \subseteq V(G)$, $|S| = k$, and

$$\text{SGut}_k(G) \cdot \text{SGut}_k(\overline{G}) \leq \frac{(n-1)^{2k+2}}{2^{2k+2}} \binom{n}{k}^2 \left[\left(\frac{\Delta(n-\delta-1)}{\delta(n-\Delta-1)} \right)^k + \left(\frac{\delta(n-\Delta-1)}{\Delta(n-\delta-1)} \right)^k + 2 \right],$$

Moreover, the equality holds if and only if G is a $\left(\frac{n-1}{2}\right)$ -regular graph with $k = n$, n is odd.

Proof. Lower bound: By Cauchy–Schwarz inequality with (1), we have

$$\text{SGut}_k(G) \cdot \text{SGut}_k(\overline{G}) \geq (k-1)^2 \sum_{\substack{S \subseteq V(G) \\ |S|=k}} \left(\prod_{v \in S} \text{deg}_G(v) \right) \sum_{\substack{S \subseteq V(\overline{G}) \\ |S|=k}} \left(\prod_{v \in S} \text{deg}_{\overline{G}}(v) \right) \tag{12}$$

$$\geq (k-1)^2 \left(\sum_{\substack{S \subseteq V(G) \\ |S|=k}} \left(\prod_{v \in S} \text{deg}_G(v) \prod_{v \in S} \text{deg}_{\overline{G}}(v) \right)^{1/2} \right)^2 \tag{13}$$

$$\geq (k-1)^2 \left(\sum_{\substack{S \subseteq V(G) \\ |S|=k}} \left(\prod_{v \in S} \text{deg}_G(v) (n-1-\text{deg}_G(v)) \right)^{1/2} \right)^2.$$

Since $\delta \leq \text{deg}_G(v) \leq \Delta$, one can easily see that

$$\text{deg}_G(v) (n-1-\text{deg}_G(v)) \geq \begin{cases} \delta(n-\delta-1) & \text{if } \Delta + \delta \leq n-1, \\ \Delta(n-\Delta-1) & \text{if } \Delta + \delta \geq n-1. \end{cases} \tag{14}$$

From the above results, we have

$$\text{SGut}_k(G) \cdot \text{SGut}_k(\overline{G}) \geq \begin{cases} (k-1)^2 \delta^k (n-\delta-1)^k \binom{n}{k}^2 & \text{if } \Delta + \delta \leq n-1, \\ (k-1)^2 \Delta^k (n-\Delta-1)^k \binom{n}{k}^2 & \text{if } \Delta + \delta \geq n-1. \end{cases}$$

The equality holds in (12) if and only if $d_G(S) = d_{\overline{G}}(S) = k-1$ for any $S \subseteq V(G)$ with $|S| = k$. By the Cauchy–Schwarz inequality, the equality holds in (13) if and only if

$$\frac{\prod_{v \in S_1} \text{deg}_G(v)}{\prod_{v \in S_1} \text{deg}_{\overline{G}}(v)} = \frac{\prod_{v \in S_2} \text{deg}_G(v)}{\prod_{v \in S_2} \text{deg}_{\overline{G}}(v)} \text{ for any } S_1, S_2 \subseteq V(G) \text{ with } |S_1| = |S_2| = k,$$

that is, if and only if $\text{deg}_G(u) = \text{deg}_G(v)$ for any $u, v \in V(G)$, that is, if and only if G is a regular graph. Hence the equality holds in (11) if and only if G is a regular graph with $d_G(S) = d_{\overline{G}}(S) = k-1$ for any $S \subseteq V(G)$, $|S| = k$.

Upper bound: Let $\overline{\Delta}$ and $\overline{\delta}$ be the maximum degree and the minimum degree of graph \overline{G} , respectively. Then $\overline{\Delta} = n-\delta-1$ and $\overline{\delta} = n-\Delta-1$. By (1) and (10), we have

$$\begin{aligned}
 & \text{SGut}_k(G) \cdot \text{SGut}_k(\overline{G}) \\
 & \leq (n-1)^2 \sum_{\substack{S \subseteq V(G) \\ |S|=k}} \left(\prod_{v \in S} \text{deg}_G(v) \right) \sum_{\substack{S \subseteq V(\overline{G}) \\ |S|=k}} \left(\prod_{v \in S} \text{deg}_{\overline{G}}(v) \right) \\
 & \leq (n-1)^2 \left(\sum_{\substack{S \subseteq V(G) \\ |S|=k}} \left(\prod_{v \in S} \text{deg}_G(v) \prod_{v \in S} \text{deg}_{\overline{G}}(v) \right)^{1/2} \right)^2 \frac{1}{4} \left(\left(\frac{\Delta \overline{\Delta}}{\delta \overline{\delta}} \right)^{k/2} + \left(\frac{\delta \overline{\delta}}{\Delta \overline{\Delta}} \right)^{k/2} \right)^2 \\
 & \leq \frac{(n-1)^2}{4} \left(\sum_{\substack{S \subseteq V(G) \\ |S|=k}} \left(\prod_{v \in S} \text{deg}_G(v) (n-1 - \text{deg}_G(v)) \right)^{1/2} \right)^2 \left(\left(\frac{\Delta \overline{\Delta}}{\delta \overline{\delta}} \right)^{k/2} + \left(\frac{\delta \overline{\delta}}{\Delta \overline{\Delta}} \right)^{k/2} \right)^2.
 \end{aligned}$$

One can easily see that

$$\text{deg}_G(v) (n-1 - \text{deg}_G(v)) \leq \frac{(n-1)^2}{4} \text{ for any } v \in V(G).$$

Using this result in the above with $\overline{\Delta} = n - \delta - 1$ and $\overline{\delta} = n - \Delta - 1$, we get

$$\text{SGut}_k(G) \cdot \text{SGut}_k(\overline{G}) \leq \frac{(n-1)^{2k+2}}{2^{2k+2}} \binom{n}{k}^2 \left[\left(\frac{\Delta(n-\delta-1)}{\delta(n-\Delta-1)} \right)^k + \left(\frac{\delta(n-\Delta-1)}{\Delta(n-\delta-1)} \right)^k + 2 \right].$$

Moreover, the above equality holds if and only if G is a $\left(\frac{n-1}{2}\right)$ -regular graph with $k = n$, n is odd (very similar proof of the Proposition 1). □

Example 2. Let $G \cong C_n$ with $k = n$. Then $\delta = 2$ and hence

$$\text{SGut}_k(G) \cdot \text{SGut}_k(\overline{G}) = (n-1)^2(n-3)^n 2^n = (k-1)^2 \delta^k (n-\delta-1)^k \binom{n}{k}^2.$$

Let G be a $\left(\frac{n-1}{2}\right)$ -regular graph of order n with $k = n$ and odd n . Then $\Delta = \delta = \frac{n-1}{2}$ and hence

$$\begin{aligned}
 \text{SGut}_k(G) \cdot \text{SGut}_k(\overline{G}) &= \frac{(n-1)^{2n+2}}{2^{2n}} \\
 &= \frac{(n-1)^{2k+2}}{2^{2k+2}} \binom{n}{k}^2 \left[\left(\frac{\Delta(n-\delta-1)}{\delta(n-\Delta-1)} \right)^k + \left(\frac{\delta(n-\Delta-1)}{\Delta(n-\delta-1)} \right)^k + 2 \right].
 \end{aligned}$$

We now give more lower and upper bounds of $\text{SGut}_k(G) + \text{SGut}_k(\overline{G})$ in terms of n, Δ and δ .

Theorem 3. Let G be a connected graph of order n with maximum degree Δ , minimum degree δ and a connected \overline{G} . Additionally, let k be an integer with $2 \leq k \leq n$. Then

$$\text{SGut}_k(G) + \text{SGut}_k(\overline{G}) \geq \begin{cases} 2(k-1) \delta^{k/2} (n-\delta-1)^{k/2} \binom{n}{k} & \text{if } \Delta + \delta \leq n-1, \\ 2(k-1) \Delta^{k/2} (n-\Delta-1)^{k/2} \binom{n}{k} & \text{if } \Delta + \delta \geq n-1 \end{cases} \tag{15}$$

with equality holding if and only if G is a $\left(\frac{n-1}{2}\right)$ -regular graph with odd n and $d_G(S) = d_{\overline{G}}(S) = k-1$ for any $S \subseteq V(G), |S| = k$, and

$$\text{SGut}_k(G) + \text{SGut}_k(\overline{G}) \leq (n-1) \left[\Delta^k + (n-\delta-1)^k \right] \binom{n}{k} \quad (16)$$

with equality holding if and only if G is a regular graph with $k = n$.

Proof. For any two real numbers a, b , we have $(a-b)^2 \geq 0$, that is, $a^2 + b^2 \geq 2ab$ with equality holding if and only if $a = b$. Therefore we have

$$\begin{aligned} \prod_{v \in S} \text{deg}_G(v) + \prod_{v \in S} \text{deg}_{\overline{G}}(v) &\geq 2 \left(\prod_{v \in S} \text{deg}_G(v) \prod_{v \in S} \text{deg}_{\overline{G}}(v) \right)^{1/2} \\ &= 2 \left(\prod_{v \in S} \text{deg}_G(v) \text{deg}_{\overline{G}}(v) \right)^{1/2} \\ &= 2 \left(\prod_{v \in S} \text{deg}_G(v) (n - \text{deg}_G(v) - 1) \right)^{1/2}. \end{aligned}$$

From the above result with (14), we get

$$\prod_{v \in S} \text{deg}_G(v) + \prod_{v \in S} \text{deg}_{\overline{G}}(v) \geq \begin{cases} 2\delta^{k/2} (n-\delta-1)^{k/2} & \text{if } \Delta + \delta \leq n-1, \\ 2\Delta^{k/2} (n-\Delta-1)^{k/2} & \text{if } \Delta + \delta \geq n-1. \end{cases}$$

Now,

$$\begin{aligned} \text{SGut}_k(G) + \text{SGut}_k(\overline{G}) &= \sum_{\substack{S \subseteq V(G) \\ |S|=k}} \left[\left(\prod_{v \in S} \text{deg}_G(v) \right) d_G(S) + \left(\prod_{v \in S} \text{deg}_{\overline{G}}(v) \right) d_{\overline{G}}(S) \right] \\ &\geq (k-1) \sum_{\substack{S \subseteq V(G) \\ |S|=k}} \left[\prod_{v \in S} \text{deg}_G(v) + \prod_{v \in S} \text{deg}_{\overline{G}}(v) \right] \\ &\geq \begin{cases} 2(k-1) \delta^{k/2} (n-\delta-1)^{k/2} \binom{n}{k} & \text{if } \Delta + \delta \leq n-1, \\ 2(k-1) \Delta^{k/2} (n-\Delta-1)^{k/2} \binom{n}{k} & \text{if } \Delta + \delta \geq n-1. \end{cases} \end{aligned}$$

From the above, one can easily see that the equality holds in (15) if and only if G is a $\left(\frac{n-1}{2}\right)$ -regular graph with odd n and $d_G(S) = d_{\overline{G}}(S) = k-1$ for any $S \subseteq V(G)$, $|S| = k$.

Upper bound: By arithmetic-geometric mean inequality, we have

$$\begin{aligned}
\text{SGut}_k(G) + \text{SGut}_k(\bar{G}) &= \sum_{\substack{S \subseteq V(G) \\ |S|=k}} \left[\left(\prod_{v \in S} \text{deg}_G(v) \right) d_G(S) + \left(\prod_{v \in S} \text{deg}_{\bar{G}}(v) \right) d_{\bar{G}}(S) \right] \\
&\leq (n-1) \sum_{\substack{S \subseteq V(G) \\ |S|=k}} \left[\prod_{v \in S} \text{deg}_G(v) + \prod_{v \in S} \text{deg}_{\bar{G}}(v) \right] \\
&\leq (n-1) \sum_{\substack{S \subseteq V(G) \\ |S|=k}} \left[\left(\frac{\sum_{v \in S} \text{deg}_G(v)}{k} \right)^k + \left(\frac{\sum_{v \in S} \text{deg}_{\bar{G}}(v)}{k} \right)^k \right] \\
&= \frac{(n-1)}{k^k} \sum_{\substack{S \subseteq V(G) \\ |S|=k}} \left[\left(\sum_{v \in S} \text{deg}_G(v) \right)^k + \left(\sum_{v \in S} (n - \text{deg}_G(v) - 1) \right)^k \right] \\
&= \frac{(n-1)}{k^k} \sum_{\substack{S \subseteq V(G) \\ |S|=k}} \left[\left(\sum_{v \in S} \text{deg}_G(v) \right)^k + \left(k(n-1) - \sum_{v \in S} \text{deg}_G(v) \right)^k \right] \\
&\leq \frac{(n-1)}{k^k} \sum_{\substack{S \subseteq V(G) \\ |S|=k}} \left[(k\Delta)^k + (k(n-1) - k\delta)^k \right] \\
&= (n-1) \left[\Delta^k + (n - \delta - 1)^k \right] \binom{n}{k}.
\end{aligned}$$

From the above, one can easily see that the equality holds in (16) if and only if G is a regular graph with $k = n$ (very similar proof of the Proposition 1). \square

Example 3. Let G be a $\left(\frac{n-1}{2}\right)$ -regular graph with odd n and $k = n$. Then $\delta = \frac{n-1}{2}$ and hence

$$\text{SGut}_k(G) + \text{SGut}_k(\bar{G}) = \frac{(n-1)^{n+1}}{2^{n-1}} = 2(k-1)\delta^{k/2}(n-\delta-1)^{k/2} \binom{n}{k}$$

Let $G \cong C_n$ with $k = n$. Then $\Delta = \delta = 2, \bar{\Delta} = \bar{\delta} = 2$ and hence

$$\text{SGut}_k(G) + \text{SGut}_k(\bar{G}) = (n-1) \left[2^n + (n-3)^n \right] = (n-1) \left[\Delta^k + (n-\delta-1)^k \right] \binom{n}{k}.$$

Author Contributions: Conceptualization, Z.W., Y.M., K.C.D. and Y.S.; writing—original draft preparation, Z.W.; methodology, Y.M., K.C.D. and Y.S.; writing—review and editing, Y.M., K.C.D. and Y.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the National Science Foundation of China grant numbers 12061059, 11601254, 11661068, 11551001, 11161037 and 11461054, and the UoA Flexible Fund from Northumbria University grant number 201920A1001.

Acknowledgments: The authors are very grateful to three anonymous referees for their valuable comments on the paper, which have considerably improved the presentation of this paper.

Conflicts of Interest: The author declares no conflict of interest.

References

1. Bondy, J.A.; Murty, U.S.R. *Graph Theory*; Springer: Berlin/Heidelberg, Germany, 2008.
2. Buckley, F.; Harary, F. *Distance in Graphs*; Addison-Wesley: Redwood, CA, USA, 1990.
3. Goddard, W.; Oellermann, O.R. Distance in graphs. In *Structural Analysis of Complex Networks*; Dehmer, M., Ed.; Birkhäuser: Dordrecht, The Netherlands, 2011; pp. 49–72.
4. Ali, P.; Dankelmann, P.; Mukwembi, S. Upper bounds on the Steiner diameter of a graph. *Discrete Appl. Math.* **2012**, *160*, 1845–1850. [[CrossRef](#)]
5. Cáceresa, J.; Mxaxrquezb, A.; Puertasa, M.L. Steiner distance and convexity in graphs. *Eur. J. Comb.* **2008**, *29*, 726–736. [[CrossRef](#)]
6. Chartrand, G.; Oellermann, O.R.; Tian, S.; Zou, H.B. Steiner distance in graphs. *Čas. Pest. Mat.* **1989**, *114*, 399–410. [[CrossRef](#)]
7. Dankelmann, P.; Oellermann, O.R.; Swart, H.C. The average Steiner distance of a graph. *J. Graph Theory* **1996**, *22*, 15–22. [[CrossRef](#)]
8. Oellermann, O.R.; Tian, S. Steiner centers in graphs. *J. Graph Theory* **1990**, *14*, 585–597. [[CrossRef](#)]
9. Li, X.; Mao, Y.; Gutman, I. The Steiner Wiener index of a graph. *Discuss. Math. Graph Theory* **2016**, *36*, 455–465.
10. Gutman, I. On Steiner degree distance of trees. *Appl. Math. Comput.* **2016**, *283*, 163–167. [[CrossRef](#)]
11. Chen, S.B.; Liu, W.J. Extremal modified Schultz index of bicyclic graphs. *MATCH Commun. Math. Comput. Chem.* **2010**, *64*, 767–782.
12. Dankelmann, P.; Gutman, I.; Mukwembi, S.; Swart, H.C. The edge-Wiener index of a graph. *Discret. Appl. Math.* **2009**, *309*, 3452–3457. [[CrossRef](#)]
13. Das, K.C.; Su, G.; Xiong, L. Relation between Degree Distance and Gutman Index of Graphs. *MATCH Commun. Math. Comput. Chem.* **2016**, *76*, 221–232.
14. Mao, Y.; Das, K.C. Steiner Gutman index. *MATCH Commun. Math. Comput. Chem.* **2018**, *79*, 779–794.
15. Shang, Y. Non-hyperbolicity of random graphs with given expected degrees. *Stoch. Models* **2013**, *29*, 451–462.
16. Shang, Y. A note on the warmth of random graphs with given expected degrees. *Int. J. Math. Math. Sci.* **2014**, *2014*, 749856. [[CrossRef](#)]
17. Mao, Y.; Wang, Z.; Das, K.C. Steiner degree distance of two graph products. *Analele Stiintifice Univ. Ovidius Constanta* **2019**, *27*, 83–99. [[CrossRef](#)]
18. Mao, Y.; Wang, Z.; Gutman, I.; Klobučar, A. Steiner degree distance. *MATCH Commun. Math. Comput. Chem.* **2017**, *78*, 221–230.
19. Mao, Y.; Wang, Z.; Gutman, I.; Li, H. Nordhaus-Gaddum-type results for the Steiner Wiener index of graphs. *Discret. Appl. Math.* **2017**, *219*, 167–175. [[CrossRef](#)]
20. Aouchiche, M.; Hansen, P. A survey of Nordhaus-Gaddum type relations. *Discret. Appl. Math.* **2013**, *161*, 466–546. [[CrossRef](#)]
21. Hua, H.; Das, K.C. On the Wiener polarity index of graphs. *Appl. Math. Comput.* **2016**, *280*, 162–167. [[CrossRef](#)]
22. Mao, Y. Nordhaus-Gaddum Type Results in Chemical Graph Theory. In *Bounds in Chemical Graph Theory—Advances*; Gutman, I., Furtula, B., Das, K.C., Milovanović, E., Milovanović, I., Eds.; University of Kragujevac and Faculty of Science Kragujevac: Kragujevac, Serbia, 2017; pp. 3–127.
23. Zhang, Y.; Hu, Y. The Nordhaus-Gaddum-type inequality for the Wiener polarity index. *Appl. Math. Comput.* **2016**, *273*, 880–884. [[CrossRef](#)]
24. Shang, Y. Bounds of distance Estrada index of graphs. *Ars Comb.* **2016**, *128*, 287–294.
25. Pólya, G.; Szegő, G. *Problems and Theorems in Analysis I, Series, Integral Calculus, Theory of Functions*; Springer: Berlin/Heidelberg, Germany, 1972.

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).