

Northumbria Research Link

Citation: Jeong, Seong-Yeop, Verscharen, Daniel, Wicks, Robert and Fazakerley, Andrew N. (2020) A Quasi-linear Diffusion Model for Resonant Wave-Particle Instability in Homogeneous Plasma. *The Astrophysical Journal*, 902 (2). p. 128. ISSN 1538-4357

Published by: The American Astronomical Society

URL: <https://doi.org/10.3847/1538-4357/abb099> <<https://doi.org/10.3847/1538-4357/abb099>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/44714/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



**Northumbria
University**
NEWCASTLE



UniversityLibrary

A Quasi-Linear Diffusion Model for Resonant Wave-Particle Instability in Homogeneous Plasma

SEONG-YEOP JEONG,¹ DANIEL VERSCHAREN,^{1,2} ROBERT T. WICKS,^{1,3} AND ANDREW N. FAZAKERLEY¹

¹*Mullard Space Science Laboratory, University College London, Dorking, RH5 6NT, UK; s.jeong.17@ucl.ac.uk, d.verscharen@ucl.ac.uk*

²*Space Science Center, University of New Hampshire, Durham, NH 03824, USA*

³*Institute for Risk and Disaster Reduction, University College London, Gower Street, London, WC1E 6BT, UK*

Abstract

In this paper, we develop a model to describe the generalized wave-particle instability in a quasi-neutral plasma. We analyze the quasi-linear diffusion equation for particles by expressing an arbitrary unstable and resonant wave mode as a Gaussian wave packet, allowing for an arbitrary direction of propagation with respect to the background magnetic field. We show that the localized energy density of the Gaussian wave packet determines the velocity-space range in which the dominant wave-particle instability and counter-acting damping contributions are effective. Moreover, we derive a relation describing the diffusive trajectories of resonant particles in velocity space under the action of such an interplay between the wave-particle instability and damping. For the numerical computation of our theoretical model, we develop a mathematical approach based on the Crank-Nicolson scheme to solve the full quasi-linear diffusion equation. Our numerical analysis solves the time evolution of the velocity distribution function under the action of a dominant wave-particle instability and counter-acting damping, and shows a good agreement with our theoretical description. As an application, we use our model to study the oblique fast-magnetosonic/whistler instability, which is proposed as a scattering mechanism for strahl electrons in the solar wind. In addition, we numerically solve the full Fokker-Planck equation to compute the time evolution of the electron-strahl distribution function under the action of Coulomb collisions with core electrons and protons after the collisionless action of the oblique fast-magnetosonic/whistler instability.

Keywords: methods: analytical — instabilities — waves — plasmas — diffusion

1. INTRODUCTION

Wave-particle resonances play an important role for the energy exchange between particles and waves in many space and astrophysical plasmas. For example, wave-particle resonances contribute to the acceleration and deceleration of particles in radiation belts (Ukhorskiy & Sitnov 2014), the deviation of the particle velocity distribution function (VDF) from a Maxwellian equilibrium in the solar wind (Marsch 2006), the thermodynamic state of the intracluster medium in galaxy clusters (Roberg-Clark et al. 2016), and the scattering and absorption of the surface radiation in neutron-star magnetospheres (Lyutikov & Gavriil 2006). Therefore, it is of great importance to study the mechanics of wave-particle resonances in order to advance our understanding of the physics of astrophysical plasmas throughout the Universe.

According to kinetic theory, wave-particle resonances can occur in the form of Landau or cyclotron resonances, which contribute to wave instability or wave damping depending on the resonance’s characteristics. The quasi-

linear theory of magnetized plasma, first established by Yakimenko (1963) and Kennel & Engelmann (1966), provides a mathematical framework to predict the evolution of the particle VDF under the action of the wave-particle resonances. The quasi-linear theory assumes that the spatially averaged VDF evolves slowly compared to the gyroperiod of the particles and the wave period. It furthermore assumes that the fluctuation amplitude is small and that the spatial average of the fluctuations vanishes. Based on this theory, numerous analytical studies have successfully explained the evolution of VDFs resulting from wave-particle resonances.

Resonant particles diffuse along specific trajectories in velocity space determined by the properties of the resonant wave (Kennel & Engelmann 1966; Gendrin 1968; Gendrin & Roux 1980; Gendrin 1981; Stix 1992; Isenberg & Lee 1996; Summers et al. 1998, 2001). Quasi-linear diffusion coefficients determine the diffusion rate of the resonant particles (Lyons et al. 1971; Lyons 1974; Albert 2004; Summers 2005; Glauert & Horne 2005; Isenberg & Vasquez 2011; Tang et al. 2020). Alternatively, quasi-

linear diffusion models based on a bi-Maxwellian VDF, in which only its moments evolve in time, describe the effective evolution of particle VDFs under the action of micro-instabilities (Yoon & Seough 2012; Seough & Yoon 2012; Yoon et al. 2015; Yoon 2017; Yoon et al. 2017). Moreover, fully non-linear simulations based on kinetic theory model the evolution of the particle VDF consistently with predictions from quasi-linear theory (Vocks & Mann 2003; Vocks et al. 2005; Gary et al. 2008; Saito et al. 2008, 2010; Saito & Peter Gary 2012).

Realistic analytical models must describe the diffusive trajectory of the resonant particles in velocity space, taking into account the localized (in wavevector space) energy density of the resonant waves. These models must also account for non-Maxwellian features in the VDF evolution in order to advance our understanding of plasma observations and kinetic simulation results. A rigorous numerical analysis of the diffusion equation, including both the diagonal and off-diagonal diffusion terms, is necessary to support the theoretical description through the quantification of the diffusion rates.

By analyzing the quasi-linear diffusion equation, we propose a novel quasi-linear diffusion model for the time evolution of VDFs under the action of a dominant wave-particle instability and counter-acting damping contributions in Section 2. Our model describes the creation and evolution of non-Maxwellian features in the particle VDF. We allow for an arbitrary type of the unstable and resonant wave mode with an arbitrary direction of propagation with respect to the background magnetic field. In our analysis, we express the electric field of this wave as a Gaussian wave packet in configuration space. The localization of such a wave packet in configuration space is the direct consequence of its generation through a linear instability, which is localized in wavevector space.

To investigate the stabilization of the VDF through quasi-linear diffusion, we apply our analysis of the quasi-linear diffusion equation to Boltzmann’s H -theorem. In this scheme, the localized energy density of the Gaussian wave packet in wavevector space defines the velocity-space range in which the dominant wave-particle instability and counter-acting damping contributions are effective. In addition, we derive a relation to describe the diffusive trajectories of resonant particles in velocity space under the action of such an instability and damping. In this way, our model accounts for the diffusive behavior of resonant particles in different regions of velocity space.

For the numerical evaluation of our theoretical description, we develop a mathematical approach based on the Crank-Nicolson scheme (for numerical details, see Appendix A) that solves the full quasi-linear diffusion equation. Because of its reliable stability, the

Crank-Nicolson scheme has been used previously to solve diffusion equations in a variety of fields (Khazanov et al. 2002; Albert 2004; Brüggemann et al. 2004; Yang et al. 2009; Klein & Chandran 2016; Taran et al. 2019). However, most standard Crank-Nicolson schemes neglect the off-diagonal terms in the diffusion equation. In our case, these off-diagonal terms are important for the description of resonant pitch-angle scattering. We note that our mathematical approach is applicable to all general two-dimensional diffusion equations, including those with off-diagonal diffusion terms.

In Section 3, as an example, we apply our model to the scattering of the electron strahl which is a field-aligned electron beam population in the solar wind (Pilipp et al. 1987; Štverák et al. 2009). Observations in the solar wind suggest that strahl electrons exchange energy with whistler waves, which ultimately leads to a scattering of strahl electrons into the halo population (Pagel et al. 2007; Lacombe et al. 2014; Graham et al. 2017). Our quasi-linear framework confirms that an instability of the fast-magnetosonic/whistler wave in oblique propagation with respect to the background magnetic field scatters the electron strahl into the electron halo, as predicted by linear theory (Vasko et al. 2019; Verscharen et al. 2019).

In Section 4, for a more realistic model of the strahl evolution after the collisionless action of the oblique fast-magnetosonic/whistler instability, we numerically solve the full Fokker-Planck equation for Coulomb collisions with our mathematical approach (for numerical details, see Appendix B). We model the time evolution of the electron-strahl VDF through the action of Coulomb collisions with core electrons and protons. This combined method allows us to compare the timescales for the strahl scattering and collisional relaxation.

In Section 5, we discuss the results of our model for the strahl scattering and electron-halo formation through the instability and Coulomb collisions. In Section 6, we summarize and conclude our treatment.

2. QUASI-LINEAR DIFFUSION MODEL

In this section, we establish our general theoretical framework for the description of a resonant wave-particle instability in quasi-linear theory. Since our work focuses on non-relativistic space plasma like the solar wind, we neglect relativistic effects throughout our study.

2.1. Analysis of Quasi-Linear Diffusion Equation

To investigate the time evolution of the particle VDF through wave-particle resonances, we study the quasi-

linear diffusion equation, given by Stix (1992)

$$\left(\frac{\partial f_j}{\partial t}\right)_{QLD} = \lim_{V \rightarrow \infty} \sum_{n=-\infty}^{\infty} \int \frac{\pi q_j^2}{V m_j^2} \times \hat{G}[k_{\parallel}] \left[\frac{v_{\perp}^2}{|v_{\parallel}|} \delta\left(k_{\parallel} - \frac{\omega_k - n\Omega_j}{v_{\parallel}}\right) |\psi_j^n|^2 \hat{G}[k_{\parallel}] f_j \right] d^3 \mathbf{k}, \quad (1)$$

where

$$\psi_j^n \equiv \frac{1}{\sqrt{2}} \left[E_k^R e^{i\phi} J_{n+1}(\rho_j) + E_k^L e^{-i\phi} J_{n-1}(\rho_j) \right] + \frac{v_{\parallel}}{v_{\perp}} E_k^z J_n(\rho_j), \quad (2)$$

and

$$\hat{G}[k_{\parallel}] \equiv \left(1 - \frac{k_{\parallel} v_{\parallel}}{\omega_k}\right) \frac{1}{v_{\perp}} \frac{\partial}{\partial v_{\perp}} + \frac{k_{\parallel}}{\omega_k} \frac{\partial}{\partial v_{\parallel}}. \quad (3)$$

The integer n determines the order of the resonance, where $n = 0$ corresponds to the Landau resonance and $n \neq 0$ corresponds to cyclotron resonances. In our equations, we label contributions from a given resonance order with a superscript n . The subscript j indicates the particle species. The particle VDF of species j is denoted as $f_j \equiv f_j(v_{\perp}, v_{\parallel}, t)$ which is spatially averaged and gyrotropic, q_j and m_j are the charge and mass of a particle of species j , v_{\perp} and v_{\parallel} are the velocity coordinates perpendicular and parallel with respect to the background magnetic field. We choose the coordinate system in which the proton bulk velocity is zero.

We denote the n^{th} -order Bessel function as $J_n(\rho_j)$ where $\rho_j \equiv k_{\perp} v_{\perp} / \Omega_j$. The cyclotron frequency of species j is defined as $\Omega_j \equiv q_j B_0 / m_j c$, \mathbf{B}_0 is the background magnetic field, c is the light speed, k_{\perp} and k_{\parallel} are the perpendicular and parallel components of the wavevector \mathbf{k} , and V is the volume in which the wave amplitude is effective so that the wave and particles undergo a significant interaction. We denote Dirac's δ -function as δ and the azimuthal angle of wavevector \mathbf{k} as ϕ . The frequency ω is a complex function of \mathbf{k} , and we define ω_k as its real part and γ_k as its imaginary part ($\omega = \omega_k + i\gamma_k$). Without loss of generality, we set $\omega_k > 0$. Furthermore, we assume that $|\gamma_k| \ll \omega_k$, i.e. the assumption of slow growth or damping that is central to quasi-linear theory.

The spatially Fourier-transformed electric field has the form of $\mathbf{E}_{\mathbf{k}} = \hat{x}E_k^x + \hat{y}E_k^y + \hat{z}E_k^z$ and is defined as (Gurnett & Bhattacharjee 2017)

$$\mathbf{E}_{\mathbf{k}} = \frac{1}{(2\pi)^{3/2}} \int \mathbf{E}_{\mathbf{r}} \exp[-i\mathbf{k} \cdot \mathbf{r}] d^3 \mathbf{r}, \quad (4)$$

where $\mathbf{k} \cdot \mathbf{r} = k_{\perp} x \cos \phi + k_{\perp} y \sin \phi + k_{\parallel} z$ and \mathbf{r} is the position vector. We take the constant background magnetic field as $\mathbf{B}_0 = \hat{z}B_0$ and define the right- and left-circularly polarized components of the electric field as

$E_k^R \equiv (E_k^x - iE_k^y)/\sqrt{2}$ and $E_k^L \equiv (E_k^x + iE_k^y)/\sqrt{2}$. We define the longitudinal component of the electric field as E_k^z .

Linear instabilities typically create fluctuations across a finite range of wavevectors. The Fourier transformation of such a wave packet in wavevector space corresponds to a wave packet in configuration space. For the sake of simplicity, we model this finite wave packet by assuming that the electric field $\mathbf{E}_{\mathbf{r}}$ of the unstable and resonant waves has the shape of a gyrotropic Gaussian wave packet

$$\mathbf{E}_{\mathbf{r}} = \mathbf{E}_0 \exp\left[-\frac{\sigma_{\perp 0}^2 x^2 + \sigma_{\perp 0}^2 y^2 + \sigma_{\parallel 0}^2 z^2}{2}\right] \exp[i\mathbf{k}_0 \cdot \mathbf{r}], \quad (5)$$

where $\mathbf{E}_0 = \hat{x}E_0^x + \hat{y}E_0^y + \hat{z}E_0^z$, $\mathbf{k}_0 \cdot \mathbf{r} = k_{\perp 0} x \cos \phi + k_{\perp 0} y \sin \phi + k_{\parallel 0} z$, and \mathbf{k}_0 is the wavevector of the Gaussian wave packet. We allow for an arbitrary angle θ_0 between \mathbf{k}_0 and \mathbf{B}_0 , which defines the orientation of the wavevector at maximum growth of the wave, and assume that $k_{\parallel 0} \neq 0$. The vector \mathbf{E}_0 represents the peak amplitude of the electric field. The free parameters $\sigma_{\perp 0}$ and $\sigma_{\parallel 0}$ characterize the width of the Gaussian envelope. Quasi-linear theory requires that $\mathbf{E}_{\mathbf{r}}$ spatially averages to zero. Therefore, we assume that $|k_{\parallel 0}| \gg \sigma_{\parallel 0}$ so that the spatial dimension of the Gaussian wave packet is large compared to the parallel wave length $2\pi/|k_{\parallel 0}|$.

The spatial Fourier transformation of Eq. (5) according to Eq. (4) then leads to

$$\mathbf{E}_{\mathbf{k}} = \frac{\mathbf{E}_0}{\sigma_{\parallel 0} \sigma_{\perp 0}^2} \exp\left[-\frac{(k_{\parallel} - k_{\parallel 0})^2}{2\sigma_{\parallel 0}^2} - \frac{(k_{\perp} - k_{\perp 0})^2}{2\sigma_{\perp 0}^2}\right]. \quad (6)$$

We identify V with the volume of the Gaussian envelope, $V = 1/(\sigma_{\parallel 0} \sigma_{\perp 0}^2)$. Eq. (6) represents the localization of the wave energy density in wavevector space. For the instability analysis through Eq. (6), we define the unstable \mathbf{k} -spectrum as the finite wavevector range in which $\gamma_k > 0$ and argue that resonant waves exist only in this unstable \mathbf{k} -spectrum. We neglect any waves outside this \mathbf{k} -spectrum since they are damped.

We define $k_{\parallel 0}$ as the value of k_{\parallel} at the center of the unstable \mathbf{k} -spectrum. We then obtain

$$k_{\perp 0} = k_{\parallel 0} \tan \theta_0. \quad (7)$$

In the case of a linear plasma instability, we identify $k_{\perp 0}$ and $k_{\parallel 0}$ with the wavevector components at which the instability has its maximum growth rate as a reasonable approximation. To approximate the wave frequency of the unstable waves at the angle θ_0 of maximum growth, we expand ω_k of the unstable and resonant waves around $k_{\parallel 0}$ as

$$\omega_k(k_{\parallel}) \approx \omega_{k0} + v_{g0} (k_{\parallel} - k_{\parallel 0}), \quad (8)$$

where

$$v_{g0} \equiv \left. \frac{\partial \omega_k}{\partial k_{\parallel}} \right|_{k_{\parallel}=k_{\parallel 0}}. \quad (9)$$

In Eqs. (8) and (9), ω_{k0} and v_{g0} are the wave frequency and parallel group velocity of the unstable and resonant waves, evaluated at $k_{\parallel} = k_{\parallel 0}$. We select the values of $\sigma_{\perp 0}$ and $\sigma_{\parallel 0}$ as the half widths of the perpendicular and parallel unstable \mathbf{k} -spectrum. In the case of a linear plasma instability, the numerical values for $k_{\perp 0}$, $k_{\parallel 0}$, $\sigma_{\perp 0}$, $\sigma_{\parallel 0}$, ω_{k0} and v_{g0} can be found from the solutions of the hot-plasma dispersion relation, which thus closes our set of equations.

By using Eq. (6) and Eq. (8), we rewrite Eq. (1) as

$$\left(\frac{\partial f_j}{\partial t} \right)_{QLD} = \sum_{n=-\infty}^{\infty} \int \hat{G}[k_{\parallel}] [D_j^n \hat{G}[k_{\parallel}] f_j] d^3 \mathbf{k}, \quad (10)$$

where

$$D_j^n \equiv \frac{\pi q_j^2 v_{\perp}^2}{\sigma_{\parallel 0} \sigma_{\perp 0}^2 m_j^2} \delta(k_{\parallel} - k_{\parallel j}^n) \times \frac{|\psi_{j0}^n|^2}{|v_{\parallel} - v_{g0}|} \exp \left[-\frac{(k_{\parallel} - k_{\parallel 0})^2}{\sigma_{\parallel 0}^2} - \frac{(k_{\perp} - k_{\perp 0})^2}{\sigma_{\perp 0}^2} \right], \quad (11)$$

$$\psi_{j0}^n \equiv \frac{1}{\sqrt{2}} \left[E_0^R e^{i\phi} J_{n+1}(\rho_j) + E_0^L e^{-i\phi} J_{n-1}(\rho_j) \right] + \frac{v_{\parallel}}{v_{\perp}} E_0^z J_n(\rho_j), \quad (12)$$

and

$$k_{\parallel j}^n \equiv \frac{\omega_{k0} - k_{\parallel 0} v_{g0} - n\Omega_j}{v_{\parallel} - v_{g0}}. \quad (13)$$

We note that $E_0^R = (E_0^x - iE_0^y)/\sqrt{2}$ and $E_0^L = (E_0^x + iE_0^y)/\sqrt{2}$ are constant, evaluated at \mathbf{k}_0 .

Eq. (10) is the quasi-linear diffusion equation describing the action of the dominant wave-particle instability and co-existing damping contributions from other resonances in a Gaussian wave packet. We define the n resonance as the contribution to the summation in Eq. (10) with only integer n . We note that any n resonance can contribute to wave instability or to wave damping depending on the resonance's characteristics.

2.2. Stabilization through a Resonant Wave-Particle Instability

We define the stabilization as the process that creates the condition in which $(\partial f_j / \partial t)_{QLD} \rightarrow 0$ for all v_{\perp} and v_{\parallel} . For our analysis of the stabilization of a VDF through a resonant wave-particle instability, including co-existing damping effects, we use Boltzmann's H -theorem, in which the quantity H is defined as

$$H(t) \equiv \int f_j(\mathbf{v}, t) \ln f_j(\mathbf{v}, t) d^3 \mathbf{v}. \quad (14)$$

By using Eq. (10), the time derivative of H is given by

$$\frac{dH}{dt} = \sum_{n=-\infty}^{\infty} \iint (\ln f_j + 1) \hat{G}[k_{\parallel}] [D_j^n \hat{G}[k_{\parallel}] f_j] d^3 \mathbf{k} d^3 \mathbf{v}. \quad (15)$$

The integrand in Eq. (15) is equivalent to

$$(\ln f_j + 1) \hat{G}[k_{\parallel}] [D_j^n \hat{G}[k_{\parallel}] f_j] = \hat{G}[k_{\parallel}] [D_j^n \hat{G}[k_{\parallel}] (f_j \ln f_j)] - D_j^n [\hat{G}[k_{\parallel}] f_j]^2 / f_j. \quad (16)$$

Upon substituting Eq. (16) into Eq. (15), the first term on the right-hand side in Eq. (16) disappears after the integration over \mathbf{v} . Then, by resolving the δ -function in D_j^n through the k_{\parallel} -integral, we obtain

$$\frac{dH}{dt} = - \sum_{n=-\infty}^{\infty} \left(\frac{dH}{dt} \right)^n, \quad (17)$$

where

$$\left(\frac{dH}{dt} \right)^n \equiv \int \left\{ \tilde{D}_j^n [\hat{G}[k_{\parallel j}^n] f_j]^2 / f_j \right\} d^3 \mathbf{v}, \quad (18)$$

$$\tilde{D}_j^n \equiv W_j^n \frac{\pi q_j^2 v_{\perp}^2}{\sigma_{\parallel 0} \sigma_{\perp 0}^2 m_j^2} \int_0^{2\pi} \int_0^{\infty} |\psi_{j0}^n|^2 \times \exp \left[-\frac{(k_{\perp} - k_{\perp 0})^2}{\sigma_{\perp 0}^2} \right] k_{\perp} dk_{\perp} d\phi, \quad (19)$$

$$W_j^n \equiv \frac{1}{|v_{\parallel} - v_{g0}|} \exp \left[-\frac{k_{\parallel 0}^2}{\sigma_{\parallel 0}^2} \left(\frac{v_{\parallel} - v_{\parallel res}^n}{v_{\parallel} - v_{g0}} \right)^2 \right], \quad (20)$$

$$v_{\parallel res}^n \equiv \frac{\omega_{k0} - n\Omega_j}{k_{\parallel 0}}, \quad (21)$$

$$\hat{G}[k_{\parallel j}^n] \equiv \left[\frac{n\Omega_j}{\omega_{k0} - k_{\parallel 0} v_{g0} - n\Omega_j} \right] \frac{v_{\parallel} - v_{g0}}{v_{ph} v_{\perp}} \frac{\partial}{\partial v_{\perp}} + \frac{1}{v_{ph}} \frac{\partial}{\partial v_{\parallel}}, \quad (22)$$

and

$$v_{ph} \equiv \frac{\omega_k(k_{\parallel j}^n)}{k_{\parallel j}^n} = \frac{(\omega_{k0} - k_{\parallel 0} v_{g0}) v_{\parallel} - n\Omega_j v_{g0}}{\omega_{k0} - k_{\parallel 0} v_{g0} - n\Omega_j}. \quad (23)$$

The function \tilde{D}_j^n in Eq. (19) plays the role of a diffusion coefficient for the n resonance. In \tilde{D}_j^n , the v_{\parallel} -function W_j^n defined in Eq. (20) serves as a window function that determines the region in v_{\parallel} -space in which the quasi-linear diffusion through the n resonance is effective. The window function W_j^n is maximum at $v_{\parallel res}^n$ defined in

Eq. (21), which is the parallel velocity of the particles that resonate with the waves at $k_{\parallel} = k_{\parallel 0}$ through the n resonance. Our window function W_j^n is linked to Dirac's δ -function in the limit

$$\lim_{v_{\parallel res}^n \rightarrow v_{g0}} W_j^n \approx \sqrt{\pi} \frac{\sigma_{\parallel 0}}{|k_{\parallel 0}|} \delta(v_{\parallel} - v_{g0}), \quad (24)$$

where $|k_{\parallel 0}| \gg \sigma_{\parallel 0}$. Due to this ordering between $|k_{\parallel 0}|$ and $\sigma_{\parallel 0}$, we assume that W_j^n restricts a finite region in v_{\parallel} -space and that the W_j^n for different resonances do not overlap with each other in v_{\parallel} -space.

Only particles distributed within W_j^n experience the n resonance and contribute to the quasi-linear diffusion which is ultimately responsible for the stabilization. Since all terms in Eq. (18) are positive semi-definite, all resonances independently stabilize f_j through quasi-linear diffusion in the v_{\parallel} -range defined by their respective W_j^n , according to Eq. (17). Therefore, H decreases and dH/dt tends towards zero during the quasi-linear diffusion through all resonances while f_j is in the process of stabilization. When f_j reaches a state of full stabilization through all n resonances, the instability has saturated and its growth ends.

The v_{\parallel} -function $k_{\parallel j}^n$ defined in Eq. (13) is the resonant parallel wavenumber, where $k_{\parallel j}^n = k_{\parallel 0}$ at $v_{\parallel} = v_{\parallel res}^n$. It quantifies the k_{\parallel} -component of the unstable \mathbf{k} -spectrum in the v_{\parallel} -range defined by W_j^n . Eq. (23) defines the phase velocity at $k_{\parallel j}^n$, which is only constant when $v_{g0} = \omega_{k0}/k_{\parallel 0}$, in which case $v_{ph} = v_{g0}$ for all v_{\parallel} . We discuss the diffusion operator $\hat{G}[k_{\parallel j}^n]$ in Eq. (22) in the next section.

2.3. Nature of Quasi-Linear Diffusion in Velocity Space

Considering Eq. (17), unless the wave amplitude is zero, the condition for the achievement of stabilization through the n resonance is

$$\hat{G}[k_{\parallel j}^n] \mathbb{F}_j^n(v_{\perp}, v_{\parallel}) = 0, \quad (25)$$

where $\mathbb{F}_j^n(v_{\perp}, v_{\parallel})$ represents the stabilized VDF of species j through the n resonance. According to Eq. (25), $\hat{G}[k_{\parallel j}^n]$ is a directional derivative along the isocontour of \mathbb{F}_j^n evaluated at a given velocity position. Considering the role of W_j^n , $\hat{G}[k_{\parallel j}^n]$ describes only the diffusion of resonant particles within W_j^n along the isocontour of \mathbb{F}_j^n . Consequently, the particles experiencing the n resonance diffuse toward the stable state so that $(dH/dt)^n \rightarrow 0$, while the isocontour of \mathbb{F}_j^n describes the diffusive trajectory for the n resonance.

To find such that trajectory, we express an infinitesimal variation of \mathbb{F}_j^n along its isocontour as

$$d\mathbb{F}_j^n = \frac{\partial \mathbb{F}_j^n}{\partial v_{\perp}} dv_{\perp} + \frac{\partial \mathbb{F}_j^n}{\partial v_{\parallel}} dv_{\parallel} = 0. \quad (26)$$

Eqs. (22) and (26) allow us to rewrite Eq. (25) as

$$v_{\perp} dv_{\perp} + \left[\frac{n\Omega_j}{n\Omega_j - \omega_{k0} + k_{\parallel 0} v_{g0}} \right] (v_{\parallel} - v_{g0}) dv_{\parallel} = 0. \quad (27)$$

By integrating Eq. (27), the diffusive trajectory for the n resonance is then defined by

$$v_{\perp}^2 + \left[\frac{n\Omega_j}{n\Omega_j - \omega_{k0} + k_{\parallel 0} v_{g0}} \right] (v_{\parallel} - v_{g0})^2 = const. \quad (28)$$

Kennel & Engelmann (1966) treat the two limiting cases in which $v_{g0} = \omega_{k0}/k_{\parallel 0}$ and $v_{g0} = 0$. Using their assumptions, our Eq. (28) is equivalent to their equation (4.8) if $v_{g0} = \omega_{k0}/k_{\parallel 0}$, and our Eq. (28) is equivalent to their equation (4.11) if $v_{g0} = 0$. Depending on the dispersion properties of the resonant waves, Eq. (28) is either an elliptic or a hyperbolic equation when $n \neq 0$. In the case of electron resonances, it is safe to assume that

$$\frac{n\Omega_j}{n\Omega_j - \omega_{k0} + k_{\parallel 0} v_{g0}} \geq 0, \quad (29)$$

in Eq. (28) if $v_{g0} < (\omega_{k0} + n|\Omega_e|)/k_{\parallel 0}$ for all positive n and $v_{g0} > (\omega_{k0} + n|\Omega_e|)/k_{\parallel 0}$ for all negative n . However, in the case of proton resonances, resonant waves are more likely to violate Eq. (29) since $\Omega_p \ll |\Omega_e|$.




Figure 1: The diffusive flux of resonant particles in velocity space under the action of two arbitrary n_1 and n_2 resonances. The dark shaded areas represent isocontours of the VDFs of two particle populations. The red and dark-blue solid curves show the diffusive trajectories, Eq. (28) with $n = n_1$ and $n = n_2$. W_j^{n1} and W_j^{n2} represent the window functions according to Eq. (20), in which the n_1 and n_2 resonances are effective. The light-blue dashed semi-circles correspond to constant-energy contours. The black solid line indicates $v_{\parallel} = v_{g0}$.

Fig. 1 illustrates the diffusive flux of particles experiencing two arbitrary resonances: the n_1 and n_2 resonances for an unstable wave. The dark shaded areas represent isocontours of the VDFs of two particle populations in velocity space. The red and dark-blue solid curves represent the diffusive trajectories according to Eq. (28) with $n = n_1$ and $n = n_2$, assuming that the resonant

wave fulfills Eq. (29). The window functions $W_j^{n_1}$ and $W_j^{n_2}$ describe the v_{\parallel} -ranges in which the n_1 and n_2 resonances are effective. The light-blue dashed semi-circles correspond to contours of constant kinetic energy in the proton rest frame, for which

$$v_{\perp}^2 + v_{\parallel}^2 = \text{const.} \quad (30)$$

In general, the diffusive flux is always directed from higher to lower phase-space densities during the process of stabilization. At point A, resonant particles in $W_j^{n_1}$ diffuse along the red solid curve towards smaller v_{\parallel} . Considering the relative alignment between the diffusive flux and constant-energy contour at point A, the diffusing particles lose kinetic energy. This energy is transferred to the resonant wave, which consequently grows in amplitude. Therefore, this situation corresponds to an instability of the resonant wave. At point B, particles do not diffuse along the red solid curve since this point lies outside $W_j^{n_1}$.

At point C, resonant particles in $W_j^{n_2}$ diffuse along the dark-blue solid curve towards greater v_{\parallel} . Considering the relative alignment between the diffusive flux and the constant-energy contour at point C, the diffusing particles gain kinetic energy. This energy is taken from the resonant wave, which consequently shrinks in amplitude. Therefore, this situation corresponds to damping of the resonant wave and counter-acts the driving of the instability through the n_1 resonance. Because the resonant wave is unstable, the n_1 resonant instability must overcome the counter-acting n_2 resonant damping.

According to Eq. (18), there are three factors that determine the diffusion rate for the action of an n resonance. The first factor is the particle density f_j within W_j^n . The second factor is \tilde{D}_j^n whose magnitude is determined by the polarization properties of the resonant waves. The third factor is the quantity $\hat{G}[k_{\parallel j}^n]f_j/f_j$ which defines the relative alignment between the isocontours of f_i and the diffusive flux along the diffusive trajectory within W_j^n . In Fig. 1, the magnitude of $|\hat{G}[k_{\parallel j}^{n_1}]f_j/f_j|$ at point A is greater than the magnitude of $|\hat{G}[k_{\parallel j}^{n_2}]f_j/f_j|$ at point C.

Since the diffusive flux is directed from higher to lower values of f_j , the quantity $\hat{G}[k_{\parallel j}^n]f_j/f_j$ resolves the ambiguity in the directions of the trajectories for resonant particles. A careful analysis of $\hat{G}[k_{\parallel j}^n]$ with the fulfillment of Eq. (29) shows that, if $(k_{\parallel}/|k_{\parallel}|)(\hat{G}[k_{\parallel j}^n]f_j/f_j) > 0$ at a given resonant velocity, resonant particles diffuse toward a smaller value of v_{\parallel} along the diffusive trajectory while, if $(k_{\parallel}/|k_{\parallel}|)(\hat{G}[k_{\parallel j}^n]f_j/f_j) < 0$ at a given resonant velocity, resonant particles diffuse toward a greater value of v_{\parallel} .

2.4. Numerical Analysis of the Quasi-Linear Diffusion Equation

To simulate the VDF evolution and to compare the diffusion rates between resonances quantitatively, a rigorous numerical analysis of Eq. (10) is necessary. For this purpose, we develop a mathematical approach based on the Crank-Nicolson scheme and present the mathematical details in Appendix A. Our approach is applicable to all two-dimensional diffusion equations with off-diagonal diffusion terms. Our numerical solution, given by Eq. (A28), evolves the VDF under the action of multiple resonances in one time step. We tested our numerical solution by showing that the diffusive flux obeys the predicted diffusion properties discussed in Section 2.3.

3. FAST-MAGNETOSONIC/WHISTLER WAVE AND ELECTRON-STRAHL SCATTERING

As an example, we apply our model developed in Section 2 to an electron resonant instability in the solar wind. The fast-magnetosonic/whistler (FM/W) wave propagating in the anti-sunward direction and with an angle of $\sim 60^\circ$ with respect to the background magnetic field scatters the electron strahl (Vasko et al. 2019; Verscharen et al. 2019). Since this prediction is based on linear theory, our quasi-linear framework is appropriate for demonstrating the action of this instability on the electron strahl.

3.1. Linear Dispersion Relation

To find the characteristics of the unstable oblique FM/W wave, we numerically solve the hot-plasma dispersion relation with the NHDS code (Verscharen & Chandran 2018). We use the same plasma parameters as Verscharen et al. (2019), which are, notwithstanding the wide range of natural variation, representative for the average electron parameters in the solar wind (Wilson et al. 2019). We assume that the initial plasma consists of isotropic Maxwellian protons, core electrons and strahl electrons. The subscripts p , e , c and s indicate protons, electrons, electron core and electron strahl, respectively.

We choose our coordinate system so that the anti-sunward and obliquely propagating FM/W waves have $k_{\parallel} > 0$. We set $\beta_c = \beta_p = 1$ and $\beta_s = 0.174$, where $\beta_j \equiv (8\pi n_j k_B T_j)/B_0^2$, n_j and T_j are the density and temperature of species j , and k_B is the Boltzmann constant. We set $n_p = n_e$, $n_c = 0.92n_p$, $n_s = 0.08n_p$, $T_c = T_p$, and $T_s = 2T_p$. In the proton rest frame, we set $n_c U_c + n_s U_s = 0$. We initialize the core and strahl bulk velocity with $U_c/v_{Ae} = -0.22$ and $U_s/v_{Ae} = 2.52$ where $v_{Ae} \equiv B_0/\sqrt{4\pi n_e m_e}$ is the electron Alfvén velocity. NHDS finds that, under these plasma parameters, $\gamma_k > 0$




Figure 2: NHDS solutions provide γ_k (dashed curves, axis on the left) and ω_k (solid curves, axis on the right) as functions of the k_{\parallel} -component of the wavevector \mathbf{k} . We show solutions for $\theta_0 = 51^\circ$, $\theta_0 = 55^\circ$, and $\theta_0 = 59^\circ$.

at angles between $\theta_0 = 51^\circ$ and $\theta_0 = 67^\circ$. Our strahl bulk velocity then provides a maximum growth rate of $\gamma_k/|\Omega_e| = 10^{-3}$ (Verscharen et al. 2019).

Fig. 2 shows γ_k and ω_k as functions of the k_{\parallel} -component of the wavevector \mathbf{k} for three different θ_0 . The oblique FM/W instability has its maximum growth rate at $\theta_0 = 55^\circ$, while $\gamma_k > 0$ for $0.21 \lesssim k_{\parallel}v_{Ae}/|\Omega_e| \lesssim 0.28$ which is the parallel unstable \mathbf{k} -spectrum. As defined in Section 2.1, we acquire $k_{\parallel 0}v_{Ae}/|\Omega_e| \approx 0.245$. This value with Eqs. (7)-(9) leads to $k_{\perp 0}v_{Ae}/|\Omega_e| = 0.35$, $\omega_{k0}/|\Omega_e| \approx 0.07$ and $v_{g0}/v_{Ae} \approx 0.86$. We also acquire $\sigma_{\parallel 0}v_{Ae}/|\Omega_e| \approx 0.035$ and $\sigma_{\perp 0}v_{Ae}/|\Omega_e| \approx 0.05$ from the obtained unstable \mathbf{k} -spectrum.

3.2. Theoretical Description of the Quasi-Linear Diffusion in the FM/W Instability

Using the wave and plasma parameters from the previous section, we describe the electron strahl and core diffusion in velocity space. In our analysis, we only consider $n = +1, -1$ and 0 resonances, neglecting higher- n resonances due to their negligible contributions.

Upon substituting our wave parameters into Eq. (20), we quantify the dimensionless window functions $W_e^n v_{Ae}$ with $n = +1, -1$ and 0 . In Fig. 3, the red, dark-blue and orange lines represent $W_e^{+1}v_{Ae}$, $W_e^{-1}v_{Ae}$ and $W_e^0v_{Ae}$, which are maximum at $v_{\parallel res}^+/v_{Ae} = 4.37$, $v_{\parallel res}^-/v_{Ae} = -3.8$ and $v_{\parallel res}^0/v_{Ae} = 0.29$, respectively. We reiterate that the superscripts indicate the n resonance. The black line indicates $v_{\parallel} = v_{g0}$. Each $W_e^n v_{Ae}$ shows the v_{\parallel} -range in which the quasi-linear diffusion through each resonance is effective. We note that $W_e^n v_{Ae}$ for the three resonances have a different width in v_{\parallel} -space and maximum value due to a different magnitude of $|v_{\parallel res}^n - v_{g0}|$ (see Eq. (24)).

By substituting our wave parameters into Eq. (28), the diffusive trajectories for the $n = +1, -1$ and 0 resonances are given by

$$(v_{\perp}/v_{Ae})^2 + 1.16 (v_{\parallel}/v_{Ae} - 0.86)^2 = const, \quad (31)$$

$$(v_{\perp}/v_{Ae})^2 + 0.88 (v_{\parallel}/v_{Ae} - 0.86)^2 = const, \quad (32)$$

and

$$(v_{\perp}/v_{Ae})^2 = const. \quad (33)$$

Eqs. (31) and (32) describe ellipses with their axes oriented along the v_{\perp} - and v_{\parallel} -directions. In Eq. (33), the perpendicular velocity of resonant particles is constant.

Fig. 4 illustrates the electron diffusion from these three resonances. We show the v_{\parallel} -ranges in which these three resonances are effective according to $W_e^{+1}v_{Ae}$, $W_e^{-1}v_{Ae}$ and $W_e^0v_{Ae}$ from Fig. 3. The red, dark-blue and orange solid lines represent the contours given by Eqs. (31)-(33), respectively. The light-blue dashed semi-circles correspond to constant-energy contours in the proton rest frame (see Eq. (30)). The black line indicates $v_{\parallel} = v_{g0}$. For the initial strahl and core VDF, we apply the plasma parameters in Section 3.1 to the dimensionless Maxwellian distribution

$$f_j^M = \frac{n_j v_{Ae}^3}{\pi^{3/2} n_p v_{th,j}^3} \exp \left[-\frac{v_{\perp}^2 + (v_{\parallel} - U_j)^2}{v_{th,j}^2} \right], \quad (34)$$

where $v_{th,j} \equiv \sqrt{2k_B T_j/m_j}$. The red and blue areas in Fig. 4 represent f_s^M and f_c^M which are normalized by the maximum value of f_c^M and plotted up to a value of 10^{-5} . In this normalization, Fig. 4 does not reflect the relative density between both electron species.

Due to the v_{\parallel} -profile of W_e^{+1} , the $n = +1$ resonance has a significant effect on f_s^M . As discussed in Section 2.3, since $(k_{\parallel}/|k_{\parallel}|)(\hat{G}[k_{\parallel}^{+1}]f_s^M/f_c^M) > 0$, this resonance leads to the diffusion of the resonant strahl electrons in W_e^{+1} along the red arrows. According to Eq. (30), the phase-space trajectory of particles that diffuse without a change in kinetic energy is described by

$$\left(\frac{dv_{\perp}}{dv_{\parallel}} \right)_E = -\frac{v_{\parallel}}{v_{\perp}}. \quad (35)$$

According to Eq. (31), the phase-space trajectory of resonant particles fulfilling the $n = +1$ resonance, indicated by superscript $+1$, is described by

$$\left(\frac{dv_{\perp}}{dv_{\parallel}} \right)^{+1} = -1.16 \frac{v_{Ae}}{v_{\perp}} \left(\frac{v_{\parallel}}{v_{Ae}} - 0.86 \right). \quad (36)$$

Evaluating Eqs. (35) and (36) in W_e^{+1} shows that $|(dv_{\perp}/dv_{\parallel})^{+1}| < |(dv_{\perp}/dv_{\parallel})_E|$ for the resonant electrons.




Figure 3: The red, dark-blue and orange plots illustrate $W_e^{+1}v_{Ae}$, $W_e^{-1}v_{Ae}$ and $W_e^0v_{Ae}$ for the oblique FM/W wave. The black solid line represents $v_{||} = v_{g0}$. Each $W_e^n v_{Ae}$ shows the $v_{||}$ -range in which each resonance is effective. Each $W_e^n v_{Ae}$ has a different width in $v_{||}$ -space and maximum value due to a different magnitude of $|v_{||res}^n - v_{g0}|$ (see Eq. (24)).




Figure 4: The red, dark-blue and orange arrows illustrate the diffusive flux according to the $n = +1$, -1 and 0 resonances for the oblique FM/W instability. The red and dark-blue filled semi-circles represent isocontours of the strahl and core VDF. This figure does not reflect the relative densities of both electron species. The light-blue dashed semi-circles correspond to constant-energy contours. The black solid line indicates $v_{||} = v_{g0}$.

Therefore, resolving the ambiguity in the directions of the trajectories, the distance of resonant strahl electrons from the origin of the coordinate system decreases. This decrease in $v_{\perp}^2 + v_{||}^2$ represents a loss of kinetic energy of the resonant strahl electrons. The $n = +1$ resonance, therefore, contributes to the driving of the FM/W instability.

Due to the $v_{||}$ -profile of W_e^{-1} , the $n = -1$ resonance has a significant effect on f_c^M . Since $(k_{||}/|k_{||}|)(\hat{G}[k_{||e}^{-1}]f_c^M/f_c^M) < 0$, this resonance leads to the diffusion of the resonant core electrons in W_e^{-1} along the dark-blue arrows. According to Eq. (32), the phase-

space trajectory of resonant particles fulfilling the $n = -1$ resonance, indicated by superscript -1 , is described by

$$\left(\frac{dv_{\perp}}{dv_{||}}\right)^{-1} = -0.88 \frac{v_{Ae}}{v_{\perp}} \left(\frac{v_{||}}{v_{Ae}} - 0.86\right). \quad (37)$$

Evaluating Eqs. (35) and (37) in W_e^{-1} shows that $|(dv_{\perp}/dv_{||})^{-1}| > |(dv_{\perp}/dv_{||})_E|$ for the resonant electrons. Therefore, resolving the ambiguity in the directions of the trajectories, the distance of resonant core electrons from the origin of the coordinate system increases. This increase in $v_{\perp}^2 + v_{||}^2$ represents a gain of kinetic energy of the resonant core electrons. The $n = -1$ resonance, therefore, counter-acts the FM/W instability through the $n = +1$ resonance.

Due to the $v_{||}$ -profile of W_e^0 , the $n = 0$ resonance has a significant effect on electrons in the $v_{||}$ -range in which $f_c^M > f_s^M$ and $\partial f_c^M/\partial v_{||} < 0$. Since $(k_{||}/|k_{||}|)(\hat{G}[k_{||e}^0]f_c^M/f_c^M) < 0$, the resonant electrons in W_e^0 diffuse along the yellow arrows. Because the distance of these electrons from the origin of the coordinate system increases, these resonant electrons diffuse towards greater kinetic energies. This diffusion removes energy from the resonant FM/W waves and thus counter-acts the driving of the FM/W instability through the $n = +1$ resonance.

The illustration in Fig. 4 describes the nature of the quasi-linear diffusion through the $n = +1$, -1 and 0 resonances in velocity space. It does not give any information regarding the relative strengths of the diffusion rates between the three resonances. Since the FM/W wave is unstable according to linear theory, the $n = +1$ resonant instability must dominate over any counter-acting contributions from the $n = -1$ and 0 resonances.

3.3. Numerical Description of the Quasi-Linear Diffusion in the FM/W Instability

We use our numerical analysis in Eq. (A28) to simulate the quasi-linear diffusion through the $n = +1$, -1 and 0 resonances, predicted in Section 3.2. According to the definitions in Appendix A, we select the discretization parameters $N_v = 45$, $v_{\perp\max}/v_{Ae} = v_{||\max}/v_{Ae} = 7$ and $|\Omega_e|\Delta t = 1$. For the computation of Eq. (A28), we use the parameters of resonant FM/W waves presented in Section 3.1 and quantify $\tilde{D}_e^{\pm 1}$ and \tilde{D}_e^0 in Eq. (19).

In \tilde{D}_e^n for each resonance, we only consider the J_0 term in ψ_{j0}^n , neglecting higher-order Bessel functions due to their small contributions. Our NHDS solutions show that $|E_0^y| \approx 0.2|E_0^x|$ and $|E_0^z| \approx 0.14|E_0^x|$ in the unstable \mathbf{k} -spectrum. Then, we set $|E_0^R| \approx |E_0^L| \approx 0.72|E_0^x|$. Faraday's law yields $E_0^x \approx [\omega_{k0}/(k_{||0}c)]B_0^y$ when neglecting the small contributions from E_0^z terms. This allows us to express E_0^x through B_0^y in ψ_{j0}^n , where

B_0^y represents the peak amplitude of the wave magnetic-field fluctuations. For simplicity, we assume that B_0^y is constant in time during the quasi-linear diffusion. Under these assumptions, we acquire

$$\begin{aligned} \tilde{D}_e^{\pm 1} &\approx W_e^{\pm 1} \frac{0.52\pi^2 |\Omega_e|^2 v_\perp^2}{\sigma_{\parallel 0} \sigma_{\perp 0}^2} \left[\frac{B_0^y \omega_{k0}}{B_0 k_{\parallel 0}} \right]^2 \\ &\times \int_0^\infty J_0(\rho_e)^2 \exp \left[-\frac{(k_\perp - k_{\perp 0})^2}{\sigma_{\perp 0}^2} \right] k_\perp dk_\perp, \end{aligned} \quad (38)$$

and

$$\begin{aligned} \tilde{D}_e^0 &\approx W_e^0 \frac{0.04\pi^2 |\Omega_e|^2 v_\parallel^2}{\sigma_{\parallel 0} \sigma_{\perp 0}^2} \left[\frac{B_0^y \omega_{k0}}{B_0 k_{\parallel 0}} \right]^2 \\ &\times \int_0^\infty J_0(\rho_e)^2 \exp \left[-\frac{(k_\perp - k_{\perp 0})^2}{\sigma_{\perp 0}^2} \right] k_\perp dk_\perp, \end{aligned} \quad (39)$$

where the relative amplitude B_0^y/B_0 is a free parameter and we set $B_0^y/B_0 = 0.001$. Then, we apply Eqs. (38) and (39) to Eq. (A28).

We initialize our numerical computation with the same f_s^M and f_c^M as defined in Section 3.2. Fig. 5a represents $f_e = f_c^M + f_s^M$, normalized by the maximum value of f_c^M and plotted up to a value of 10^{-5} . Fig. 5b shows f_e evolved through the $n = +1$, -1 and 0 resonances, resulting from the iterative calculation of Eq. (A28). Considering the maximum value of the instability's growth rate, $\gamma_k/|\Omega_e| = 4.8 \times 10^{-3}$ in Fig. 2, we finish the evaluation of our numerical computation at $|\Omega_e|t = 3 \times 10^2$ which corresponds to $\gamma_k t \sim 1$ and thus a reasonable total growth of the unstable FM/W waves.

The strahl electrons at around $v_\parallel/v_{Ae} \approx 4.4$ diffuse through the $n = +1$ resonance, as theoretically predicted in Fig. 4. This diffusion increases the pitch-angle of the resonant strahl electrons and generates a strong pitch-angle gradient at $v_\parallel/v_{Ae} \approx 3.8$. During this process, the v_\perp of the scattered strahl electrons increases while their v_\parallel decreases.

Since the longitudinal component of the electric-field fluctuations is much weaker than their transverse components, the diffusion through the $n = 0$ resonance is not noticeable over the modeled time interval. The diffusion through the $n = -1$ resonance is not noticeable even though \tilde{D}_e^{-1} and \tilde{D}_e^{+1} have a similar magnitude. This is because the magnitude of $|\hat{G}[k_{\parallel e}^{-1}]f_e/f_e|$ in W_e^{-1} is much smaller than the magnitude of $|\hat{G}[k_{\parallel e}^{+1}]f_e/f_e|$ in W_e^{+1} , as discussed in Section 2.3, and the number of core electrons in W_e^{-1} is very small (see Fig. 4 and 5).

4. THE SECONDARY EFFECT OF COULOMB COLLISIONS

Since the collisionless action of resonant wave-particle instabilities often form strong pitch-angle gradients (see,




Figure 5: Fig. 5a: the initial electron VDF; Fig. 5b: the electron VDF evolved through the $n = +1$, -1 and 0 resonances. Compared to Fig. 4, only the effect of the $n = +1$ resonance is noticeable during the time $\gamma_k t \sim 1$. It causes a significant pitch-angle gradient at $v_\parallel/v_{Ae} \approx 3.8$ through the scattering of strahl electrons. An animation of this figure is available in the HTML version of the article. The animation shows the time evolution of the distribution function from $|\Omega_e|t = 0$ to $|\Omega_e|t = 3 \times 10^2$. During this evolution, the strahl scattering towards larger v_\perp is visible.

for example, Fig. 5), the collisions can be enhanced in the plasma. Therefore, a more realistic evolution of the total electron VDF must account for the action of Coulomb collisions of strahl electrons with core electrons and protons. For this purpose, we adopt the Fokker-Planck equation given by Ljepojevic et al. (1990) with Rosenbluth potentials (Rosenbluth et al. 1957), and normalize it in our dimensionless system of units as

$$\begin{aligned} \left(\frac{\partial f_j}{\partial t} \right)_{\text{Fokker-Planck}} &= \sum_b \Gamma_{jb} \left\{ 4\pi \frac{m_j}{m_b} f_b f_j \right. \\ &\left. + \frac{\partial h}{\partial v^\alpha} \frac{\partial f_j}{\partial v^\alpha} + \frac{1}{2} \frac{\partial^2 g}{\partial v^\alpha \partial v^\beta} \frac{\partial^2 f_j}{\partial v^\alpha \partial v^\beta} \right\}, \end{aligned} \quad (40)$$

where

$$g(\mathbf{v}) \equiv \int f_b(\mathbf{v}') |\mathbf{v} - \mathbf{v}'| d^3 \mathbf{v}', \quad (41)$$

$$h(\mathbf{v}) \equiv \frac{m_b - m_j}{m_b} \int f_b(\mathbf{v}') |\mathbf{v} - \mathbf{v}'|^{-1} d^3 \mathbf{v}', \quad (42)$$

and

$$\Gamma_{jb} = \frac{4\pi n_b}{v_{Ae}^3 |\Omega_e|} \left(\frac{Z_j Z_b q_j^2}{m_j} \right)^2 \ln \Lambda_{jb}. \quad (43)$$

The subscript b indicates the species of background particles, with which the particles of species j Coulomb-collide. The quantity $\ln \Lambda_{jb}$ is the Coulomb logarithm and typically $\ln \Lambda_{jb} \approx 25$ in space plasmas. The parameters Z_j and Z_b are the atomic masses of a particle of species j and b . The superscripts α and β indicate the component of the velocity in cylindrical coordinates and the summation convention holds.

We assume that the timescale of Coulomb collisions is much longer than the timescale of the quasi-linear diffusion in the solar wind under our set of parameters. This assumption allows us to model the resonant wave-particle instability first and to use the resulting VDF as the input for the model of the subsequent, secondary effects of collisions.

Based on our mathematical approach presented in Appendix A, we numerically solve the Fokker-Planck equation, Eq. (40), in Appendix B. We tested our numerical solutions, Eq. (B31), by showing that a set of arbitrary test VDFs diffuses toward f_b with time.

For the computation of Eq. (B31), we set isotropic Maxwellian electron-core and proton VDFs as background species, $f_b = f_c^M$ and $f_b = f_p^M$, for which we apply the plasma parameters presented in Section 3.1 to Eq. (34). In this numerical computation, we select the discretization parameters $N_v = 45$, $v_{\perp \max}/v_{Ae} = v_{\parallel \max}/v_{Ae} = 7$ and $|\Omega_e| \Delta t = 10$. Moreover, we set $B_0 = 5 \times 10^{-4} G$ and $n_b = 10^2 cm^{-3}$ in Eq. (43), which are representative for the conditions in the solar wind at a distance of 0.3 au from the Sun. We initialize f_j with the electron-strahl VDF as f_s evolved by the oblique FM/W instability at time $|\Omega_e|t = 3 \times 10^2$ from our quasi-linear analysis. In this setup, our initial electron VDF for the Coulomb collision analysis is same as the electron VDF shown in Fig. 5b.

The iterative calculation of Eq. (B31) results in the time evolution of the electron-strahl VDF under the action of Coulomb collisions with core electrons and protons. The result of this computation at the time $|\Omega_e|t = 7 \times 10^7$ is shown in Fig. 6.

A detailed comparison of the distribution function before (Fig. 6a) and after (Fig. 6b) our calculation of the effect of Coulomb collisions reveals that Coulomb collisions relax the strong pitch-angle gradient at $v_{\parallel}/v_{Ae} \approx 3.8$, which resulted from the action of the oblique FM/W instability. However, the Coulomb collisions are only




Figure 6: Fig. 6a: the electron VDF as initial condition for our collision analysis; Fig. 6b: the electron VDF evolved through Coulomb collisions of strahl electrons with core electrons and protons. The strong pitch-angle gradient at $v_{\parallel}/v_{Ae} \approx 3.8$ (shown in Fig. 6a and Fig. 5b) is relaxed through the Coulomb collisions. However, the required time for a noticeable collisional effect on that gradient is around 10^5 times longer than the timescale of the strahl scattering. An animation of this figure is available in the HTML version of the article. The animation shows the time evolution of the distribution function from $|\Omega_e|t = 3 \times 10^2$ to $|\Omega_e|t = 7 \times 10^7$. During this evolution, the collisional smoothing of the pitch-angle gradients is visible.

capable of affecting strong pitch-angle gradients in the modified electron VDF under our plasma parameters. In addition, the required time for a noticeable collisional effect on this pitch-angle gradient is of order 10^5 times longer than the characteristic timescale of the quasi-linear diffusion.

5. DISCUSSION FOR THE STRAHL SCATTERING

The numerical computation of Eq. (17) shows that dH/dt is negative and asymptotically tends towards zero as the electron VDF evolves through the oblique FM/W

instability and the counter-acting damping effects until the time $|\Omega_e|t = 3 \times 10^2$, which is presented in Fig. 5. Therefore, our quasi-linear diffusion model reflects the stabilization of the particle VDF through the participating wave-particle resonances.

During the action of the oblique FM/W instability, the scattered strahl electrons reduce their collimation along the \mathbf{B}_0 -direction and become more isotropic. Even though this instability does not cause significant strahl scattering, we argue that it contributes to the initial formation of the halo population. However, other mechanisms must be considered to account for the full strahl scattering in agreement with observations (Gurgiolo et al. 2012; Gurgiolo & Goldstein 2016).

Alternative models describing Coulomb-collisional effects on the strahl VDF suggest that an anomalous-diffusion process must be considered in order to achieve an agreement with observations (Lemons & Feldman 1983; Horaites et al. 2018, 2019). We note that our analysis includes the subsequent action of Coulomb collisions after the action of collisionless wave-particle resonances assuming plasma parameters consistent with the solar wind at a distance of 0.3 au from the Sun. Our collisional effects are similar to those proposed by Vocks et al. (2005). However, our model predicts that the collisional relaxation is so subtle that the strahl scattering through collisions is barely noticeable for the analyzed phase of the VDF evolution.

The clear separation of timescales between wave-particle effects and Coulomb-collisional effects complicates the description of the VDF evolution on heliospheric scales, since other processes act on comparable timescales. These additional processes, which our analysis neglects, include turbulence, shocks, plasma mixing, plasma expansion, and magnetic focusing (Yoon et al. 2012; Ryu et al. 2007; Feldman et al. 1983; Fitzenreiter et al. 2003; Tang et al. 2020). A complete model for the radial evolution of the VDF must quantify and account for these processes as well. In the context of our work, these processes can potentially push a VDF that has undergone stabilization as shown in Fig. 5b into the unstable regime again. In this case, dH/dt in Eq. (17) returns to a non-zero value, which signifies a new onset of wave-particle resonances and further scattering of resonant particles.

6. CONCLUSIONS

Wave-particle resonances are important plasma-physics processes in many astrophysical plasmas. Often, fully non-linear simulations with codes solving the equations of kinetic plasma theory are used to model the evolution of the distribution function under the action of wave-particle resonances. However, quasi-linear theory

augments this approach since it allows us to study the contributions of different processes to these resonances. Therefore, quasi-linear theory is a very helpful tool to improve our understanding of wave-particle resonances in astrophysical plasmas.

We propose a quasi-linear diffusion model for any generalized wave-particle instability. We analyze the quasi-linear diffusion equation by expressing the electric field of an arbitrary unstable and resonant wave mode as a Gaussian wave packet. From Boltzmann's H -theorem with our quasi-linear analysis, we define a window function that determines the specific velocity-space range in which a dominant wave-particle instability and counter-acting damping contributions are effective. This window function is the consequence of the localized energy density of our Gaussian wave packet both in configuration space and in wavevector space.

Moreover, we derive a relation describing the diffusive trajectories of the resonant particles for such an instability in velocity space. These trajectories evolve the particle VDF into a stable state in which no further quasi-linear diffusion occurs. Therefore, our theoretical model illustrates the diffusion and stabilization which resonant particles, depending on their location in velocity space, experience in wave-particle resonances.

For the computational quantification of our theoretical model, we introduce a mathematical approach based on the Crank-Nicolson scheme to numerically solve the full quasi-linear diffusion equation. We highlight that this mathematical approach applies to all general two-dimensional diffusion equations, including those with off-diagonal diffusion terms.

As an example, we apply our model to the oblique FM/W instability that scatters strahl electrons in the solar wind. Our model shows that the $n = +1$ resonant instability of FM/W waves propagating with an angle of $\sim 55^\circ$ with respect to the background magnetic field scatters strahl electrons towards larger v_\perp and smaller v_\parallel . The strahl scattering instability through the $n = +1$ resonance dominates over the counter-acting damping contributions through the $n = -1$ and $n = 0$ resonances. This instability creates a strong pitch-angle gradient in the electron-strahl VDF.

By numerically solving the Fokker-Planck equation with our mathematical approach, we show that Coulomb collisions of strahl electrons with core electrons and protons relax this strong pitch-angle gradient on a timescale 10^5 times longer than the timescale of the collisionless strahl scattering. This finding suggests that collisional effects are negligible in the strahl-driven oblique FM/W instability, which is a representative example for a resonant wave-particle instability in the solar wind.

Our predicted evolution of the electron VDF is consistent with the observed formation of a proto-halo through strahl scattering (Gurgiolo et al. 2012). However, further observations are ambiguous regarding the exact source of the proto-halo (Gurgiolo & Goldstein 2016). Future high-resolution electron observations with Solar Orbiter and Parker Solar Probe at different distances from the Sun may help us to resolve these ambiguities.

Our general quasi-linear diffusion model applies to all non-relativistic collisionless plasmas, such as planetary magnetosphere (e.g. Mourenas et al. 2015). It also applies to other types of wave-particle instabilities in plasmas

such as the resonant instabilities driven by temperature anisotropy or by relative drift. We especially note that our model is capable of describing ion-driven instabilities.

ACKNOWLEDGMENTS

We appreciate helpful discussions with Georgios Nicolaou, Konstantinos Horaites and Jung Joon Seough. D.V. is supported by the STFC Ernest Rutherford Fellowship ST/P003826/1. D.V., R.T.W., and A.N.F. are supported by STFC Consolidated Grant ST/S000240/1.

APPENDIX

A. NUMERICAL ANALYSIS OF QUASI-LINEAR DIFFUSION EQUATION

Eq. (10) is a second-order differential equation which includes cross-derivative operators such as $\partial^2/\partial v_{\parallel}\partial v_{\perp}$. In order to simultaneously evaluate the $\partial^2/\partial v_{\parallel}\partial v_{\perp}$ operators with the $\partial^2/\partial v_{\parallel}^2$ and $\partial^2/\partial v_{\perp}^2$ operators in Eq. (10), we divide velocity space into $2N_v \times 2N_v$ steps with equal step sizes of $\Delta v/2$ by defining the outer boundaries of velocity space as $\pm v_{\perp\max}$ and $\pm v_{\parallel\max}$. The v_{\perp} -index M and the v_{\parallel} -index N both step through $1, 3/2, 2, \dots, N_v, N_v + 1/2$. We define the discrete velocity coordinates as $v_{\perp M} = -v_{\perp\max} + (M - 1)\Delta v$ and $v_{\parallel N} = -v_{\parallel\max} + (N - 1)\Delta v$. We note that this definition introduces negative v_{\perp} -values that, although they simplify our numerical analysis, we neglect in our computational results. We divide the time t with equal step sizes of Δt and the t -index T steps through $1, 2, 3, \dots$. We define the discrete time as $t^T = (T - 1)\Delta t$. We then define the discrete VDF as $f_{M,N}^T = f_j(v_{\perp M}, v_{\parallel N}, t^T)$. For the discretization of the velocity derivatives, we adopt the two-point central difference operator (Gilat & Subramaniam 2011)

$$\frac{\partial f_j(v_{\perp M}, v_{\parallel N}, t^T)}{\partial v_{\perp}} \approx \frac{f_{M+1/2,N}^T - f_{M-1/2,N}^T}{\Delta v}, \quad (\text{A1})$$

and

$$\frac{\partial f_j(v_{\perp M}, v_{\parallel N}, t^T)}{\partial v_{\parallel}} \approx \frac{f_{M,N+1/2}^T - f_{M,N-1/2}^T}{\Delta v}. \quad (\text{A2})$$

For the discretization of the time derivative, we adopt the forward difference operator

$$\frac{\partial f_j(v_{\perp M}, v_{\parallel N}, t^T)}{\partial t} \approx \frac{f_{M,N}^{T+1} - f_{M,N}^T}{\Delta t}. \quad (\text{A3})$$

By using Eqs. (A2) and (A1), we discretize the right-hand side of Eq. (10) and express it as $(\partial f/\partial t)_{M,N}^T$

$$\begin{aligned} \left(\frac{\partial f}{\partial t}\right)_{M,N}^T &\equiv \sum_{n=-\infty}^{\infty} \int \left[\left(1 - \frac{k_{\parallel} v_{\parallel N}}{\omega_{k0} + v_{g0}(k_{\parallel} - k_{\parallel 0})}\right) \frac{1}{v_{\perp M}} \frac{D_{M+1/2,N}^n [\hat{G}f]_{M+1/2,N}^T - D_{M-1/2,N}^n [\hat{G}f]_{M-1/2,N}^T}{\Delta v} \right. \\ &\quad \left. + \frac{k_{\parallel}}{\omega_{k0} + v_{g0}(k_{\parallel} - k_{\parallel 0})} \frac{D_{M,N+1/2}^n [\hat{G}f]_{M,N+1/2}^T - D_{M,N-1/2}^n [\hat{G}f]_{M,N-1/2}^T}{\Delta v} \right] d^3 \mathbf{k}, \end{aligned} \quad (\text{A4})$$

where

$$[\hat{G}f]_{M,N}^T \equiv \left(1 - \frac{k_{\parallel} v_{\parallel N}}{\omega_{k0} + v_{g0}(k_{\parallel} - k_{\parallel 0})}\right) \frac{1}{v_{\perp M}} \frac{f_{M+1/2,N}^T - f_{M-1/2,N}^T}{\Delta v} + \frac{k_{\parallel}}{\omega_{k0} + v_{g0}(k_{\parallel} - k_{\parallel 0})} \frac{f_{M,N+1/2}^T - f_{M,N-1/2}^T}{\Delta v}, \quad (\text{A5})$$

and

$$D_{M,N}^n \equiv D_j^n \Big|_{\substack{v_{\perp} = v_{\perp M} \\ v_{\parallel} = v_{\parallel N}}} \quad (\text{A6})$$

According to the Crank-Nicolson scheme (Iserles 2008), the full discretization of Eq. (10) in its time and velocity derivatives is then given by

$$f_{M,N}^{T+1} - \frac{\Delta t}{2} \left(\frac{\partial f}{\partial t} \right)_{M,N}^{T+1} = f_{M,N}^T + \frac{\Delta t}{2} \left(\frac{\partial f}{\partial t} \right)_{M,N}^T. \quad (\text{A7})$$

By using Eqs. (A4)-(A6) and resolving the δ -functions in $D_{M\pm 1/2,N}^n$ and $D_{M,N\pm 1/2}^n$ through the k_{\parallel} -integral, we rewrite Eq. (A7) as

$$\begin{aligned} & f_{M,N}^{T+1} - \sum_{n=-\infty}^{\infty} \left\{ \frac{\mu}{2} P_{M,N}^n \tilde{D}_{M+1/2,N}^n \left[P_{M+1/2,N}^n \left(f_{M+1,N}^{T+1} - f_{M,N}^{T+1} \right) + Q_N^n \left(f_{M+1/2,N+1/2}^{T+1} - f_{M+1/2,N-1/2}^{T+1} \right) \right] \right. \\ & \quad - \frac{\mu}{2} P_{M,N}^n \tilde{D}_{M-1/2,N}^n \left[P_{M-1/2,N}^n \left(f_{M,N}^{T+1} - f_{M-1,N}^{T+1} \right) + Q_N^n \left(f_{M-1/2,N+1/2}^{T+1} - f_{M-1/2,N-1/2}^{T+1} \right) \right] \\ & \quad + \frac{\mu}{2} Q_{N+1/2}^n \tilde{D}_{M,N+1/2}^n \left[P_{M,N+1/2}^n \left(f_{M+1/2,N+1/2}^{T+1} - f_{M-1/2,N+1/2}^{T+1} \right) + Q_{N+1/2}^n \left(f_{M,N+1}^{T+1} - f_{M,N}^{T+1} \right) \right] \\ & \quad \left. - \frac{\mu}{2} Q_{N-1/2}^n \tilde{D}_{M,N-1/2}^n \left[P_{M,N-1/2}^n \left(f_{M+1/2,N-1/2}^{T+1} - f_{M-1/2,N-1/2}^{T+1} \right) + Q_{N-1/2}^n \left(f_{M,N}^{T+1} - f_{M,N-1}^{T+1} \right) \right] \right\} \\ & = f_{M,N}^T + \sum_{n=-\infty}^{\infty} \left\{ \frac{\mu}{2} P_{M,N}^n \tilde{D}_{M+1/2,N}^n \left[P_{M+1/2,N}^n \left(f_{M+1,N}^T - f_{M,N}^T \right) + Q_N^n \left(f_{M+1/2,N+1/2}^T - f_{M+1/2,N-1/2}^T \right) \right] \right. \\ & \quad - \frac{\mu}{2} P_{M,N}^n \tilde{D}_{M-1/2,N}^n \left[P_{M-1/2,N}^n \left(f_{M,N}^T - f_{M-1,N}^T \right) + Q_N^n \left(f_{M-1/2,N+1/2}^T - f_{M-1/2,N-1/2}^T \right) \right] \\ & \quad + \frac{\mu}{2} Q_{N+1/2}^n \tilde{D}_{M,N+1/2}^n \left[P_{M,N+1/2}^n \left(f_{M+1/2,N+1/2}^T - f_{M-1/2,N+1/2}^T \right) + Q_{N+1/2}^n \left(f_{M,N+1}^T - f_{M,N}^T \right) \right] \\ & \quad \left. - \frac{\mu}{2} Q_{N-1/2}^n \tilde{D}_{M,N-1/2}^n \left[P_{M,N-1/2}^n \left(f_{M+1/2,N-1/2}^T - f_{M-1/2,N-1/2}^T \right) + Q_{N-1/2}^n \left(f_{M,N}^T - f_{M,N-1}^T \right) \right] \right\}, \end{aligned} \quad (\text{A8})$$

where

$$P_{M,N}^n \equiv \frac{n\Omega_j [v_{\parallel N} - v_{g0}]}{[(\omega_{k0} - k_{\parallel 0} v_{g0}) v_{\parallel N} - n\Omega_j v_{g0}] v_{\perp M}}, \quad (\text{A9})$$

$$Q_N^n \equiv \frac{\omega_{k0} - k_{\parallel 0} v_{g0} - n\Omega_j}{(\omega_{k0} - k_{\parallel 0} v_{g0}) v_{\parallel N} - n\Omega_j v_{g0}}, \quad (\text{A10})$$

$$\tilde{D}_{M,N}^n \equiv \tilde{D}_j^n \Big|_{\substack{v_{\perp} = v_{\perp M} \\ v_{\parallel} = v_{\parallel N}}}, \quad (\text{A11})$$

and $\mu \equiv \Delta t / (\Delta v)^2$. Eq. (A8) is a two-dimensional set of algebraic equations, the solution of which, $f_{M,N}^{T+1}$, for all v_{\perp} - and v_{\parallel} -indexes describes the VDF at time $T + 1$ based on $f_{M,N}^T$ for all v_{\perp} - and v_{\parallel} -indexes.

In order to transform Eq. (A8) into a single matrix equation with a tridiagonal matrix, we introduce the concept of a double matrix. On both sides of Eq. (A8), we group the terms by the same v_{\perp} -index in the VDF and rearrange these groups in increasing order in v_{\perp} -index. In each group, we then rearrange terms in increasing order in v_{\parallel} -index in the VDF. Then, we have

$$\begin{aligned} & -\eta(\mu)_{M,N}^{(1)} f_{M-1,N}^{T+1} - \xi(\mu)_{M,N}^{(12)} f_{M-1/2,N-1/2}^{T+1} + \xi(\mu)_{M,N}^{(11)} f_{M-1/2,N+1/2}^{T+1} - \alpha(\mu)_{M,N}^{(2)} f_{M,N-1}^{T+1} + \alpha(\mu)_{M,N} f_{M,N}^{T+1} \\ & - \alpha(\mu)_{M,N}^{(1)} f_{M,N+1}^{T+1} + \xi(\mu)_{M,N}^{(22)} f_{M+1/2,N-1/2}^{T+1} - \xi(\mu)_{M,N}^{(21)} f_{M+1/2,N+1/2}^{T+1} - \eta(\mu)_{M,N}^{(2)} f_{M+1,N}^{T+1} \\ & = \\ & -\eta(-\mu)_{M,N}^{(1)} f_{M-1,N}^T - \xi(-\mu)_{M,N}^{(12)} f_{M-1/2,N-1/2}^T + \xi(-\mu)_{M,N}^{(11)} f_{M-1/2,N+1/2}^T - \alpha(-\mu)_{M,N}^{(2)} f_{M,N-1}^T + \alpha(-\mu)_{M,N} f_{M,N}^T \\ & - \alpha(-\mu)_{M,N}^{(1)} f_{M,N+1}^T + \xi(-\mu)_{M,N}^{(22)} f_{M+1/2,N-1/2}^T - \xi(-\mu)_{M,N}^{(21)} f_{M+1/2,N+1/2}^T - \eta(-\mu)_{M,N}^{(2)} f_{M+1,N}^T, \end{aligned} \quad (\text{A12})$$

where

$$\alpha(\mu)_{M,N} = 1 + \frac{\mu}{2} \sum_{n=-\infty}^{\infty} \left[\left(P_{M,N}^n P_{M+1/2,N}^n \right) \tilde{D}_{M+1/2,N}^n + \left(P_{M,N}^n P_{M-1/2,N}^n \right) \tilde{D}_{M-1/2,N}^n + \left(Q_{N+1/2}^n \right)^2 \tilde{D}_{M,N+1/2}^n + \left(Q_{N-1/2}^n \right)^2 \tilde{D}_{M,N-1/2}^n \right], \quad (\text{A13})$$

$$\alpha(\mu)_{M,N}^{(1)} \equiv \frac{\mu}{2} \sum_{n=-\infty}^{\infty} \left[\left(Q_{N+1/2}^n \right)^2 \tilde{D}_{M,N+1/2}^n \right], \quad (\text{A14})$$

$$\alpha(\mu)_{M,N}^{(2)} \equiv \frac{\mu}{2} \sum_{n=-\infty}^{\infty} \left[\left(Q_{N-1/2}^n \right)^2 \tilde{D}_{M,N-1/2}^n \right], \quad (\text{A15})$$

$$\xi(\mu)_{M,N}^{(11)} \equiv \frac{\mu}{2} \sum_{n=-\infty}^{\infty} \left[\left(P_{M,N}^n Q_N^n \right) \tilde{D}_{M-1/2,N}^n + \left(P_{M,N+1/2}^n Q_{N+1/2}^n \right) \tilde{D}_{M,N+1/2}^n \right], \quad (\text{A16})$$

$$\xi(\mu)_{M,N}^{(12)} \equiv \frac{\mu}{2} \sum_{n=-\infty}^{\infty} \left[\left(P_{M,N}^n Q_N^n \right) \tilde{D}_{M-1/2,N}^n + \left(P_{M,N-1/2}^n Q_{N-1/2}^n \right) \tilde{D}_{M,N-1/2}^n \right], \quad (\text{A17})$$

$$\xi(\mu)_{M,N}^{(21)} \equiv \frac{\mu}{2} \sum_{n=-\infty}^{\infty} \left[\left(P_{M,N}^n Q_N^n \right) \tilde{D}_{M+1/2,N}^n + \left(P_{M,N+1/2}^n Q_{N+1/2}^n \right) \tilde{D}_{M,N+1/2}^n \right], \quad (\text{A18})$$

$$\xi(\mu)_{M,N}^{(22)} \equiv \frac{\mu}{2} \sum_{n=-\infty}^{\infty} \left[\left(P_{M,N}^n Q_N^n \right) \tilde{D}_{M+1/2,N}^n + \left(P_{M,N-1/2}^n Q_{N-1/2}^n \right) \tilde{D}_{M,N-1/2}^n \right], \quad (\text{A19})$$

$$\eta(\mu)_{M,N}^{(1)} \equiv \frac{\mu}{2} \sum_{n=-\infty}^{\infty} \left[\left(P_{M,N}^n P_{M-1/2,N}^n \right) \tilde{D}_{M-1/2,N}^n \right], \quad (\text{A20})$$

and

$$\eta(\mu)_{M,N}^{(2)} \equiv \frac{\mu}{2} \sum_{n=-\infty}^{\infty} \left[\left(P_{M,N}^n P_{M+1/2,N}^n \right) \tilde{D}_{M+1/2,N}^n \right]. \quad (\text{A21})$$

All terms in both sides of Eq. (A12) with a constant v_{\perp} -index account for variations in the v_{\parallel} -space only. Therefore, they can be grouped into a single system of one-dimensional algebraic equations.

We transform all terms with v_{\perp} -index of M in both sides of Eq. (A12) into the tridiagonal matrices $[A(\mu)_M][F_M^{T+1}]$ and $[A(-\mu)_M][F_M^T]$, where $F_M^T = [f_{M,1}^T, f_{M,3/2}^T, f_{M,2}^T, \dots, f_{M,N_v}^T]_{1 \times 2N_v}^T$ (\mathbf{T} represents the transpose of a matrix), and

$$A(\mu)_M \equiv \begin{bmatrix} \alpha(\mu)_{M,1} & 0 & -\alpha(\mu)_{M,2}^{(1)} & 0 & 0 & \dots & 0 \\ 0 & \alpha(\mu)_{M,3/2} & 0 & -\alpha(\mu)_{M,5/2}^{(1)} & 0 & \dots & 0 \\ -\alpha(\mu)_{M,1}^{(2)} & 0 & \alpha(\mu)_{M,2} & 0 & -\alpha(\mu)_{M,3}^{(1)} & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & \dots & 0 & -\alpha(\mu)_{M,N_v-1}^{(2)} & 0 & \alpha(\mu)_{M,N_v} & 0 \\ 0 & \dots & 0 & 0 & -\alpha(\mu)_{M,N_v-1/2}^{(2)} & 0 & \alpha(\mu)_{M,N_v+1/2} \end{bmatrix}_{2N_v \times 2N_v}. \quad (\text{A22})$$

We transform all terms with v_{\perp} -index of $M-1/2$ in both sides of Eq. (A12) into the tridiagonal matrices $[B(\mu)_M^{(1)}][F_{M-1/2}^{T+1}]$ and $[B(-\mu)_M^{(1)}][F_{M-1/2}^T]$, where

$$B(\mu)_M^{(1)} \equiv \begin{bmatrix} 0 & \xi(\mu)_{M,3/2}^{(11)} & 0 & \dots & 0 \\ -\xi(\mu)_{M,1}^{(12)} & 0 & \xi(\mu)_{M,2}^{(11)} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & \dots & -\xi(\mu)_{M,N_v-1/2}^{(12)} & 0 & \xi(\mu)_{M,N_v+1/2}^{(11)} \\ 0 & \dots & 0 & -\xi(\mu)_{M,N_v}^{(12)} & 0 \end{bmatrix}_{2N_v \times 2N_v}. \quad (\text{A23})$$

We transform all terms with v_{\perp} -index of $M+1/2$ in both sides of Eq. (A12) into the tridiagonal matrices $[B(\mu)_M^{(2)}][F_{M+1/2}^{T+1}]$ and $[B(-\mu)_M^{(2)}][F_{M+1/2}^T]$, where

$$B(\mu)_M^{(2)} \equiv \begin{bmatrix} 0 & -\xi(\mu)_{M,3/2}^{(21)} & 0 & \cdots & 0 \\ \xi(\mu)_{M,1}^{(22)} & 0 & -\xi(\mu)_{M,2}^{(21)} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & \cdots & \xi(\mu)_{M,N_v-1/2}^{(22)} & 0 & -\xi(\mu)_{M,N_v+1/2}^{(21)} \\ 0 & \cdots & 0 & \xi(\mu)_{M,N_v}^{(22)} & 0 \end{bmatrix}_{2N_v \times 2N_v}. \quad (\text{A24})$$

We transform all terms with v_{\perp} -index of $M-1$ in both sides of Eq. (A12) into the tridiagonal matrices $[C(\mu)_M^{(1)}][F_{M-1}^{T+1}]$ and $[C(-\mu)_M^{(1)}][F_{M-1}^T]$, where

$$C(\mu)_M^{(1)} \equiv \begin{bmatrix} -\eta(\mu)_{M,1}^{(1)} & 0 & \cdots & 0 \\ 0 & -\eta(\mu)_{M,3/2}^{(1)} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & -\eta(\mu)_{M,N_v+1/2}^{(1)} \end{bmatrix}_{2N_v \times 2N_v}. \quad (\text{A25})$$

Lastly, we transform all terms with v_{\perp} -index of $M+1$ in both sides of Eq. (A12) into the tridiagonal matrices $[C(\mu)_M^{(2)}][F_{M+1}^{T+1}]$ and $[C(-\mu)_M^{(2)}][F_{M+1}^T]$, where

$$C(\mu)_M^{(2)} \equiv \begin{bmatrix} -\eta(\mu)_{M,1}^{(2)} & 0 & \cdots & 0 \\ 0 & -\eta(\mu)_{M,3/2}^{(2)} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & -\eta(\mu)_{M,N_v+1/2}^{(2)} \end{bmatrix}_{2N_v \times 2N_v}. \quad (\text{A26})$$

This strategy allows us to express Eq. (A12) as a single system of one-dimensional algebraic equations

$$\begin{aligned} & [C(\mu)_M^{(1)}][F_{M-1}^{T+1}] + [B(\mu)_M^{(1)}][F_{M-1/2}^{T+1}] + [A(\mu)_M][F_M^{T+1}] + [B(\mu)_M^{(2)}][F_{M+1/2}^{T+1}] + [C(\mu)_M^{(2)}][F_{M+1}^{T+1}] \\ & = [C(-\mu)_M^{(1)}][F_{M-1}^T] + [B(-\mu)_M^{(1)}][F_{M-1/2}^T] + [A(-\mu)_M][F_M^T] + [B(-\mu)_M^{(2)}][F_{M+1/2}^T] + [C(-\mu)_M^{(2)}][F_{M+1}^T]. \end{aligned} \quad (\text{A27})$$

Eq. (A27) only describes the VDF evolution in the v_{\perp} -space. However, each matrix term itself includes the VDF evolution in the v_{\parallel} -space. We transform Eq. (A27) into a single tridiagonal matrix

$$E(\mu)_{QLD} \begin{bmatrix} F_1^{T+1} \\ F_{3/2}^{T+1} \\ F_2^{T+1} \\ \vdots \\ F_{N_v+1/2}^{T+1} \end{bmatrix}_{(2N_v)^2 \times 1} = E(-\mu)_{QLD} \begin{bmatrix} F_1^T \\ F_{3/2}^T \\ F_2^T \\ \vdots \\ F_{N_v+1/2}^T \end{bmatrix}_{(2N_v)^2 \times 1}, \quad (\text{A28})$$

where

$$E(\mu)_{QLD} \equiv \begin{bmatrix} A(\mu)_1 & B(\mu)_1^{(2)} & C(\mu)_1^{(2)} & 0 & 0 & \cdots & 0 \\ B(\mu)_{3/2}^{(1)} & A(\mu)_{3/2} & B(\mu)_{3/2}^{(2)} & C(\mu)_{3/2}^{(2)} & 0 & \cdots & 0 \\ C(\mu)_2^{(1)} & B(\mu)_2^{(1)} & A(\mu)_2 & B(\mu)_2^{(2)} & C(\mu)_2^{(2)} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & \cdots & 0 & C(\mu)_{N_v}^{(1)} & B(\mu)_{N_v}^{(1)} & A(\mu)_{N_v} & B(\mu)_{N_v}^{(2)} \\ 0 & \cdots & 0 & 0 & C(\mu)_{N_v+1/2}^{(1)} & B(\mu)_{N_v+1/2}^{(1)} & A(\mu)_{N_v+1/2} \end{bmatrix}_{(2N_v)^2 \times (2N_v)^2}. \quad (\text{A29})$$

Eq. (A28) is in the form of a double matrix and $E(\mu)_{QLD}$ in Eq. (A29) defines the evolution matrix. The inner matrices of $E(\mu)_{QLD}$ evolve $f_{M,N}^T$ in the v_{\parallel} -space while the outer matrices of $E(\mu)_{QLD}$ evolve $f_{M,N}^T$ in the v_{\perp} -space during each time step. By multiplying Eq. (A28) with the inverse of $E(\mu)_{QLD}$ on both sides, Eq. (A28) provides the time evolution of $f_{M,N}^T$ in one time step simultaneously in the v_{\perp} - and v_{\parallel} -spaces. Therefore, it represents the numerical solution of Eq. (10) which describes the quasi-linear diffusion of a VDF through the all resonances.

B. NUMERICAL ANALYSIS OF FOKKER-PLANCK EQUATION

In this Appendix, we present our numerical strategy to solve the Fokker-Planck equation for Coulomb collisions in Eq. (40). Using the Crank-Nicolson scheme presented in Appendix A, we discretize Eq. (40) as

$$\begin{aligned}
& f_{M,N}^{T+1} - \sum_b \frac{\Gamma_{jb}}{2} \left[4\pi(\Delta v)^2 \mu \frac{m_j}{m_b} f_b(v_{\perp M}, v_{\parallel N}) f_{M,N}^{T+1} + \mu g_{M,N}^{\perp\perp} \left(f_{M+1/2,N+1/2}^{T+1} - f_{M-1/2,N+1/2}^{T+1} - f_{M+1/2,N-1/2}^{T+1} \right. \right. \\
& \quad \left. \left. + f_{M-1/2,N-1/2}^{T+1} \right) + \frac{\mu g_{M,N}^{\perp\parallel}}{2} \left(f_{M+1,N}^{T+1} - 2f_{M,N}^{T+1} + f_{M-1,N}^{T+1} \right) + \frac{\mu g_{M,N}^{\parallel\parallel}}{2} \left(f_{M,N+1}^{T+1} - 2f_{M,N}^{T+1} + f_{M,N-1}^{T+1} \right) \right. \\
& \quad \left. + (\Delta v) \mu h_{M,N}^{\perp\parallel} \left(f_{M+1/2,N}^{T+1} - f_{M-1/2,N}^{T+1} \right) + (\Delta v) \mu h_{M,N}^{\parallel\parallel} \left(f_{M,N+1/2}^{T+1} - f_{M,N-1/2}^{T+1} \right) \right] \\
& = f_{M,N}^T + \sum_b \frac{\Gamma_{jb}}{2} \left[4\pi(\Delta v)^2 \mu \frac{m_j}{m_b} f_b(v_{\perp M}, v_{\parallel N}) f_{M,N}^T + \mu g_{M,N}^{\perp\perp} \left(f_{M+1/2,N+1/2}^T - f_{M-1/2,N+1/2}^T - f_{M+1/2,N-1/2}^T \right. \right. \\
& \quad \left. \left. + f_{M-1/2,N-1/2}^T \right) + \frac{\mu g_{M,N}^{\perp\parallel}}{2} \left(f_{M+1,N}^T - 2f_{M,N}^T + f_{M-1,N}^T \right) + \frac{\mu g_{M,N}^{\parallel\parallel}}{2} \left(f_{M,N+1}^T - 2f_{M,N}^T + f_{M,N-1}^T \right) \right. \\
& \quad \left. + (\Delta v) \mu h_{M,N}^{\perp\parallel} \left(f_{M+1/2,N}^T - f_{M-1/2,N}^T \right) + (\Delta v) \mu h_{M,N}^{\parallel\parallel} \left(f_{M,N+1/2}^T - f_{M,N-1/2}^T \right) \right], \tag{B30}
\end{aligned}$$

where $g_{M,N}^{\perp\perp} \equiv \partial^2 g / \partial v_{\perp}^2$, $g_{M,N}^{\parallel\parallel} \equiv \partial^2 g / \partial v_{\parallel}^2$, $g_{M,N}^{\perp\parallel} \equiv \partial^2 g / \partial v_{\parallel} \partial v_{\perp}$, $h_{M,N}^{\perp\parallel} \equiv \partial h / \partial v_{\perp}$ and $h_{M,N}^{\parallel\parallel} \equiv \partial h / \partial v_{\parallel}$, estimated at $v_{\perp} = v_{\perp M}$ and $v_{\parallel} = v_{\parallel N}$.

Eq. (B30) represents a system of two-dimensional algebraic equations. Therefore, we transform Eq. (B30) into a single tridiagonal matrix using the same strategy for a double matrix as presented in Appendix A

$$E(\mu)_F \begin{bmatrix} F_1^{T+1} \\ F_{3/2}^{T+1} \\ F_2^{T+1} \\ \vdots \\ F_{N_v+1/2}^{T+1} \end{bmatrix}_{(2N_v)^2 \times 1} = E(-\mu)_F \begin{bmatrix} F_1^T \\ F_{3/2}^T \\ F_2^T \\ \vdots \\ F_{N_v+1/2}^T \end{bmatrix}_{(2N_v)^2 \times 1}, \tag{B31}$$

where

$$E(\mu)_F \equiv \begin{bmatrix} X(\mu)_1 & -Y(\mu)_1 & Z(\mu)_1 & 0 & 0 & \cdots & 0 \\ Y(\mu)_{3/2} & X(\mu)_{3/2} & -Y(\mu)_{3/2} & Z(\mu)_{3/2} & 0 & \cdots & 0 \\ Z(\mu)_2 & Y(\mu)_2 & X(\mu)_2 & -Y(\mu)_2 & Z(\mu)_2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & \cdots & 0 & Z(\mu)_{N_v} & Y(\mu)_{N_v} & X(\mu)_{N_v} & -Y(\mu)_{N_v} \\ 0 & \cdots & 0 & 0 & Z(\mu)_{N_v+1/2} & Y(\mu)_{N_v+1/2} & X(\mu)_{N_v+1/2} \end{bmatrix}_{(2N_v)^2 \times (2N_v)^2}, \tag{B32}$$

$$X(\mu)_M \equiv \begin{bmatrix} \varepsilon(\mu)_{M,1} & -\varepsilon(\mu)_{M,3/2}^{(2)} & -\varepsilon(\mu)_{M,2}^{(1)} & 0 & 0 & \cdots & 0 \\ \varepsilon(\mu)_{M,1}^{(2)} & \varepsilon(\mu)_{M,3/2} & -\varepsilon(\mu)_{M,2}^{(2)} & -\varepsilon(\mu)_{M,5/2}^{(1)} & 0 & \cdots & 0 \\ -\varepsilon(\mu)_{M,1}^{(1)} & \varepsilon(\mu)_{M,3/2}^{(2)} & \varepsilon(\mu)_{M,2} & -\varepsilon(\mu)_{M,5/2}^{(2)} & -\varepsilon(\mu)_{M,3}^{(1)} & \cdots & 0 \\ & \vdots & & \ddots & & \vdots & \\ 0 & \cdots & 0 & -\varepsilon(\mu)_{M,N_v-1}^{(1)} & \varepsilon(\mu)_{M,N_v-1/2}^{(2)} & \varepsilon(\mu)_{M,N_v} & -\varepsilon(\mu)_{M,N_v+1/2}^{(2)} \\ 0 & \cdots & 0 & 0 & -\varepsilon(\mu)_{M,N_v-1/2}^{(1)} & \varepsilon(\mu)_{M,N_v}^{(2)} & \varepsilon(\mu)_{M,N_v+1/2} \end{bmatrix}_{2N_v \times 2N_v}, \quad (\text{B33})$$

$$Y(\mu)_M \equiv \begin{bmatrix} \varrho(\mu)_{M,1}^{(2)} & \varrho(\mu)_{M,3/2}^{(1)} & 0 & 0 & \cdots & 0 \\ -\varrho(\mu)_{M,1}^{(1)} & \varrho(\mu)_{M,3/2}^{(2)} & \varrho(\mu)_{M,2}^{(1)} & 0 & \cdots & 0 \\ & \vdots & \ddots & & & \vdots \\ 0 & \cdots & 0 & -\varrho(\mu)_{M,N_v-1/2}^{(1)} & \varrho(\mu)_{M,N_v}^{(2)} & \varrho(\mu)_{M,N_v+1/2}^{(1)} \\ 0 & \cdots & 0 & 0 & -\varrho(\mu)_{M,N_v}^{(1)} & \varrho(\mu)_{M,N_v+1/2}^{(2)} \end{bmatrix}_{2N_v \times 2N_v}, \quad (\text{B34})$$

$$Z(\mu)_M \equiv \begin{bmatrix} -\tau(\mu)_{M,1} & 0 & \cdots & 0 \\ 0 & -\tau(\mu)_{M,3/2} & \cdots & 0 \\ \vdots & & \ddots & \vdots \\ 0 & \cdots & 0 & -\tau(\mu)_{M,N_v+1/2} \end{bmatrix}_{2N_v \times 2N_v}, \quad (\text{B35})$$

$$\varepsilon(\mu)_{M,N} \equiv 1 - \mu \sum_b \Gamma_{jb} \left[2\pi(\Delta v)^2 \frac{m_j}{m_b} f_b(v_{\perp M}, v_{\parallel N}) - \frac{g_{M,N}^{\perp\perp}}{2} - \frac{g_{M,N}^{\parallel\parallel}}{2} \right], \quad (\text{B36})$$

$$\varepsilon(\mu)_{M,N}^{(1)} \equiv \mu \sum_b \frac{\Gamma_{jb} g_{M,N}^{\parallel\parallel}}{4}, \quad (\text{B37})$$

$$\varepsilon(\mu)_{M,N}^{(2)} \equiv \mu \sum_b \frac{\Gamma_{jb} h_{M,N}^{\parallel}(\Delta v)}{2}, \quad (\text{B38})$$

$$\varrho(\mu)_{M,N}^{(1)} \equiv \mu \sum_b \frac{\Gamma_{jb} g_{M,N}^{\perp\perp}}{2}, \quad (\text{B39})$$

$$\varrho(\mu)_{M,N}^{(2)} \equiv \mu \sum_b \frac{\Gamma_{jb} h_{M,N}^{\perp}(\Delta v)}{2}, \quad (\text{B40})$$

and

$$\tau(\mu)_{M,N} \equiv \mu \sum_b \frac{\Gamma_{jb} g_{M,N}^{\perp\perp}}{4}. \quad (\text{B41})$$

Like Eq. (A28), Eq. (B31) provides the time evolution of $f_{M,N}^T$ in one time step simultaneously in the v_{\perp} - and v_{\parallel} -spaces. Therefore, it represents the numerical solution of Eq. (40) which describes the action of Coulomb collisions of particles in f_j with particles in f_b .

REFERENCES

- Albert, J. M. 2004, Using quasi-linear diffusion to model acceleration and loss from wave-particle interactions, doi: [10.1029/2004SW000069](https://doi.org/10.1029/2004SW000069)
- Brügmann, B., Tichy, W., & Jansen, N. 2004, Physical Review Letters, 92, doi: [10.1103/physrevlett.92.211101](https://doi.org/10.1103/physrevlett.92.211101)

- Feldman, W. C., Anderson, R. C., Bame, S. J., et al. 1983, *Journal of Geophysical Research: Space Physics*, 88, 9949, doi: [10.1029/JA088iA12p09949](https://doi.org/10.1029/JA088iA12p09949)
- Fitzenreiter, R. J., Ogilvie, K. W., Bale, S. D., & Viñas, A. F. 2003, *Journal of Geophysical Research: Space Physics*, 108, doi: [10.1029/2003JA009865](https://doi.org/10.1029/2003JA009865)
- Gary, S. P., Saito, S., & Li, H. 2008, Cascade of whistler turbulence: Particle-in-cell simulations, doi: [10.1029/2007GL032327](https://doi.org/10.1029/2007GL032327)
- Gendrin, R. 1968, Pitch angle diffusion of low energy protons due to gyroresonant interaction with hydromagnetic waves, doi: [10.1016/S0021-9169\(68\)91158-6](https://doi.org/10.1016/S0021-9169(68)91158-6)
- Gendrin, R. 1981, General relationships between wave amplification and particle diffusion in a magnetoplasma, doi: [10.1029/RG019i001p00171](https://doi.org/10.1029/RG019i001p00171)
- Gendrin, R., & Roux, A. 1980, Energization of helium ions by proton-induced hydromagnetic waves, doi: [10.1029/JA085iA09p04577](https://doi.org/10.1029/JA085iA09p04577)
- Gilat, A., & Subramaniam, V. V. 2011, Numerical methods : an introduction with applications using MATLAB
- Glauert, S. A., & Horne, R. B. 2005, Calculation of pitch angle and energy diffusion coefficients with the PADIE code, doi: [10.1029/2004JA010851](https://doi.org/10.1029/2004JA010851)
- Graham, G. A., Rae, I. J., Owen, C. J., et al. 2017, The evolution of solar wind strahl with heliospheric distance, doi: [10.1002/2016JA023656](https://doi.org/10.1002/2016JA023656)
- Gurgiolo, C., & Goldstein, M. L. 2016, Observations of diffusion in the electron halo and strahl, doi: [10.5194/angeo-34-1175-2016](https://doi.org/10.5194/angeo-34-1175-2016)
- Gurgiolo, C., Goldstein, M. L., Viñas, A. F., & Fazakerley, A. N. 2012, Direct observations of the formation of the solar wind halo from the strahl, doi: [10.5194/angeo-30-163-2012](https://doi.org/10.5194/angeo-30-163-2012)
- Gurnett, D. A., & Bhattacharjee, A. 2017, *Introduction to Plasma Physics: With Space, Laboratory and Astrophysical Applications*, Cambridge University Press, doi: [10.1017/9781139226059](https://doi.org/10.1017/9781139226059)
- Horaites, K., Boldyrev, S., & Medvedev, M. V. 2019, *MNRAS*, 484, 2474, doi: [10.1093/mnras/sty3504](https://doi.org/10.1093/mnras/sty3504)
- Horaites, K., Boldyrev, S., Wilson, Lynn B., I., Viñas, A. F., & Merka, J. 2018, *MNRAS*, 474, 115, doi: [10.1093/mnras/stx2555](https://doi.org/10.1093/mnras/stx2555)
- Isenberg, P., & Vasquez, B. 2011, A Kinetic Model of Solar Wind Generation by Oblique Ion-cyclotron Waves, doi: [10.1088/0004-637X/731/2/88](https://doi.org/10.1088/0004-637X/731/2/88)
- Isenberg, P. A., & Lee, M. A. 1996, A dispersive analysis of bispherical pickup ion distributions, doi: [10.1029/96JA00293](https://doi.org/10.1029/96JA00293)
- Iserles, A. 2008, *A First Course in the Numerical Analysis of Differential Equations*, Cambridge University Press, doi: [10.1017/CBO9780511995569](https://doi.org/10.1017/CBO9780511995569)
- Kennel, C. F., & Engelmann, F. 1966, *Velocity Space Diffusion from Weak Plasma Turbulence in a Magnetic Field*, doi: [10.1063/1.1761629](https://doi.org/10.1063/1.1761629)
- Khazanov, G. V., Gamayunov, K. V., Jordanova, V. K., & Krivorutsky, E. N. 2002, *Journal of Geophysical Research: Space Physics*, 107, SMP 14, doi: [10.1029/2001JA000180](https://doi.org/10.1029/2001JA000180)
- Klein, K. G., & Chandran, B. D. G. 2016, *ApJ*, 820, 47, doi: [10.3847/0004-637X/820/1/47](https://doi.org/10.3847/0004-637X/820/1/47)
- Lacombe, C., Alexandrova, O., Matteini, L., et al. 2014, Whistler Mode Waves and the Electron Heat Flux in the Solar Wind: Cluster Observations, doi: [10.1088/0004-637X/796/1/5](https://doi.org/10.1088/0004-637X/796/1/5)
- Lemons, D. S., & Feldman, W. C. 1983, *J. Geophys. Res.*, 88, 6881, doi: [10.1029/JA088iA09p06881](https://doi.org/10.1029/JA088iA09p06881)
- Ljepojevic, N. N., Burgess, A., & Moffatt, H. K. 1990, Calculation of the electron velocity distribution function in a plasma slab with large temperature and density gradients, doi: [10.1098/rspa.1990.0026](https://doi.org/10.1098/rspa.1990.0026)
- Lyons, L. R. 1974, *General relations for resonant particle diffusion in pitch angle and energy*, Cambridge University Press, doi: [10.1017/S0022377800024910](https://doi.org/10.1017/S0022377800024910)
- Lyons, L. R., Thorne, R. M., & Kennel, C. F. 1971, *Electron pitch-angle diffusion driven by oblique whistler-mode turbulence*, Cambridge University Press, doi: [10.1017/S0022377800006310](https://doi.org/10.1017/S0022377800006310)
- Lyutikov, M., & Gavriil, F. P. 2006, Resonant cyclotron scattering and Comptonization in neutron star magnetospheres, doi: [10.1111/j.1365-2966.2006.10140.x](https://doi.org/10.1111/j.1365-2966.2006.10140.x)
- Marsch, E. 2006, *Kinetic Physics of the Solar Corona and Solar Wind*, doi: [10.12942/lrsp-2006-1](https://doi.org/10.12942/lrsp-2006-1)
- Mourenas, D., Artemyev, A. V., Agapitov, O. V., Krasnoselskikh, V., & Mozer, F. S. 2015, *Journal of Geophysical Research: Space Physics*, 120, 3665, doi: [10.1002/2015JA021135](https://doi.org/10.1002/2015JA021135)
- Pagel, C., Gary, S. P., de Koning, C. A., Skoug, R. M., & Steinberg, J. T. 2007, Scattering of suprathermal electrons in the solar wind: ACE observations, doi: [10.1029/2006JA011967](https://doi.org/10.1029/2006JA011967)
- Pilipp, W. G., Miggenrieder, H., Montgomery, M. D., et al. 1987, Characteristics of electron velocity distribution functions in the solar wind derived from the Helios Plasma Experiment, doi: [10.1029/JA092iA02p01075](https://doi.org/10.1029/JA092iA02p01075)

- Roberg-Clark, G. T., Drake, J. F., Reynolds, C. S., & Swisdak, M. 2016, SUPPRESSION OF ELECTRON THERMAL CONDUCTION IN THE HIGH INTRACLUSTER MEDIUM OF GALAXY CLUSTERS, American Astronomical Society, doi: [10.3847/2041-8205/830/1/19](https://doi.org/10.3847/2041-8205/830/1/19)
- Rosenbluth, M. N., MacDonald, W. M., & Judd, D. L. 1957, Fokker-Planck Equation for an Inverse-Square Force, American Physical Society, doi: [10.1103/PhysRev.107.1](https://doi.org/10.1103/PhysRev.107.1)
- Ryu, C.-M., Rhee, T., Umeda, T., Yoon, P. H., & Omura, Y. 2007, Physics of Plasmas, 14, 100701, doi: [10.1063/1.2779282](https://doi.org/10.1063/1.2779282)
- Saito, S., Gary, S. P., Li, H., & Narita, Y. 2008, Whistler turbulence: Particle-in-cell simulations, doi: [10.1063/1.2997339](https://doi.org/10.1063/1.2997339)
- Saito, S., Gary, S. P., & Narita, Y. 2010, Wavenumber spectrum of whistler turbulence: Particle-in-cell simulation, doi: [10.1063/1.3526602](https://doi.org/10.1063/1.3526602)
- Saito, S., & Peter Gary, S. 2012, Beta dependence of electron heating in decaying whistler turbulence: Particle-in-cell simulations, doi: [10.1063/1.3676155](https://doi.org/10.1063/1.3676155)
- Seough, J., & Yoon, P. H. 2012, Quasilinear theory of anisotropy-beta relations for proton cyclotron and parallel firehose instabilities, doi: [10.1029/2012JA017645](https://doi.org/10.1029/2012JA017645)
- Stix, T. H. 1992, Waves in plasmas
- Štverák, Š., Maksimovic, M., Trávníček, P. M., et al. 2009, Radial evolution of nonthermal electron populations in the low-latitude solar wind: Helios, Cluster, and Ulysses Observations, doi: [10.1029/2008JA013883](https://doi.org/10.1029/2008JA013883)
- Summers, D. 2005, Quasi-linear diffusion coefficients for field-aligned electromagnetic waves with applications to the magnetosphere, doi: [10.1029/2005JA011159](https://doi.org/10.1029/2005JA011159)
- Summers, D., Thorne, R. M., & Xiao, F. 1998, Relativistic theory of wave-particle resonant diffusion with application to electron acceleration in the magnetosphere, doi: [10.1029/98JA01740](https://doi.org/10.1029/98JA01740)
- . 2001, Gyroresonant acceleration of electrons in the magnetosphere by superluminous electromagnetic waves, doi: [10.1029/2000JA000309](https://doi.org/10.1029/2000JA000309)
- Tang, B., Zank, G., & Kolobov, V. 2020, Numerical Modeling of Suprathermal Electron Transport in the Solar Wind: Effects of Whistler Turbulence, doi: [10.3847/1538-4357/ab7a93](https://doi.org/10.3847/1538-4357/ab7a93)
- Taran, S., Safari, H., & Daei, F. 2019, ApJ, 882, 157, doi: [10.3847/1538-4357/ab372b](https://doi.org/10.3847/1538-4357/ab372b)
- Ukhorskiy, A. Y., & Sitnov, M. I. 2014, Dynamics of Radiation Belt Particles, Boston, MA: Springer US, doi: [10.1007/978-1-4899-7433-4_17](https://doi.org/10.1007/978-1-4899-7433-4_17)
- Vasko, I. Y., Krasnoselskikh, V., Tong, Y., et al. 2019, Whistler Fan Instability Driven by Strahl Electrons in the Solar Wind, doi: [10.3847/2041-8213/ab01bd](https://doi.org/10.3847/2041-8213/ab01bd)
- Verscharen, D., & Chandran, B. D. G. 2018, NHDS: The New Hampshire Dispersion Relation Solver, American Astronomical Society, doi: [10.3847/2515-5172/aabfe3](https://doi.org/10.3847/2515-5172/aabfe3)
- Verscharen, D., Chandran, B. D. G., Jeong, S.-Y., et al. 2019, Self-induced Scattering of Strahl Electrons in the Solar Wind. <https://arxiv.org/abs/1906.02832>
- Vocks, C., & Mann, G. 2003, Generation of Suprathermal Electrons by Resonant Wave-Particle Interaction in the Solar Corona and Wind, doi: [10.1086/376682](https://doi.org/10.1086/376682)
- Vocks, C., Salem, C., Lin, R. P., & Mann, G. 2005, Electron Halo and Strahl Formation in the Solar Wind by Resonant Interaction with Whistler Waves, IOP Publishing, doi: [10.1086/430119](https://doi.org/10.1086/430119)
- Wilson, Lynn B., I., Chen, L.-J., Wang, S., et al. 2019, ApJS, 245, 24, doi: [10.3847/1538-4365/ab5445](https://doi.org/10.3847/1538-4365/ab5445)
- Yakimenko, V. 1963, ABSORPTION OF WAVES IN A PLASMA (QUASILINEAR APPROXIMATION)
- Yang, X., Wang, W., & Duan, Y. 2009, Journal of Computational and Applied Mathematics, 225, 31, doi: <https://doi.org/10.1016/j.cam.2008.06.013>
- Yoon, P. H. 2017, Kinetic instabilities in the solar wind driven by temperature anisotropies, doi: [10.1007/s41614-017-0006-1](https://doi.org/10.1007/s41614-017-0006-1)
- Yoon, P. H., López, R. A., Seough, J., & Sarfraz, M. 2017, Velocity moment-based quasilinear theory and particle-in-cell simulation of parallel electron firehose instability, doi: [10.1063/1.4997666](https://doi.org/10.1063/1.4997666)
- Yoon, P. H., & Seough, J. 2012, Quasilinear theory of anisotropy-beta relation for combined mirror and proton cyclotron instabilities, doi: [10.1029/2012JA017697](https://doi.org/10.1029/2012JA017697)
- Yoon, P. H., Seough, J., Hwang, J., & Nariyuki, Y. 2015, Macroscopic quasi-linear theory and particle-in-cell simulation of helium ion anisotropy instabilities, doi: [10.1002/2015JA021495](https://doi.org/10.1002/2015JA021495)
- Yoon, P. H., Ziebell, L. F., Gaelzer, R., Lin, R. P., & Wang, L. 2012, SSRv, 173, 459, doi: [10.1007/s11214-012-9867-3](https://doi.org/10.1007/s11214-012-9867-3)