

# Northumbria Research Link

Citation: Grandner, Michael A., Alfonso-Miller, Pamela, Fernandez-Mendoza, Julio, Shetty, Safal, Shenoy, Sundeep and Combs, Daniel (2016) Sleep: Important considerations for the prevention of cardiovascular disease. *Current Opinion in Cardiology*, 31 (5). pp. 551-565. ISSN 1531-7080

Published by: Lippincott, Williams & Wilkins

URL: <https://doi.org/10.1097/HCO.0000000000000324>  
<<https://doi.org/10.1097/HCO.0000000000000324>>

This version was downloaded from Northumbria Research Link:  
<http://nrl.northumbria.ac.uk/id/eprint/47090/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



Published in final edited form as:

*Curr Opin Cardiol.* 2016 September ; 31(5): 551–565. doi:10.1097/HCO.0000000000000324.

## Sleep: Important Considerations for the Prevention of Cardiovascular Disease

Michael A. Grandner, PhD MTR<sup>1,2,3,\*</sup>, Pamela Alfonso-Miller, MD<sup>1</sup>, Julio Fernandez-Mendoza, PhD<sup>4</sup>, Safal Shetty, MD<sup>3</sup>, Sundeep Shenoy, MD<sup>3</sup>, and Daniel Combs, MD<sup>3</sup>

<sup>1</sup>Sleep and Health Research Program, Department of Psychiatry, University of Arizona College of Medicine, Tucson, AZ

<sup>2</sup>Sarver Heart Center, University of Arizona College of Medicine, Tucson, AZ

<sup>3</sup>Department of Medicine, University of Arizona College of Medicine, Tucson, AZ

<sup>4</sup>Department of Psychiatry, Penn State Hershey Medical Center, Hershey, PA

### Abstract

**Purpose of Review**—Sleep plays many roles in maintenance of cardiovascular health. This review summarizes the literature across several areas of sleep and sleep disorders in relation to cardiometabolic disease risk factors.

**Current Findings**—Insufficient sleep duration is prevalent in the population and is associated with weight gain and obesity, inflammation, cardiovascular disease, diabetes, and mortality. Insomnia is also highly present and represents an important risk factor for cardiovascular disease, especially when accompanied by short sleep duration. Sleep apnea is a well-characterized risk factor for cardiometabolic disease and cardiovascular mortality. Other issues are relevant as well. For example, sleep disorders in pediatric populations may convey cardiovascular risks. Also, sleep may play an important role in cardiovascular health disparities.

**Summary**—Sleep and sleep disorders are implicated in cardiometabolic disease risk. This review addresses these and other issues, concluding with recommendations for research and clinical practice.

### Keywords

Sleep; cardiovascular; diabetes; insomnia; sleep apnea

## INTRODUCTION

Sleep is responsible for many regulatory and maintenance functions in human physiology. Although this has been well-established, the specific mechanisms of these functions, and the

\*CORRESPONDING AUTHOR: Michael A. Grandner PhD MTR, University of Arizona Department of Psychiatry, 1501 N Campbell Ave, Suite 8326, PO Box 245002, Tucson, AZ 85724-5002, (520) 626-6346, grandner@email.arizona.edu.

### CONFLICTS OF INTEREST

Dr. Grandner has received consulting fees from FitBit, Bayer, Nexalin, and the National Sleep Foundation, and grant support from Kemlin Industries. The remaining authors have no conflicts of interest.

relation of this physiology to relevant clinical outcomes, has only more recently received increased attention. With this in mind, much recent work has focused on the role of sleep in the maintenance of cardiometabolic health. This includes laboratory, clinical, and epidemiologic research, focusing on issues such as insufficient sleep and sleep disorders such as sleep apnea and insomnia. Also, research has focused on issues pertaining to special groups, such as pediatric populations and racial/ethnic minorities. This review summarizes the work in these areas and proposes some considerations for addressing sleep duration and sleep disorders in the context of preventing cardiovascular disease.

## HABITUAL SLEEP DURATION AND CARDIOVASCULAR DISEASE

Recently, the American Academy of Sleep Medicine and the Sleep Research Society jointly established a consensus panel to determine recommended guidelines for sleep duration. As a result of this process, a final guideline document was released(1-3) in addition to a companion manuscript that detailed methodological and other considerations(4, 5). In a parallel effort, the National Sleep Foundation developed its own guideline document(6) and companion manuscript(7). Both of these efforts agreed that a typical adult should get at least 7 hours of sleep on a typical night in order to maintain optimal health and functioning. (The document by the National Sleep Foundation also suggested an upper limit of 9 hours.) The American Thoracic Society also recently released a document that made a similar statement(8). Taken together, these documents identify 7 hours as the minimum amount of habitual sleep duration that is recommended for health and functioning. As a part of all of these efforts, the respective panels examined a growing body of literature that identifies associations between insufficient habitual sleep duration and cardiometabolic disease risk.

### Prevalence of insufficient sleep

Data from two survey cycles (2005-2006 and 2007-2008) of the National Health and Nutrition Examination Survey (NHANES) show that adults aged 20-39 and adults aged 40-59 reported sleeping less than 7 hours in a 24 hour period more frequently (37.0% and 40.3% respectively) than those aged over 60 years (32%)(9). These results are further supported by a recent CDC report, utilizing data from the Behavioral Risk Factor Surveillance System, that over 30% of US adults report less than 7 hours of habitual sleep duration(10).

Rather than being distributed evenly across the geographic landscape, Grandner and colleagues found unevenness in the distribution of such phenomenon. Using county-level data from the 2009 Behavioral Risk Factor Surveillance System (BRFSS), Grandner et al. identified insufficient sleep ‘hotspots’ in the Appalachian region that connects Tennessee, Virginia, West Virginia, Kentucky, and Ohio. Similar hotspots were found in the Midwest and southeast United States(11).

### Sleep duration, cardiovascular events and mortality

Sleep duration and mortality are associated in a U-shaped fashion, where lowest risk is usually found in those who sleep between 7-8 hours, and mortality risk increasing with further deviation on either end of the range. This pattern of findings has been explored in

multiple narrative reviews(12, 13) and meta-analyses(14, 15). Mechanisms for this relationship are still not completely delineated, but short sleep duration has been linked to chronic illness, cardiovascular disease (CVD), increased BMI and risk for obesity, accidents, hypertension, hypercholesterolemia heart attack and stroke(12, 16-26) [3-11]. To investigate the association between short and long sleep duration and total CVD and coronary heart disease incidence, Hoevenaar-Blom et al. conducted a 12-year prospective study of 20,432 healthy men and women in the Netherlands and found that those individuals who slept 6 hours had a 15% higher risk of CVD incidence and a 23% higher risk of CHD incidence compared to people who slept 7-8 hours. When sleep quality was compared, individuals with short sleep duration and poor subjective sleep has a 63% higher risk of CVD and 79% higher risk of CHD than those with normal sleep duration and good sleep quality(22). In another relevant study, Altman and colleagues examined data from the 2009 BRFFS and found that short sleep duration was associated with elevated prevalence of hypertension, hyperlipidemia and history of heart attack and stroke(16).

### **Sleep duration and obesity**

While the relationship between sleep duration and BMI can vary across age and age-related sleep need(19), insufficient sleep has been linked to lower energy expenditure, altered glucose homeostasis and alterations in hormones that mediate hunger(24, 25, 27-31). Altered sleep patterns may also be tied to lifestyle choices that are related to higher BMI and obesity(32). One possible mechanism for this could be explained by the metabolic hormones leptin and ghrelin. Leptin is secreted by adipose cells and it acts on the hypothalamus by regulating appetite through hunger inhibition. In laboratory sleep deprivation studies, decreased leptin secretion has been shown(26, 33). Ghrelin is a neuropeptide produced in the gastrointestinal tract. It acts on hypothalamic cells to increase hunger and has been shown to be increased in deprived sleep states in laboratory studies(26, 28, 33). This hormonal interaction may lead to an eating pattern that may lead to a dysregulation of energy balance through appetite stimulation that might predispose to cardiometabolic disease and weight gain. Along with the situational aspect of being awake longer leading to increased calorie intake, sleep deprived individuals are prone to indulge in late night snacks and make unhealthy food choices(34-40).

### **Sleep duration and behavioral risk factors**

A variety of behavioral risk factors have been positively linked to insufficient sleep, such as unemployment, increased alcohol consumption, self-reported poor mental health days, and active cigarette smoking(41, 42). Additionally, sleep deprivation leads to cognitive impairment and reduced attention(43-46). Chronic sleep insufficiency has played a role in motor vehicle accidents, industrial disasters, medical and other occupational errors, for instance(41, 42, 47-51). Others have found that short self-reported sleep duration is associated with risk of depression(4, 52-54), the same association has been found for long sleepers (>8-9 hours) (13).

### **Sleep duration and inflammation**

Inflammation is a well-established key mechanism in CVD risk and could be a possible mechanism linking sleep duration and CVD(55). Sleep deprivation has been associated with

increased inflammation and negative cardiovascular outcomes. Proinflammatory processes that promote the development of atherosclerotic plaques and Proinflammatory markers (such as tumor necrosis factor  $\alpha$  [TNF $\alpha$ ], interleukins 1 (IL-1), 6 (IL-6), and 17 (IL-17), C-reactive protein (CRP), cellular adhesion molecules, and visfatin) have demonstrated or suggested an association with sleep deprivation in laboratory studies(55). Cytokines (specifically IL-6) and elevated CRP are a clinically useful measure of systemic inflammations that have been shown to be present in healthy individuals exposed to partial and total sleep deprivation(56).

A recent meta-analysis did not find a consistent association between short sleep and inflammation(57). A study using the 2007-2008 NHANES(56) (not included in that meta-analysis) and a subsequent review(55) present some potential explanations. For example, using the 2007-2008 NHANES, levels of CRP were distributed in a U-shape across the US population. A few elements of this relationship suggest reasons why other studies could not find this relationship. First, these elevations were only seen at the extreme ends of the sleep duration spectrum (<5 and >9 hours), which most studies are not powered to examine. Second, there were significant race/ethnicity and sex patterns, suggesting that the makeup of the sample also played a role in findings. Third, since CRP does not demonstrate a circadian rhythm(58), a single measurement (common in epidemiologic studies) is more appropriate for this analyte. Finally, the NHANES sample may be one of the few samples that is sufficiently large and diverse (regarding age, sex, race/ethnicity, and geography) to detect the elevation associated with short sleep.

### **Sleep duration and Diabetes Mellitus (DM) risk**

Sleep deprivation alters glucose homeostasis, leading to insulin resistance and risk of diabetes(17, 18, 24, 59-61). Sleep extension in restricted sleepers may lessen this relationship by improving glucose tolerance(17). Current research is exploring mechanistic pathways involved, including adipose tissue physiology and the role of circadian genetics.

In the Western New York Health Study, Rafalson and colleagues found that sleep duration <6 h was associated with impaired fasting glucose [30]. In another study, Shankar and colleagues found that self-reported insufficient sleep was positively associated with prevalence of Diabetes(62). In a cross sectional study, Hall et al. found that both short (>7h) and long sleepers (>8) were associated with metabolic syndrome in midlife adults despite adjusting for traditional risk factor components(63).

Several reviews on this topic have been published elsewhere(24, 60, 64, 65), including several meta-analyses(3, 66-69), which show that habitual short sleep duration was associated with a 30-50% increased incidence of diabetes.

### **Sleep duration and blood pressure**

In a study of ambulatory blood pressure in 18 healthy subjects, Lusardi and colleagues observed that when sleep was restricted to the second half of the night for only a week, elevated diastolic and systolic blood pressure were present, compared to the week prior where a typical night of sleep was observed with no alterations(70).

At the population level, Altman and colleagues found increased risk of hypertension associated with both short ( 5-6 hours)(16). In an analysis of the 2004–2005 NHIS data, Buxton and Marcelli found that short sleep duration is associated with increased prevalence of hypertension and CVD(21). Using the 2007-2008 NHANES data, Grandner and colleagues established an association between short sleepers and elevated risk for self-reported history of hypertension, but not recorded blood pressure(20).

### **Sleep duration and lipids**

Increased CVD morbidity risk is tied to a decrease in High-Density Lipoprotein (HDL) and an increase in Low-Density lipoprotein (LDL). This relationship might be mediated by age and gender, as lipoprotein metabolism is strongly associated with sex hormones. In one study, elevated triglyceride and LDL profiles were found in Japanese women who slept <6 hours and >8 hours and were lowest among women who slept 6-7 hours. Risk of high LDL was low in men sleeping 8 hours(71). Short sleep duration has also been identified as a risk factor for hyperlipidemia and hypercholesterolemia in adolescents(72). Although a different study found relationships to hyperlipidemia only in very short sleepers, this association was most evident in non-Hispanic whites and Asians/others(20). It should be noted that in this study population, women were less likely to report as short or long sleepers. Very short sleep (>5) and increased incidence of hyperlipidemia was also found by Altman et al(16).

Taken together, habitual short sleep duration is highly prevalent in the US population. Regarding cardiometabolic risk, short sleep duration (6 hours or less), and especially very short sleep duration (less than 5 hours) is associated with increased cardiometabolic morbidity and mortality, including weight gain and obesity, hypertension, hyperlipidemia, and diabetes.

## **SLEEP APNEA AND CARDIOVASCULAR DISEASE**

### **Prevalence of Sleep Apnea**

In a recent review of 11 published epidemiological studies, the prevalence of obstructive sleep apnea (OSA), defined as an Apnea-Hypopnea Index (AHI) greater or equal to 5, was estimated to be 22% in men and 17% in women(73). More recent studies indicate that this is increasing, with an estimated prevalence of 37% in men and 50% in women(74, 75). The reason for the rising prevalence is thought to be secondary to differences in diagnostic equipment, the definition of OSA, study design, and patient characteristics such as age and obesity.(73) The prevalence of OSA in obesity is about 50% (76), with much higher prevalence in morbidly obese individuals.(77)

Drager and colleagues, studied 152 consecutive patients with metabolic syndrome (MetS) with a standard polysomnogram (PSG) and found the prevalence of moderate-severe sleep apnea, defined as an AHI greater than or equal to 15, to be about 60%. Patients with MetS and OSA were found to have greater cardiometabolic morbidity in comparison with those without OSA.(78)

## Obstructive sleep apnea (OSA) and cardiovascular mortality

The association between OSA and cardiovascular mortality is complicated due to associated factors such as obesity, insulin resistance, and dyslipidemia. However, OSA is an independent risk factor for atherosclerotic heart disease(79). Epidemiological data supports the idea that OSA can initiate not only cardiovascular conditions such as hypertension and ischemic heart disease but also worsen existing disease(80).

The key pathology resulting in the adverse cardiovascular outcomes stems from the intermittent hypoxia, hypercapnia, intrathoracic pressure changes and the resulting autonomic changes(80). The intermittent hypoxia induces oxygen free radical production and sets off an inflammatory response, over time causing adverse cardiovascular damage(81, 82).

A prospective study from Punjabi and colleagues, showed increased all-cause mortality, specifically from coronary artery disease, in men aged 40-70 years with severe OSA characterized by apnea-hypopnea index (AHI) of greater than 30(83). Similar results were noted in an observational study where severe OSA was linked to an increased risk of fatal cardiovascular events (odds ratio of 2.87; 1.17 – 7.51) as well as non-fatal cardiovascular events (odds ratio of 3.17, 1.12 – 7.51) compared to a control of healthy individuals(84). The intermittent hypoxia-hypercapnia results in myocardial oxygen demand-supply mismatch and can trigger nocturnal angina as well as ischemic electrocardiographic (EKG) changes(85).

OSA is a common treatable cause of drug-resistant hypertension(86). Independent of confounders, the odds of developing hypertension was 2.89 (95% CI; 1.46 – 5.64) for patients with an AHI of >15 compared to the control group(87). A higher prevalence of arrhythmias such as paroxysmal supraventricular tachycardia and atrial fibrillation is noted in patients with OSA. An observational study reported patients with untreated OSA had twice the risk of recurrence of atrial fibrillation compared to the group who were treated with Continuous Positive Airway Pressure (CPAP) therapy(88).

Patients with congestive heart failure can have a component of central sleep apnea atop their obstructive component(89). Mansfield and colleagues, in a randomized trial, showed that patients treated with CPAP in addition to standard heart failure therapy had improved ejection fraction as well as improved quality of life(90). Similar results were noted by Kaneko et al., who noticed an improvement in ejection fraction as well morning blood pressures with CPAP treatment. This improvement was observed as early as one month from the initiation of CPAP therapy(91). Promising results were also noted by Ryan and colleagues who reported a 58% reduction in the frequency of ventricular arrhythmias during sleep in patients with OSA and heart failure(92).

A prospective observational study involving patients with CAD and moderate OSA with AHI >15 showed patients on CPAP had a significantly decreased cardiovascular mortality, acute coronary syndrome, hospitalization from heart failure exacerbation as well as need for coronary revascularization compared to the group which did not use the CPAP(93). Hence, a

consistent finding is that there is an improved cardiovascular outcome with regular use of CPAP in patients with OSA.

### **Syndrome Z and Metabolic Syndrome**

Syndrome Z describes the association between OSA and metabolic syndrome (MetS). The odds for the presence of MetS in OSA ranges from five to nine fold when compared to individuals without OSA.(94) OSA may be an independent risk factor for metabolic syndrome.(95)

### **Sleep Apnea and Heart Failure**

The prevalence of OSA among heart failure patients is about 30-50% while that of central sleep apnea (CSA) is about 25-40%. The presence of Congestive heart failure (CHF) may exacerbate or contribute to the development of OSA(96). Nocturnal fluid shift rostrally from the lower extremity towards the neck is believed to contribute to the pathogenesis of both OSA and CSA.(97)

Episodes of apnea and hypopnea related to pharyngeal collapse associated with OSA leads to negative inspiratory intrathoracic pressure. As a consequence, the venous return to heart increases increasing the right ventricular (RV) preload. Intermittent hypoxia, causes pulmonary vasoconstriction and hence increases RV after load. This may lead to the development of pulmonary hypertension and RV dysfunction. The post apneic related increase in sympathetic drive can persist during the day and lead to neurohormonal dysregulation and can adversely affect cardiac remodeling(96). Patients with HF may not present with typical symptoms of sleep apnea, such as excessive daytime sleepiness due to the heightened sympathetic activity(98). Intermittent hypoxia results in oxidative stress and release of inflammatory mediators involved in the progression of atherosclerosis. Wang and colleagues, in a study of patients with HF to determine the effect of untreated moderate to severe OSA versus mild to no OSA, found that the mortality was significantly higher in the untreated moderate to severe OSA group(99). Treatment with CPAP therapy has been shown to improve left ventricular ejection fraction, functional class, reduce hospitalization and consequently survival.(96)

Patients with HF have a high prevalence of Central sleep apnea (CSA). Cheyne stokes respiration (CSR), a pattern of breathing characterized by a cyclic crescendo decrescendo pattern of hyperventilation followed by apnea in the absence of upper airway obstruction(100). CSR in low output HF is thought to be due to an impaired feedback mechanism with delayed messaging to the respiratory control center resulting in respiratory oscillation in response to a fluctuating PCO<sub>2</sub> level(101). The presence of wake hypocapnia, as demonstrated on an arterial blood gas may help predict the presence of CSR during sleep. (101) Treatment consists of targeted optimization of medical therapy to HF status. CPAP improves CSR by multiple mechanisms, such as by increasing lung volume, stabilization of the upper airway, and improved oxygenation and cardiac function.(102)



### Sleep Apnea and Diabetes Risk

The prevalence of OSA in patients with history of diabetes mellitus (DM) ranges from 58-87%.(94) OSA has been linked with increased risk of developing DM due to deregulation of glucose metabolism and alteration in the hypothalamic –pituitary adrenal axis (HPA) axis(103, 104). Intermittent hypoxia and sleep fragmentation related to OSA have been shown to independently increase insulin resistance, increase sympathetic activity and serum cortisol, and impair glucose clearance(94, 105) . The severity of OSA has a direct correlation with the worsening of glucose metabolism. In a study done to assess the effect of the severity of OSA and glycemic control by measurement of Hemoglobin A1c, it was found that HbA1c increased by 1.49% in mild OSA, 1.93% in moderate OSA, and 3.69% in severe OSA, compared with individuals without OSA.(106)

Improvement in glycemic control has been demonstrated after 3 months of CPAP use, with improvement related to number of hours of nightly CPAP use(107, 108). In a recent randomized control trial, patients who received CPAP when compared to the control group who did not receive CPAP were found to have improved glucose control and insulin sensitivity.(109) Also, CPAP therapy has been shown to improve insulin sensitivity in patients with OSA with or without diabetes(110, 111).

### Sleep Apnea and Hypertension

Epidemiologic studies support the strong relationship between OSA and hypertension.(112) The prevalence of OSA in individuals with hypertension is estimated to be about 40-50%(112) and approaches 83% in patients with resistant hypertension(86). In a large prospective longitudinal study on participants of the Wisconsin sleep cohort study, the odds for the presence of hypertension were 2.03 for apnea-hypopnea index (AHI) between 5.0 to 14.9 events per hour and 2.89 for AHI greater than 15 events per hour(113). Epidemiologic studies suggest the relationship between hypertension and OSA to be bidirectional. Activation of the sympathetic nervous system and renin-angiotension-aldosterone system, along with oxidative stress and endothelial dysfunction, implicate OSA as an independent risk factor for the development of hypertension. Inhibition of upper airway muscles, volume overload and rostral shift of fluid nocturnally can contribute to pharyngeal edema and the development of OSA in hypertensive patients.(114)

Schein and colleagues performed a recent met analysis and systematic review of 16 RCTs involving over 1000 patients, to study the effects of CPAP therapy on hypertension in OSA patients. Modest reduction in mean systolic blood pressure (SBP) (–4.92 mm Hg; 95%CI –8.70 to –1.14) and mean (–2.56 mm Hg; 95% CI –4.43 to –0.68 mmHg) nighttime blood pressure were seen. No significant change in daytime SBP (–0.74 mm Hg; 95% CI –3.9 to 2.41) and daytime diastolic blood pressure (–1.86 mm Hg; 95% CI –4.55 to 0.83) were observed, however.(115)

The effect of CPAP therapy is more pronounced in patients with resistant hypertension. In a recent meta-analysis, to study the effect of CPAP therapy on OSA patients on this subset of patients, showed the mean improvement of ambulatory SBP and DBP of –6.74 mm Hg (95% CI –9.98 to –3.49;P value <0.001) and –5.94 mm Hg (95% CI;–9.40 to

-2.47;P=0.001) respectively(116). The improvement correlated with the duration of CPAP use. Blood Pressure improved by 1.3 mm Hg for each additional hour of CPAP use.(117)

### **Sleep Apnea and Dyslipidemia**

Murine experiments reveal that it is the intermittent hypoxia which upregulates triglyceride and phospholipid synthesis and inhibits cholesterol uptake in the liver resulting in dyslipidemia(118, 119). OSA has been associated with dyslipidemia and its associated higher CV mortality. However, the causal relation between OSA and dyslipidemia remains to be elucidated due confounders such as obesity, diet, and physical activity.

The evidence is limited to cross-sectional studies and a few randomized studies. The major limitation of all these studies being their duration of follow-up. The majority of the studies indicate that the total cholesterol and LDL were unchanged. However, an increase in triglyceride levels and decrease in the HDL levels were noted in patients with OSA(120, 121). Non-randomized trials studying the effects of treatment of OSA with CPAP revealed a decrease in total cholesterol, LDL, homocysteine, and triglycerides, while on treatment; however, there was mixed data on HDL(122-127). Robinson and colleagues demonstrated a decrease in total cholesterol while on treatment with CPAP(128). Similarly, Phillips and colleagues, in a randomized, placebo-controlled trial looking at the role of CPAP on postprandial lipid levels in patients with OSA, concluded that treatment of severe OSA with CPAP improved postprandial triglyceride levels as well as total cholesterol levels(129). These trials, although randomized, have limitations such as a lack of adjustment for CPAP adherence and the role of lipid-lowering medications. An observational study in Portugal followed patients for an eight year period and concluded that CPAP treatment improves daytime sleepiness in patients with OSA but does not play as significant a role in improving their lipid profile comparable to the lipid-lowering medications(130). Although a direct causal relation of OSA and dyslipidemia is yet to be determined, there is increasing evidence that an association exists(131). The actual impact of the treatment of OSA in the improvement of the lipid profile is debatable and will require large RCTs.

### **INSOMNIA AS A CARDIOVASCULAR DISEASE RISK FACTOR**

Insomnia has long been viewed as a symptom of the “worried well.” However, accumulating evidence indicates that insomnia disorder is a significant risk factor for CVD(132, 133).

Insomnia, especially when coupled with short sleep duration, is associated with hyperactivity of the hypothalamic-pituitary-adrenal axis (e.g., increased cortisol secretion) and sympathetic system (e.g., impaired heart rate variability) and, consequently, with a clinically significant risk of CVD (e.g., hypertension, type-2 diabetes, myocardial infarction, and coronary heart disease).

#### **Epidemiology of Insomnia: Scope of the Problem**

The prevalence of insomnia in the general population ranges between 8-40%, depending on the definition used. While 20-30% of the general population has “poor sleep” (i.e., difficulty falling asleep, staying asleep, early morning awakening or non-restorative sleep at any given time), another 8-10% of the population has insomnia(134, 135).

Natural history studies indeed show that insomnia is a highly persistent condition, whereas the course of “poor sleep” is more unstable and likely to remit(136-140), which indicates that insomnia is a disorder while “poor sleep” is a symptom of underlying physical or mental health problems(137, 138, 140). Despite the fact that the objective sleep of people with insomnia is different than that of normal sleepers(135, 141) and that objective short sleep duration has been shown to predict the development of “poor sleep” into insomnia(138) and the persistence of insomnia(140), polysomnographic (PSG) studies are not required for the diagnosis of insomnia(141), which contrasts with the current indication of PSG for the evaluation of patients with sleep disordered breathing and their associated CVD risk. Actually, the associations between insomnia and significant CVD risk has not been systematically examined until recently.

### **Insomnia as a Behavioral Risk Factor**

Insomnia may be associated with increased CVD risk through its association with poor health behaviors. Indeed, individuals with “poor sleep” or insomnia are more likely to report increased alcohol and caffeine use, smoking, and lack of physical activity and to show lower cardiorespiratory fitness as compared to normal sleepers(142). Moreover, individuals with insomnia may be at increased CVD risk through poor diet, as subjective and objective sleep has been associated with dietary macronutrient and micronutrient composition(143). For example, a recent study of the 2007-2008 National Health and Nutrition Examination Survey (NHANES) found inadequate intake of alpha-carotene, calcium, selenium, salt, carbohydrates, vitamin D, lycopene, dodecanoic, hexadecanoic, butanoic or hexanoic acids to be associated with “poor sleep”(144). It is likely that the relationship between insomnia and “poor sleep” with many of these health behaviors is bidirectional(137, 138, 140). Interestingly, however, sleep hygiene therapy alone, which targets poor health behaviors, is not effective in the treatment of insomnia. Importantly, most studies on the independent association of insomnia with CVD risk have controlled for alcohol and smoking but not diet or physical activity. More work is needed to establish the relative contribution of health behaviors to the CVD risk associated with insomnia.

### **Insomnia Pathophysiology and Cardiovascular Risk**

Current models of insomnia posit the joint effect of stressful life events(145) and stress-related psychological factors as central to the pathogenesis of insomnia(146, 147); thus, activation of the stress system, i.e., hypothalamic-pituitary-adrenal (HPA) axis and sympathetic nervous system, should be a key pathophysiologic mechanism of the persistence of insomnia. While early studies reported no difference between “poor sleepers” and controls in cortisol levels, later studies found that 24-h urinary free cortisol, norepinephrine, and catecholamine metabolites levels were increased in patients with insomnia or were correlated with PSG indices of sleep disturbance in insomnia patients(148-151). Also, other studies, with a few exceptions(152-155), have found that insomnia is associated with increased nocturnal heart rate, blunted heart rate variability (HRV)(156-160), altered sympathovagal balance, as measured by impedance cardiography(156, 161), increased overall oxygen consumption (VO<sub>2</sub>), a measure of increased whole-body metabolic rate(162, 163), increased pupil size, indicative of sympathetic system activation(164), increased central nervous system activation during sleep(165-167), increased or altered systemic

inflammation(57, 168), and increased daytime sleep latencies in the Multiple Sleep Latency Test (MSLT), a test of physiologic sleep propensity(155, 169). The shorter the objective sleep duration, the longer the sleep latencies in the MSLT in persons with insomnia(170, 171).

In summary, increased peripheral and central markers of HPA axis dysfunction and sympathetic activation are primarily found in insomniacs with objective short sleep duration, who suffer from a disorder of 24-hour hyperarousal(172). Thus, proposed physiologic mechanisms of the association of insomnia with CVD risk include neuroendocrine dysregulation, autonomic imbalance, and increased inflammation.

### **Insomnia and Cardiovascular Disease Risk**

Although the association of insomnia with cardiovascular disease has been reported for several years(173), its association with clinical risk factors (e.g., hypertension, diabetes, metabolic syndrome) and cardiovascular disease has remained largely unexplored until recently(174). In fact, only in the past few years several systematic reviews and meta-analyses have been published on the association of insomnia with CVD risk. For example, the estimated risk for incident hypertension associated with “poor sleep” or insomnia ranges between 5% and 20%(175), that for type-2 diabetes (T2D) about 57%(3), and that for incident myocardial infarction (MI) and coronary heart disease (CHD) between 41% and 55%, respectively(176, 177). Although these studies have shown a significant association between insomnia, mainly “poor sleep”, and CVD risk, the effect sizes reported were modest and most studies did not include PSG and could not control for SDB.

Based on the physiologic mechanisms reviewed above, recent work from the Penn State Adult Cohort has shown a synergistic effect between insomnia and PSG-measured short sleep duration (i.e., < 6 hours) on the risk for hypertension and T2D even after controlling for multiple confounders, including SDB(178-180). Specifically, the odds of T2D in individuals with insomnia who slept < 6 hours were 2.1- to 3.0-fold cross-sectionally and the odds of hypertension were 3.5- to 5.1-fold cross-sectionally<sup>71</sup> and 3.8-fold longitudinally, while the odds of hypertension or T2D in individuals who reported insomnia but slept objectively > 6 hours were not significantly increased (e.g., OR = 1.3 and 0.85). Importantly, these findings have been replicated in two recent studies in patients with insomnia using PSG-measured short sleep duration(181) and MSLT-measured increased sleep latency(182).

While SDB is strongly associated with the metabolic syndrome (MetS), the evidence for insomnia is limited and inconsistent(63, 183, 184). A potential explanation may be related to different physiologic mechanisms in SDB and insomnia; for example, central obesity is strongly associated with incident SDB and “poor sleep”, but not as much with incident insomnia. In fact, patients with insomnia are typically non-obese(185) and in a recent longitudinal study(186), despite sleeping objectively shorter than controls or “poor sleepers”, non-obese insomniacs were less likely to become obese than controls or “poor sleepers”, a finding consistent with the presence of increased whole-body metabolic rate(162). These data indicate that insomnia with short sleep duration may be linked to insulin resistance and CVD independent of central obesity.

Current literature indicates that insomnia is a premorbid risk factor of CVD. Accumulating evidence suggests that the association of insomnia with CVD risk is more pronounced when insomnia is associated with objective short sleep duration or other measures of physiologic arousal. Evidence suggests that measures of insomnia, and especially objective sleep measures, should be included in the estimation of CVD risk in the clinical and general population.

## **SLEEP AND CARDIOVASCULAR DISEASE RISK IN PEDIATRICS**

Factors that increase cardiovascular risk have been shown to emerge early in life. For example, blood pressure and body mass index in adolescence predict future adult cardiovascular risk (187). Similar to what has been shown in adults; sleep-related factors are important contributors to cardiovascular risk in childhood. These factors include both sleep duration, as well as factors that disrupt sleep quality, particularly sleep-disordered breathing.

### **Sleep duration and cardiovascular disease risk in children**

Sleep curtailment and short sleep duration have been linked to increased adiposity and elevated BMI in longitudinal studies spanning from infancy to adolescence (188, 189). Additionally, there is robust evidence linking short sleep duration to impaired glucose metabolism. A recent study showed that shorter sleep duration, as well as worse sleep efficiency, was associated with impaired glucose tolerance (190). This is consistent with most prior literature, which generally has shown that short sleep in children and adolescents is associated with impaired glucose metabolism (191). One prior study identified a U-shaped relationship, with short and long sleep duration both predicting impaired glucose tolerance, similar to adults (192). A recent cross-sectional study has also identified that short sleep duration in obese children is associated with elevated homocysteine levels compared to obese children who obtain sufficient sleep (193).

### **Pediatric sleep-disordered breathing and cardiovascular disease risk**

SDB is a prevalent condition in children, with obstructive sleep apnea seen in 1.2-5.7% of all children (194). In children with cardiovascular risk factors, the prevalence may be even higher. A recent retrospective study found that 55% of children with hypertension referred for polysomnography were found to have sleep-disordered breathing (195). Sleep-disordered breathing in children is associated with adverse cardiovascular effects including dysregulation of blood pressure (196, 197) and cardiac dysfunction (198, 199). Children with sleep-disordered breathing have significantly elevated pulmonary arterial pressures as well as right ventricular hypertrophy and dysfunction (199). There is also evidence for left ventricular dysfunction (198). The underlying pathophysiologic effects of sleep-disordered breathing that may cause these deleterious effects include endothelial dysfunction (200), autonomic dysfunction (201), and chronic inflammation (202). Importantly, these adverse cardiovascular effects appear to be reversible with treatment (199). Sleep disordered breathing is also associated with significant metabolic adverse effects. The presence of OSA has been shown to be associated with increased serum triglycerides, blood glucose and insulin resistance (203).

In general, recent evidence in children has continued to identify sleep as an important factor in cardiovascular risk. Both short sleep duration and sleep-disordered breathing appear to be important modifiable risk factors that contribute towards overall cardiovascular risk. Targeted sleep interventions during childhood may be an important strategy to lower cardiovascular risk throughout the lifetime.

## **SLEEP AND CARDIOVASCULAR HEALTH DISPARITIES**

### **Cardiovascular Health Disparities as a Public Health Priority**

Health disparities exist when one group of people is at a disadvantage as a virtue of their socioedemographic category and not inherent physiologic risk. In these cases, increased risk for morbidity and mortality are a product of social-environmental influences and as such represent a societal concern. Although health care advances have proceeded at a dramatic pace over the last half-century, racial/ethnic and socioeconomic health disparities are still well-documented in the United States.(204-206). In particular many racial/ethnic minority groups (particularly Blacks/African-Americans and Hispanics-Latinos) are at greater risk of obesity, heart disease, diabetes, and cardiovascular events (204, 206, 207). These effects persist even after accounting for socioeconomics (204, 206). It is likely that a complex combination of structural, physiological, psychological and behavioral differences drive these relationships (205, 206, 208-210). In addition, environmental factors such as neighborhood may play a significant role through differential exposures to social, psychological, and physical influences (211, 212). Understanding and reducing health disparities has been identified as a key public health priority(206, 213).

### **Sleep Disparities in the Population**

Several studies have documented sleep disparities in the US population. A meta-analysis by Ruitter and colleagues(214) showed that Blacks/African-Americans obtained less polysomnographic sleep than Non-Hispanic Whites. Blacks/African-Americans also obtained less Slow Wave Sleep. Other studies have also found that Blacks/African-Americans obtained less Slow Wave Sleep(215-219) and had poorer sleep efficiency(217, 219) than Non-Hispanic Whites.

Several epidemiologic studies have examined population-level sleep disparities. Using the 2007-2008 NHANES, Whinnery and colleagues(220) found that relative to non-Hispanic Whites, Blacks/African-Americans were 3.5 times as likely to report very short (<5 hours) sleep duration and twice as likely to report short (5-6 hours) sleep duration. In this same sample, non-Mexican Hispanics/Latinos were 87% more likely to report very short sleep and 35% more likely to report short sleep. And Asians/Others were 2.5 times as likely to report very short sleep and 67% more likely to report short sleep. These findings are mirrored in many other studies, which have shown increased prevalence of habitual short sleep duration among racial/ethnic minority groups(221-225).

Regarding sleep disorders, Ruitter and colleagues(226) found that African-Americans were more likely to have sleep disordered breathing, of greater severity, compared to Whites. Data from the 2007-2008 NHANES(227) also showed that Blacks/African-Americans were more

likely to report sleep latency >30 minutes, but self-reported difficulties initiating or maintaining sleep were all reported less frequently among minority groups, suggesting that despite differences in sleep, these are not perceived as “difficulties.” In this sample, Non-Mexican Hispanics/Latinos were more likely to report choking/gasping during sleep and snoring(228). Other studies have found inconsistent relationships between race/ethnicity and sleep symptoms. Although some studies have found that sleep quality is worse among racial/ethnic minorities(229-232), others have not(233-235). A potential explanation is the variable, interactive roles of race/ethnicity, socioeconomic, and cultural factors.

### **Could Sleep be a Modifiable Risk Factor for Cardiovascular Health Disparities?**

If racial/ethnic minorities are at increased risk of cardiometabolic disease, and if insufficient sleep is also associated with cardiometabolic disease, and if racial/ethnic minorities are more likely to experience insufficient sleep, it is plausible to suggest that sleep may play a role in disparities and therefore may serve as an intervention target. Although this has not yet been directly tested, several studies have shown that the relationship between sleep and cardiometabolic disease risk does depend on race/ethnicity. Knutson and colleagues(236) found that racial differences in 5-year blood pressure change were explained by differences in sleep duration. Also, Grandner and colleagues(56) found that the relationship between sleep duration and C-reactive protein (a cardiovascular risk marker) differed by race/ethnicity. This was followed up by another analysis that showed that the relationship between sleep duration and obesity, diabetes, hypertension and hyperlipidemia depended on self-identified race/ethnicity(20).

## **PREVENTING CARDIOVASCULAR DISEASE: A ROLE FOR SLEEP DISORDERS SCREENING**

Sleep disorders have been robustly implicated in cardiometabolic disease risk. In particular, sleep apnea is a well-characterized risk factor for cardiometabolic morbidity and even cardiovascular events. Strikingly, as described above, studies that evaluated the prevalence of sleep apnea in obesity, cardiology, and diabetes clinics routinely find that most patients meet criteria for sleep apnea, though few are diagnosed and even fewer are successfully treated. For this reason, clinics that treat patients with cardiovascular disease, diabetes, and/or obesity should routinely screen patients for sleep apnea risk and refer for diagnosis appropriately.

Insomnia is also shown to be an important marker of cardiometabolic disease risk. It is, however, often ignored in the context of treatment for these conditions. Screening for insomnia is not difficult, though, with brief instruments such as the Insomnia Severity Index(237) and other brief screening measures. Of note, the recommended first line treatment for insomnia is not pharmacotherapy, although pharmacotherapy remains the most widely-utilized approach(238). This particularly important since prescription hypnotic medications may carry significant risks(239-243). The recommended first-line treatment, Cognitive Behavioral Therapy for Insomnia (CBTI) does not carry these risks and is therefore preferable for reasons beyond demonstrated efficacy(244-248), efficacy relative to medications(249-253), and effectiveness(254). Despite this, knowledge about the existence

of CBTI is limited in the medical community, and providers may be difficult to locate(255). Still, patients with insomnia should be directed to CBTI when it is available.

## **PREVENTING CARDIOVASCULAR DISEASE: A ROLE FOR HEALTHY SLEEP DURATION**

In addition to sleep disorders, it has been repeatedly shown that insufficient sleep duration is an important risk factor for cardiometabolic disease risk. In addition to many cross-sectional studies, prospective studies show that insufficient sleep is a risk factor for incident obesity, hypertension, and diabetes. For this reason (and others), achieving adequate sleep duration has been included as a Healthy People 2020 goal(213). This should be reflected in practice. Clinicians should be asking about sleep duration and consider short sleep duration among other behavioral risk factors such as poor diet, inactivity, smoking, and excessive alcohol use.

## **CONCLUSIONS**

Over 50 years ago, epidemiologic studies began to show that habitual sleep was associated with mortality(256). Since that time, substantial knowledge about the role of sleep in cardiometabolic health has emerged. Habitual short sleep duration is associated with weight gain and obesity, hypertension, hyperlipidemia, inflammation, diabetes, heart attack, and stroke. Sleep apnea is associated with cardiac and vascular dysfunction, hypertension, dyslipidemia, diabetes, inflammation, heart attack and stroke. Even insomnia, which has been traditionally seen as relatively benign, is a risk factor for cardiometabolic morbidity and mortality. Taken together, sleep represents an important domain of cardiometabolic health risk. Future studies are needed to better characterize the mechanisms by which sleep leads to morbidity and to identify and refine interventions to better ameliorate this risk. In the meantime, healthy sleep should be a consideration in the clinical care of heart disease patients.

## **ACKNOWLEDGEMENTS**

The authors wish to acknowledge the support of Dr. Sairam Parthasarathy of the University of Arizona for his assistance with the manuscript.

### **FINANCIAL SUPPORT**

Dr. Grandner is supported by K23HL110216. No other financial support was provided.

## **REFERENCES**

- \*1. Watson NF, Badr MS, Belenky G, Bliwise DL, Buxton OM, Buysse D, et al. Recommended Amount of Sleep for a Healthy Adult: A Joint Consensus Statement of the American Academy of Sleep Medicine and Sleep Research Society. *Sleep*. 2015; 38(6):843–4. [PubMed: 26039963] [This statement represents the first time that either (and in this case, both) major scientific sleep organizations has released a consensus document. This document, and the accompanying manuscript (references 4 and 5 below), provide the most complete description of the literature, conclusions based on the literature, and limitations of the existing literature, on the issue of short sleep duration.]



2. Consensus Conference P. Watson NF, Badr MS, Belenky G, Bliwise DL, Buxton OM, et al. Recommended Amount of Sleep for a Healthy Adult: A Joint Consensus Statement of the American Academy of Sleep Medicine and Sleep Research Society. *J Clin Sleep Med*. 2015; 11(6):591–2. [PubMed: 25979105]
3. Cappuccio FP, D'Elia L, Strazzullo P, Miller MA. Quantity and quality of sleep and incidence of type 2 diabetes: a systematic review and meta-analysis. *Diabetes Care*. 2010; 33(2):414–20. [PubMed: 19910503]
4. Consensus Conference P. Watson NF, Badr MS, Belenky G, Bliwise DL, Buxton OM, et al. Joint Consensus Statement of the American Academy of Sleep Medicine and Sleep Research Society on the Recommended Amount of Sleep for a Healthy Adult: Methodology and Discussion. *J Clin Sleep Med*. 2015; 11(8):931–52. [PubMed: 26235159]
5. Consensus Conference P. Watson NF, Badr MS, Belenky G, Bliwise DL, Buxton OM, et al. Joint Consensus Statement of the American Academy of Sleep Medicine and Sleep Research Society on the Recommended Amount of Sleep for a Healthy Adult: Methodology and Discussion. *Sleep*. 2015; 38(8):1161–83. [PubMed: 26194576]
6. Hirshkowitz M, Whiton K, Alpert SM, Alessi C, Bruni O, DonCarlos L, et al. National Sleep Foundation's updated sleep duration recommendations: Final report. *Sleep Health*. 2015; 1:233–43.
7. Hirshkowitz M, Whiton K, Alpert SM, Alessi C, Bruni O, DonCarlos L, et al. National Sleep Foundation's sleep time duration recommendations: Methodology and results summary. *Sleep Health*. 2015; 1:40–3.
8. Mukherjee S, Patel SR, Kales SN, Ayas NT, Strohl KP, Gozal D, et al. An Official American Thoracic Society Statement: The Importance of Healthy Sleep. Recommendations and Future Priorities. *Am J Respir Crit Care Med*. 2015; 191(12):1450–8. [PubMed: 26075423]
9. McKnight-Eily LR, Liu Y, Wheaton AG, Croft JB, Perry GS, Okoro CA, et al. Unhealthy sleep-related behaviors --- 12 States, 2009. *MMWR Morb Mortal Wkly Rep*. 2011; 60(8):233–8. [PubMed: 21368738]
10. Liu Y, Wheaton AG, Chapman DP, Cunningham TJ, Lu H, Croft JB. Prevalence of Healthy Sleep Duration among Adults - United States, 2014. *MMWR Morb Mortal Wkly Rep*. 2016; 65(6):137–41. [PubMed: 26890214]
11. Grandner MA, Smith TE, Jackson N, Jackson T, Burgard S, Branas C. Geographic distribution of insufficient sleep across the United States: a county-level hotspot analysis. *Sleep Health*. 2015; 1(3):158–65. [PubMed: 26989761]
12. Grandner MA, Patel NP, Hale L, Moore M. Mortality associated with sleep duration: The evidence, the possible mechanisms, and the future. *Sleep Med Rev*. 2010; 14:191–203. [PubMed: 19932976]
13. Youngstedt SD, Kripke DF. Long sleep and mortality: rationale for sleep restriction. *Sleep Med Rev*. 2004; 8(3):159–74. [PubMed: 15144959]
14. Gallicchio L, Kalesan B. Sleep Duration and Mortality: A Systematic Review and Meta-analysis. *J Sleep Res*. 2009; 18(2):148–58. [PubMed: 19645960]
15. Cappuccio FP, D'Elia L, Strazzullo P, Miller MA. Sleep duration and all-cause mortality: a systematic review and meta-analysis of prospective studies. *Sleep*. 2010; 33(5):585–92. [PubMed: 20469800]
16. Altman NG, Izci-Balserak B, Schopfer E, Jackson N, Rattanaumpawan P, Gehrman PR, et al. Sleep duration versus sleep insufficiency as predictors of cardiometabolic health outcomes. *Sleep Med*. 2012; 13(10):1261–70. [PubMed: 23141932]
17. Arble DM, Bass J, Behn CD, Butler MP, Challet E, Czeisler C, et al. Impact of Sleep and Circadian Disruption on Energy Balance and Diabetes: A Summary of Workshop Discussions. *Sleep*. 2015; 38(12):1849–60. [PubMed: 26564131]
18. Knutson KL. Does inadequate sleep play a role in vulnerability to obesity? *Am J Hum Biol*. 2012; 24(3):361–71. [PubMed: 22275135]
19. Grandner MA, Schopfer EA, Sands-Lincoln M, Jackson N, Malhotra A. Relationship between sleep duration and body mass index depends on age. *Obesity (Silver Spring)*. 2015; 23(12):2491–8. [PubMed: 26727118]

20. Grandner MA, Chakravorty S, Perlis ML, Oliver L, Gurubhagavatula I. Habitual sleep duration associated with self-reported and objectively determined cardiometabolic risk factors. *Sleep Med.* 2014; 15(1):42–50. [PubMed: 24333222]
21. Buxton OM, Marcelli E. Short and long sleep are positively associated with obesity, diabetes, hypertension, and cardiovascular disease among adults in the United States. *Soc Sci Med.* 2010; 71(5):1027–36. [PubMed: 20621406]
22. Hoevenaar-Blom MP, Spijkerman AM, Kromhout D, Verschuren WM. Sufficient sleep duration contributes to lower cardiovascular disease risk in addition to four traditional lifestyle factors: the MORGEN study. *European journal of preventive cardiology.* 2013
23. Hoevenaar-Blom MP, Spijkerman AM, Kromhout D, van den Berg JF, Verschuren WM. Sleep duration and sleep quality in relation to 12-year cardiovascular disease incidence: the MORGEN study. *Sleep.* 2011; 34(11):1487–92. [PubMed: 22043119]
24. Morselli LL, Guyon A, Spiegel K. Sleep and metabolic function. *Pflugers Arch.* 2012; 463(1):139–60. [PubMed: 22101912]
25. Knutson KL, Spiegel K, Penev P, Van Cauter E. The metabolic consequences of sleep deprivation. *Sleep Med Rev.* 2007; 11(3):163–78. [PubMed: 17442599]
26. Taheri S, Lin L, Austin D, Young T, Mignot E. Short sleep duration is associated with reduced leptin, elevated ghrelin, and increased body mass index. *PLoS Med.* 2004; 1(3):e62. [PubMed: 15602591]
27. Tasali E, Leproult R, Spiegel K. Reduced sleep duration or quality: relationships with insulin resistance and type 2 diabetes. *Prog Cardiovasc Dis.* 2009; 51(5):381–91. [PubMed: 19249444]
28. Van Cauter E, Spiegel K, Tasali E, Leproult R. Metabolic consequences of sleep and sleep loss. *Sleep Med.* 2008; 9(Suppl 1):S23–8. [PubMed: 18929315]
29. Spiegel K, Knutson K, Leproult R, Tasali E, Van Cauter E. Sleep loss: a novel risk factor for insulin resistance and Type 2 diabetes. *J Appl Physiol.* 2005; 99(5):2008–19. [PubMed: 16227462]
30. Spiegel K, Leproult R, L'Hermite-Baleriaux M, Copinschi G, Penev PD, Van Cauter E. Leptin levels are dependent on sleep duration: relationships with sympathovagal balance, carbohydrate regulation, cortisol, and thyrotropin. *J Clin Endocrinol Metab.* 2004; 89(11):5762–71. [PubMed: 15531540]
31. Spiegel K, Leproult R, Van Cauter E. Impact of sleep debt on metabolic and endocrine function. *Lancet.* 1999; 354(9188):1435–9. [PubMed: 10543671]
32. Patel SR. Reduced sleep as an obesity risk factor. *Obes Rev.* 2009; 10(Suppl 2):61–8. [PubMed: 19849803]
33. Spiegel K, Tasali E, Penev P, Van Cauter E. Brief communication: Sleep curtailment in healthy young men is associated with decreased leptin levels, elevated ghrelin levels, and increased hunger and appetite. *Ann Intern Med.* 2004; 141(11):846–50. [PubMed: 15583226]
34. Spaeth AM, Dinges DF, Goel N. Sex and race differences in caloric intake during sleep restriction in healthy adults. *Am J Clin Nutr.* 2014; 100(2):559–66. [PubMed: 24965304]
35. Spaeth AM, Dinges DF, Goel N. Effects of Experimental Sleep Restriction on Weight Gain, Caloric Intake, and Meal Timing in Healthy Adults. *Sleep.* 2013; 36(7):981–90. [PubMed: 23814334]
36. Greer SM, Goldstein AN, Walker MP. The impact of sleep deprivation on food desire in the human brain. *Nat Commun.* 2013; 4:2259. [PubMed: 23922121]
37. Shechter A, Grandner MA, St-Onge MP. The role of sleep in the control of food intake. *Am J Lifestyle Med.* 2014
38. St-Onge MP. The role of sleep duration in the regulation of energy balance: effects on energy intakes and expenditure. *J Clin Sleep Med.* 2013; 9(1):73–80. [PubMed: 23319909]
39. St-Onge MP, Roberts AL, Chen J, Kelleman M, O'Keefe M, RoyChoudhury A, et al. Short sleep duration increases energy intakes but does not change energy expenditure in normal-weight individuals. *Am J Clin Nutr.* 2011; 94(2):410–6. [PubMed: 21715510]
40. Markwald RR, Melanson EL, Smith MR, Higgins J, Perreault L, Eckel RH, et al. Impact of insufficient sleep on total daily energy expenditure, food intake, and weight gain. *Proc Natl Acad Sci U S A.* 2013; 110(14):5695–700. [PubMed: 23479616]

41. Grandner MA, Jackson NJ, Izci-Balserak B, Gallagher RA, Murray-Bachmann R, Williams NJ, et al. Social and Behavioral Determinants of Perceived Insufficient Sleep. *Front Neurol.* 2015; 6:112. [PubMed: 26097464]
42. Grandner MA. Addressing sleep disturbances: An opportunity to prevent cardiometabolic disease? *International review of psychiatry.* 2014; 26(2):155–76. [PubMed: 24892892]
43. Goel N, Basner M, Rao H, Dinges DF. Circadian rhythms, sleep deprivation, and human performance. *Progress in molecular biology and translational science.* 2013; 119:155–90. [PubMed: 23899598]
44. Dinges, DF.; Banks, S. Sleep deprivation: Cognitive performance.. In: Amlaner, CJ.; Fuller, PM., editors. *Basics of Sleep Guide.* 2 ed.. Sleep Research Society; Westchester, IL: 2009. p. 257-64.
45. Lim J, Dinges DF. Sleep deprivation and vigilant attention. *Ann N Y Acad Sci.* 2008; 1129:305–22. [PubMed: 18591490]
46. Banks S, Dinges DF. Behavioral and physiological consequences of sleep restriction. *J Clin Sleep Med.* 2007; 3(5):519–28. [PubMed: 17803017]
47. Maia Q, Grandner MA, Findley J, Gurubhagavatula I. Short and long sleep duration and risk of drowsy driving and the role of subjective sleep insufficiency. *Accid Anal Prev.* 2013; 59:618–22. [PubMed: 23973762]
48. Akerstedt T, Folkard S, Portin C. Predictions from the three-process model of alertness. *Aviat Space Environ Med.* 2004; 75(3 Suppl):A75–83. [PubMed: 15018267]
49. Akerstedt T, Hume K, Minors D, Waterhouse J. The subjective meaning of good sleep, an intraindividual approach using the Karolinska Sleep Diary. *Percept Mot Skills.* 1994; 79(1 Pt 1): 287–96. [PubMed: 7991323]
50. Anund A, Kecklund G, Kircher A, Tapani A, Akerstedt T. The effects of driving situation on sleepiness indicators after sleep loss: a driving simulator study. *Ind Health.* 2009; 47(4):393–401. [PubMed: 19672013]
51. Dawson D, Ian Noy Y, Harma M, Akerstedt T, Belenky G. Modelling fatigue and the use of fatigue models in work settings. *Accid Anal Prev.* 2011; 43(2):549–64. [PubMed: 21130216]
52. Goodwin RD, Marusic A. Association between short sleep and suicidal ideation and suicide attempt among adults in the general population. *Sleep.* 2008; 31(8):1097–101. [PubMed: 18714781]
53. Ryu SY, Kim KS, Han MA. Factors associated with sleep duration in Korean adults: results of a 2008 community health survey in Gwangju metropolitan city, Korea. *J Korean Med Sci.* 2011; 26(9):1124–31. [PubMed: 21935265]
54. Gangwisch JE, Malaspina D, Posner K, Babiss LA, Heymsfield SB, Turner JB, et al. Insomnia and sleep duration as mediators of the relationship between depression and hypertension incidence. *Am J Hypertens.* 2010; 23(1):62–9. [PubMed: 19893498]
55. Grandner MA, Sands-Lincoln MR, Pak VM, Garland SN. Sleep duration, cardiovascular disease, and proinflammatory biomarkers. *Nat Sci Sleep.* 2013; 5:93–107. [PubMed: 23901303]
56. Grandner MA, Buxton OM, Jackson N, Sands-Lincoln M, Pandey A, Jean-Louis G. Extreme sleep durations and increased C-reactive protein: effects of sex and ethnoracial group. *Sleep.* 2013; 36(5):769–79. [PubMed: 23633760]
57. Irwin MR, Olmstead R, Carroll JE. Sleep Disturbance, Sleep Duration, and Inflammation: A Systematic Review and Meta-Analysis of Cohort Studies and Experimental Sleep Deprivation. *Biol Psychiatry.* 2015
58. Meier-Ewert HK, Ridker PM, Rifai N, Price N, Dinges DF, Mullington JM. Absence of diurnal variation of C-reactive protein concentrations in healthy human subjects. *Clin Chem.* 2001; 47(3): 426–30. [PubMed: 11238292]
59. Ioja S, Weir ID, Rennert NJ. Relationship between sleep disorders and the risk for developing type 2 diabetes mellitus. *Postgrad Med.* 2012; 124(4):119–29. [PubMed: 22913900]
60. Lucassen EA, Rother KI, Cizza G. Interacting epidemics? Sleep curtailment, insulin resistance, and obesity. *Ann N Y Acad Sci.* 2012; 1264(1):110–34. [PubMed: 22827862]
61. Barone MT, Menna-Barreto L. Diabetes and sleep: a complex cause-and-effect relationship. *Diabetes Res Clin Pract.* 2011; 91(2):129–37. [PubMed: 20810183]

62. Rafalson L, Donahue RP, Stranges S, Lamonte MJ, Dmochowski J, Dorn J, et al. Short sleep duration is associated with the development of impaired fasting glucose: the Western New York Health Study. *Ann Epidemiol.* 2010; 20(12):883–9. [PubMed: 20620078]
63. Hall MH, Okun ML, Sowers M, Matthews KA, Kravitz HM, Hardin K, et al. Sleep is associated with the metabolic syndrome in a multi-ethnic cohort of midlife women: the SWAN Sleep Study. *Sleep.* 2012; 35(6):783–90. [PubMed: 22654197]
64. Bopparaju S, Surani S. Sleep and diabetes. *Int J Endocrinol.* 2010; 2010:759509. [PubMed: 20224753]
65. Knutson KL. Sleep duration and cardiometabolic risk: a review of the epidemiologic evidence. *Best Pract Res Clin Endocrinol Metab.* 2010; 24(5):731–43. [PubMed: 21112022]
66. Upala S, Sanguaneko A, Congrete S, Romphothong K. Sleep duration and insulin resistance in individuals without diabetes mellitus: A systematic review and meta-analysis. *Diabetes Res Clin Pract.* 2015; 109(3):e11–2. [PubMed: 26163439]
- \*67. Anothaisintawee T, Reutrakul S, Van Cauter E, Thakkinstian A. Sleep disturbances compared to traditional risk factors for diabetes development: Systematic review and meta-analysis. *Sleep Med Rev.* 2015; 30:11–24. [PubMed: 26687279] [This meta-analysis provides one of the most thorough overviews of the relationship between insufficient sleep and diabetes risk.]
68. Shan Z, Ma H, Xie M, Yan P, Guo Y, Bao W, et al. Sleep duration and risk of type 2 diabetes: a meta-analysis of prospective studies. *Diabetes Care.* 2015; 38(3):529–37. [PubMed: 25715415]
69. Holliday EG, Magee CA, Kritharides L, Banks E, Attia J. Short sleep duration is associated with risk of future diabetes but not cardiovascular disease: a prospective study and meta-analysis. *PLoS One.* 2013; 8(11):e82305. [PubMed: 24282622]
70. Lusardi P, Zoppi A, Preti P, Pesce RM, Piazza E, Fogari R. Effects of insufficient sleep on blood pressure in hypertensive patients: a 24-h study. *Am J Hypertens.* 1999; 12(1 Pt 1):63–8. [PubMed: 10075386]
71. Kaneita Y, Uchiyama M, Yoshiike N, Ohida T. Associations of usual sleep duration with serum lipid and lipoprotein levels. *Sleep.* 2008; 31(5):645–52. [PubMed: 18517035]
72. Gangwisch JE, Malaspina D, Babiss LA, Opler MG, Posner K, Shen S, et al. Short sleep duration as a risk factor for hypercholesterolemia: analyses of the National Longitudinal Study of Adolescent Health. *Sleep.* 2010; 33(7):956–61. [PubMed: 20614855]
73. Franklin KA, Lindberg E. Obstructive sleep apnea is a common disorder in the population—a review on the epidemiology of sleep apnea. *Journal of thoracic disease.* 2015; 7(8):1311–22. [PubMed: 26380759]
74. Nakayama-Ashida Y, Takegami M, Chin K, Sumi K, Nakamura T, Takahashi K, et al. Sleep-disordered breathing in the usual lifestyle setting as detected with home monitoring in a population of working men in Japan. *Sleep.* 2008; 31(3):419–25. [PubMed: 18363319]
75. Franklin KA, Sahlin C, Stenlund H, Lindberg E. Sleep apnoea is a common occurrence in females. *Eur Respir J.* 2013; 41(3):610–5. [PubMed: 22903961]
76. Resta O, Foschino-Barbaro MP, Legari G, Talamo S, Bonfitto P, Palumbo A, et al. Sleep-related breathing disorders, loud snoring and excessive daytime sleepiness in obese subjects. *Int J Obes Relat Metab Disord.* 2001; 25(5):669–75. [PubMed: 11360149]
77. Valencia-Flores M, Orea A, Castano VA, Resendiz M, Rosales M, Rebollar V, et al. Prevalence of sleep apnea and electrocardiographic disturbances in morbidly obese patients. *Obes Res.* 2000; 8(3):262–9. [PubMed: 10832770]
78. Drager LF, Lopes HF, Maki-Nunes C, Trombetta IC, Toschi-Dias E, Alves MJ, et al. The impact of obstructive sleep apnea on metabolic and inflammatory markers in consecutive patients with metabolic syndrome. *PLoS One.* 2010; 5(8):e12065. [PubMed: 20711453]
79. Pack AI, Gislason T. Obstructive sleep apnea and cardiovascular disease: a perspective and future directions. *Prog Cardiovasc Dis.* 2009; 51(5):434–51. [PubMed: 19249449]
80. Bradley TD, Floras JS. Obstructive sleep apnoea and its cardiovascular consequences. *Lancet.* 2009; 373(9657):82–93. [PubMed: 19101028]
81. Kohler M, Stradling JR. Mechanisms of vascular damage in obstructive sleep apnea. *Nat Rev Cardiol.* 2010; 7(12):677–85. [PubMed: 21079639]

82. Levy P, Ryan S, Oldenburg O, Parati G. Sleep apnoea and the heart. *European respiratory review* : an official journal of the European Respiratory Society. 2013; 22(129):333–52. [PubMed: 23997061]
83. Punjabi NM, Caffo BS, Goodwin JL, Gottlieb DJ, Newman AB, O'Connor GT, et al. Sleep-disordered breathing and mortality: a prospective cohort study. *PLoS Med*. 2009; 6(8):e1000132. [PubMed: 19688045]
84. Marin JM, Carrizo SJ, Vicente E, Agusti AG. Long-term cardiovascular outcomes in men with obstructive sleep apnoea-hypopnoea with or without treatment with continuous positive airway pressure: an observational study. *Lancet*. 2005; 365(9464):1046–53. [PubMed: 15781100]
85. Franklin KA, Nilsson JB, Sahlin C, Naslund U. Sleep apnoea and nocturnal angina. *Lancet*. 1995; 345(8957):1085–7. [PubMed: 7715342]
86. Logan AG, Perlikowski SM, Mente A, Tisler A, Tkacova R, Niroumand M, et al. High prevalence of unrecognized sleep apnoea in drug-resistant hypertension. *J Hypertens*. 2001; 19(12):2271–7. [PubMed: 11725173]
87. Mehra R, Benjamin EJ, Shahar E, Gottlieb DJ, Nawab R, Kirchner HL, et al. Association of nocturnal arrhythmias with sleep-disordered breathing: The Sleep Heart Health Study. *Am J Respir Crit Care Med*. 2006; 173(8):910–6. [PubMed: 16424443]
88. Kanagala R, Murali NS, Friedman PA, Ammash NM, Gersh BJ, Ballman KV, et al. Obstructive sleep apnea and the recurrence of atrial fibrillation. *Circulation*. 2003; 107(20):2589–94. [PubMed: 12743002]
89. Ferrier K, Campbell A, Yee B, Richards M, O'Meeghan T, Weatherall M, et al. Sleep-disordered breathing occurs frequently in stable outpatients with congestive heart failure. *Chest*. 2005; 128(4):2116–22. [PubMed: 16236863]
90. Mansfield DR, Gollogly NC, Kaye DM, Richardson M, Bergin P, Naughton MT. Controlled trial of continuous positive airway pressure in obstructive sleep apnea and heart failure. *Am J Respir Crit Care Med*. 2004; 169(3):361–6. [PubMed: 14597482]
91. Kaneko Y, Floras JS, Usui K, Plante J, Tkacova R, Kubo T, et al. Cardiovascular effects of continuous positive airway pressure in patients with heart failure and obstructive sleep apnea. *N Engl J Med*. 2003; 348(13):1233–41. [PubMed: 12660387]
92. Ryan CM, Usui K, Floras JS, Bradley TD. Effect of continuous positive airway pressure on ventricular ectopy in heart failure patients with obstructive sleep apnoea. *Thorax*. 2005; 60(9):781–5. [PubMed: 15994252]
93. Milleron O, Pilliere R, Foucher A, de Roquefeuil F, Aegerter P, Jondeau G, et al. Benefits of obstructive sleep apnoea treatment in coronary artery disease: a long-term follow-up study. *Eur Heart J*. 2004; 25(9):728–34. [PubMed: 15120882]
94. Morgenstern M, Wang J, Beatty N, Batemarco T, Sica AL, Greenberg H. Obstructive sleep apnea: an unexpected cause of insulin resistance and diabetes. *Endocrinol Metab Clin North Am*. 2014; 43(1):187–204. [PubMed: 24582098]
95. Kono M, Tatsumi K, Saibara T, Nakamura A, Tanabe N, Takiguchi Y, et al. Obstructive sleep apnea syndrome is associated with some components of metabolic syndrome. *Chest*. 2007; 131(5):1387–92. [PubMed: 17494788]
96. Zhai AB, Yip A, Haddad H. Heart failure and sleep-disordered breathing. *Curr Opin Cardiol*. 2016; 31(2):224–8. [PubMed: 26766163]
97. Yumino D, Redolfi S, Ruttanaumpawan P, Su MC, Smith S, Newton GE, et al. Nocturnal rostral fluid shift: a unifying concept for the pathogenesis of obstructive and central sleep apnea in men with heart failure. *Circulation*. 2010; 121(14):1598–605. [PubMed: 20351237]
98. Naughton MT, Benard DC, Liu PP, Rutherford R, Rankin F, Bradley TD. Effects of nasal CPAP on sympathetic activity in patients with heart failure and central sleep apnea. *Am J Respir Crit Care Med*. 1995; 152(2):473–9. [PubMed: 7633695]
99. Wang H, Parker JD, Newton GE, Floras JS, Mak S, Chiu KL, et al. Influence of obstructive sleep apnea on mortality in patients with heart failure. *J Am Coll Cardiol*. 2007; 49(15):1625–31. [PubMed: 17433953]
100. Naughton MT, Lorenzi-Filho G. Sleep in heart failure. *Prog Cardiovasc Dis*. 2009; 51(4):339–49. [PubMed: 19110135]

101. Hanly P, Zuberi N, Gray R. Pathogenesis of Cheyne-Stokes respiration in patients with congestive heart failure. Relationship to arterial PCO<sub>2</sub>. *Chest*. 1993; 104(4):1079–84. [PubMed: 8404170]
102. Sands SA, Owens RL. Congestive Heart Failure and Central Sleep Apnea. *Sleep Med Clin*. 2016; 11(1):127–42. [PubMed: 26972039]
103. Lattova Z, Keckeis M, Maurovich-Horvat E, Wetter TC, Wilde-Frenz J, Schuld A, et al. The stress hormone system in various sleep disorders. *J Psychiatr Res*. 2011; 45(9):1223–8. [PubMed: 21501849]
104. Lanfranco F, Motta G, Minetto MA, Baldi M, Balbo M, Ghigo E, et al. Neuroendocrine alterations in obese patients with sleep apnea syndrome. *Int J Endocrinol*. 2010; 2010:474518. [PubMed: 20182553]
105. Louis M, Punjabi NM. Effects of acute intermittent hypoxia on glucose metabolism in awake healthy volunteers. *J Appl Physiol*. 2009; 106(5):1538–44. [PubMed: 19265062]
106. Aronsohn RS, Whitmore H, Van Cauter E, Tasali E. Impact of untreated obstructive sleep apnea on glucose control in type 2 diabetes. *Am J Respir Crit Care Med*. 2010; 181(5):507–13. [PubMed: 20019340]
107. Hassaballa HA, Tulaimat A, Herdegen JJ, Mokhlesi B. The effect of continuous positive airway pressure on glucose control in diabetic patients with severe obstructive sleep apnea. *Sleep Breath*. 2005; 9(4):176–80. [PubMed: 16283228]
108. Babu AR, Herdegen J, Fogelfeld L, Shott S, Mazzone T. Type 2 diabetes, glycemic control, and continuous positive airway pressure in obstructive sleep apnea. *Arch Intern Med*. 2005; 165(4):447–52. [PubMed: 15738376]
109. Martinez-Ceron E, Barquiel B, Bezos AM, Casitas R, Galera R, Garcia-Benito C, et al. Effect of CPAP on Glycemic Control in Patients with Obstructive Sleep Apnea and Type 2 Diabetes. A Randomized Clinical Trial. *Am J Respir Crit Care Med*. 2016
110. Iftikhar IH, Khan MF, Das A, Magalang UJ. Meta-analysis: continuous positive airway pressure improves insulin resistance in patients with sleep apnea without diabetes. *Annals of the American Thoracic Society*. 2013; 10(2):115–20. [PubMed: 23607839]
111. Salord N, Fortuna AM, Monasterio C, Gasa M, Perez A, Bonsignore MR, et al. A Randomized Controlled Trial of Continuous Positive Airway Pressure on Glucose Tolerance in Obese Patients with Obstructive Sleep Apnea. *Sleep*. 2016; 39(1):35–41. [PubMed: 26350474]
112. Silverberg DS, Oksenberg A. Are sleep-related breathing disorders important contributing factors to the production of essential hypertension? *Curr Hypertens Rep*. 2001; 3(3):209–15. [PubMed: 11353571]
113. Peppard PE, Young T, Palta M, Skatrud J. Prospective study of the association between sleep-disordered breathing and hypertension. *N Engl J Med*. 2000; 342(19):1378–84. [PubMed: 10805822]
114. Torres G, Sanchez-de-la-Torre M, Barbe F. Relationship Between OSA and Hypertension. *Chest*. 2015; 148(3):824–32. [PubMed: 25879807]
115. Schein AS, Kerkhoff AC, Coronel CC, Plentz RD, Sbruzzi G. Continuous positive airway pressure reduces blood pressure in patients with obstructive sleep apnea; a systematic review and meta-analysis with 1000 patients. *J Hypertens*. 2014; 32(9):1762–73. [PubMed: 24979300]
116. Iftikhar IH, Valentine CW, Bittencourt LR, Cohen DL, Fedson AC, Gislason T, et al. Effects of continuous positive airway pressure on blood pressure in patients with resistant hypertension and obstructive sleep apnea: a meta-analysis. *J Hypertens*. 2014; 32(12):2341–50. discussion 50. [PubMed: 25243523]
117. Martinez-Garcia MA, Capote F, Campos-Rodriguez F, Lloberes P, Diaz de Atauri MJ, Somoza M, et al. Effect of CPAP on blood pressure in patients with obstructive sleep apnea and resistant hypertension: the HIPARCO randomized clinical trial. *JAMA*. 2013; 310(22):2407–15. [PubMed: 24327037]
118. Li J, Savransky V, Nanayakkara A, Smith PL, O'Donnell CP, Polotsky VY. Hyperlipidemia and lipid peroxidation are dependent on the severity of chronic intermittent hypoxia. *J Appl Physiol* (1985). 2007; 102(2):557–63. [PubMed: 17082365]
119. Li J, Thorne LN, Punjabi NM, Sun CK, Schwartz AR, Smith PL, et al. Intermittent hypoxia induces hyperlipidemia in lean mice. *Circ Res*. 2005; 97(7):698–706. [PubMed: 16123334]

120. Roche F, Xuong AN, Court-Fortune I, Costes F, Pichot V, Duverney D, et al. Relationship among the severity of sleep apnea syndrome, cardiac arrhythmias, and autonomic imbalance. *Pacing Clin Electrophysiol.* 2003; 26(3):669–77. [PubMed: 12698666]
121. Newman AB, Nieto FJ, Guidry U, Lind BK, Redline S, Pickering TG, et al. Relation of sleep-disordered breathing to cardiovascular disease risk factors: the Sleep Heart Health Study. *Am J Epidemiol.* 2001; 154(1):50–9. [PubMed: 11434366]
122. Chin K, Nakamura T, Shimizu K, Mishima M, Nakamura T, Miyasaka M, et al. Effects of nasal continuous positive airway pressure on soluble cell adhesion molecules in patients with obstructive sleep apnea syndrome. *Am J Med.* 2000; 109(7):562–7. [PubMed: 11063958]
123. Chin K, Shimizu K, Nakamura T, Narai N, Masuzaki H, Ogawa Y, et al. Changes in intra-abdominal visceral fat and serum leptin levels in patients with obstructive sleep apnea syndrome following nasal continuous positive airway pressure therapy. *Circulation.* 1999; 100(7):706–12. [PubMed: 10449691]
124. Cuhadaroglu C, Utkusavas A, Ozturk L, Salman S, Ece T. Effects of nasal CPAP treatment on insulin resistance, lipid profile, and plasma leptin in sleep apnea. *Lung.* 2009; 187(2):75–81. [PubMed: 19127383]
125. Dorkova Z, Petrasova D, Molcanyiova A, Popovnakova M, Tkacova R. Effects of continuous positive airway pressure on cardiovascular risk profile in patients with severe obstructive sleep apnea and metabolic syndrome. *Chest.* 2008; 134(4):686–92. [PubMed: 18625666]
126. Gozal D, Capdevila OS, Kheirandish-Gozal L. Metabolic alterations and systemic inflammation in obstructive sleep apnea among nonobese and obese prepubertal children. *Am J Respir Crit Care Med.* 2008; 177(10):1142–9. [PubMed: 18276939]
127. Steiropoulos P, Tsara V, Nena E, Fiteli C, Kataropoulou M, Froudarakis M, et al. Effect of continuous positive airway pressure treatment on serum cardiovascular risk factors in patients with obstructive sleep apnea-hypopnea syndrome. *Chest.* 2007; 132(3):843–51. [PubMed: 17573492]
128. Robinson GV, Pepperell JC, Segal HC, Davies RJ, Stradling JR. Circulating cardiovascular risk factors in obstructive sleep apnoea: data from randomised controlled trials. *Thorax.* 2004; 59(9):777–82. [PubMed: 15333855]
129. Phillips CL, Yee BJ, Marshall NS, Liu PY, Sullivan DR, Grunstein RR. Continuous positive airway pressure reduces postprandial lipidemia in obstructive sleep apnea: a randomized, placebo-controlled crossover trial. *Am J Respir Crit Care Med.* 2011; 184(3):355–61. [PubMed: 21527567]
130. Rebelo S, Drummond M, Marques JA. Lipid profile after long-term APAP in OSA patients. *Sleep Breath.* 2015; 19(3):931–7. [PubMed: 25510508]
131. Adedayo AM, Olafiranye O, Smith D, Hill A, Zizi F, Brown C, et al. Obstructive sleep apnea and dyslipidemia: evidence and underlying mechanism. *Sleep Breath.* 2012
132. Grandner MA, Perlis ML. Insomnia as a cardiometabolic risk factor. *Sleep.* 2013; 36(1):11–2. [PubMed: 23288965]
133. Grandner MA, Perlis ML. Treating Insomnia Disorder in the Context of Medical and Psychiatric Comorbidities. *JAMA internal medicine.* 2015; 175(9):1472–3. [PubMed: 26147221]
134. Ohayon MM. Epidemiology of insomnia: what we know and what we still need to learn. *Sleep Med Rev.* 2002; 6(2):97–111. [PubMed: 12531146]
135. Bixler EO, Vgontzas AN, Lin HM, Vela-Bueno A, Kales A. Insomnia in central Pennsylvania. *J Psychosom Res.* 2002; 53(1):589–92. [PubMed: 12127176]
136. LeBlanc M, Merette C, Savard J, Ivers H, Baillargeon L, Morin CM. Incidence and risk factors of insomnia in a population-based sample. *Sleep.* 2009; 32(8):1027–37. [PubMed: 19725254]
137. Singareddy R, Vgontzas AN, Fernandez-Mendoza J, Liao D, Calhoun S, Shaffer ML, et al. Risk factors for incident chronic insomnia: a general population prospective study. *Sleep Med.* 2012; 13(4):346–53. [PubMed: 22425576]
138. Fernandez-Mendoza J, Vgontzas AN, Bixler EO, Singareddy R, Shaffer ML, Calhoun SL, et al. Clinical and polysomnographic predictors of the natural history of poor sleep in the general population. *Sleep.* 2012; 35(5):689–97. [PubMed: 22547895]

139. Morin CM, Belanger L, LeBlanc M, Ivers H, Savard J, Espie CA, et al. The natural history of insomnia: a population-based 3-year longitudinal study. *Arch Intern Med.* 2009; 169(5):447–53. [PubMed: 19273774]
140. Vgontzas AN, Fernandez-Mendoza J, Bixler EO, Singareddy R, Shaffer ML, Calhoun SL, et al. Persistent insomnia: the role of objective short sleep duration and mental health. *Sleep.* 2012; 35(1):61–8. [PubMed: 22215919]
141. American Academy of Sleep Medicine. *International Classification of Sleep Disorders.* 3rd ed.. American Academy of Sleep Medicine; Darien, IL: 2014.
142. Strand LB, Laugsand LE, Wisloff U, Nes BM, Vatten L, Janszky I. Insomnia symptoms and cardiorespiratory fitness in healthy individuals: the Nord-Trøndelag Health Study (HUNT). *Sleep.* 2013; 36(1):99–108. [PubMed: 23288976]
143. Grandner MA, Kripke DF, Naidoo N, Langer RD. Relationships among dietary nutrients and subjective sleep, objective sleep, and napping in women. *Sleep Med.* 2010; 11(2):180–4. [PubMed: 20005774]
144. Grandner MA, Jackson N, Gerstner JR, Knutson KL. Sleep symptoms associated with intake of specific dietary nutrients. *J Sleep Res.* 2014; 23(1):22–34. [PubMed: 23992533]
145. Healey ES, Kales A, Monroe LJ, Bixler EO, Chamberlin K, Soldatos CR. Onset of insomnia: role of life-stress events. *Psychosom Med.* 1981; 43(5):439–51. [PubMed: 7313035]
146. Riemann D, Spiegelhalder K, Feige B, Voderholzer U, Berger M, Perlis M, et al. The hyperarousal model of insomnia: a review of the concept and its evidence. *Sleep Med Rev.* 2010; 14(1):19–31. [PubMed: 19481481]
147. Fernandez-Mendoza J, Vela-Bueno A, Vgontzas AN, Ramos-Platon MJ, Olavarrieta-Bernardino S, Bixler EO, et al. Cognitive-emotional hyperarousal as a premorbid characteristic of individuals vulnerable to insomnia. *Psychosom Med.* 2010; 72(4):397–403. [PubMed: 20368477]
148. Vgontzas AN, Tsigos C, Bixler EO, Stratakis CA, Zachman K, Kales A, et al. Chronic insomnia and activity of the stress system: a preliminary study. *J Psychosom Res.* 1998; 45(1 Spec No):21–31. [PubMed: 9720852]
149. Vgontzas AN, Bixler EO, Lin HM, Prolo P, Mastorakos G, Vela-Bueno A, et al. Chronic insomnia is associated with nyctohemeral activation of the hypothalamic-pituitary-adrenal axis: clinical implications. *J Clin Endocrinol Metab.* 2001; 86(8):3787–94. [PubMed: 11502812]
150. Rodenbeck A, Cohrs S, Jordan W, Huether G, Ruther E, Hajak G. The sleep-improving effects of doxepin are paralleled by a normalized plasma cortisol secretion in primary insomnia. A placebo-controlled, double-blind, randomized, cross-over study followed by an open treatment over 3 weeks. *Psychopharmacology (Berl).* 2003; 170(4):423–8. [PubMed: 13680082]
151. Irwin M, Clark C, Kennedy B, Christian Gillin J, Ziegler M. Nocturnal catecholamines and immune function in insomniacs, depressed patients, and control subjects. *Brain Behav Immun.* 2003; 17(5):365–72. [PubMed: 12946658]
152. Riemann D, Klein T, Rodenbeck A, Feige B, Horny A, Hummel R, et al. Nocturnal cortisol and melatonin secretion in primary insomnia. *Psychiatry Res.* 2002; 113(1-2):17–27. [PubMed: 12467942]
153. Varkevisser M, Van Dongen HP, Kerkhof GA. Physiologic indexes in chronic insomnia during a constant routine: evidence for general hyperarousal? *Sleep.* 2005; 28(12):1588–96. [PubMed: 16408419]
154. Lichstein KL, Johnson RS, Sen Gupta S, O’Laughlin DL, Dykstra TA. Are insomniacs sleepy during the day? A pupillometric assessment. *Behav Res Ther.* 1992; 30(3):283–92. [PubMed: 1586365]
155. Lichstein KL, Wilson NM, Noe SL, Aguillard RN, Bellur SN. Daytime sleepiness in insomnia: behavioral, biological and subjective indices. *Sleep.* 1994; 17(8):693–702. [PubMed: 7701180]
156. de Zambotti M, Covassin N, De Min Tona G, Sarlo M, Stegagno L. Sleep onset and cardiovascular activity in primary insomnia. *J Sleep Res.* 2011; 20(2):318–25. [PubMed: 20673289]
157. Johansson JK, Kronholm E, Jula AM. Variability in home-measured blood pressure and heart rate: associations with self-reported insomnia and sleep duration. *J Hypertens.* 2011; 29(10):1897–905. [PubMed: 21841496]



158. Jurysta F, Lanquart JP, Sputaels V, Dumont M, Migeotte PF, Leistedt S, et al. The impact of chronic primary insomnia on the heart rate--EEG variability link. *Clin Neurophysiol.* 2009; 120(6):1054–60. [PubMed: 19403330]
159. Spiegelhalder K, Fuchs L, Ladwig J, Kyle SD, Nissen C, Voderholzer U, et al. Heart rate and heart rate variability in subjectively reported insomnia. *J Sleep Res.* 2011; 20(1 Pt 2):137–45. [PubMed: 20626615]
160. Stein PK, Pu Y. Heart rate variability, sleep and sleep disorders. *Sleep Med Rev.* 2012; 16(1):47–66. [PubMed: 21658979]
161. de Zambotti M, Covassin N, Sarlo M, De Min Tona G, Trinder J, Stegagno L. Nighttime cardiac sympathetic hyper-activation in young primary insomniacs. *Clinical autonomic research : official journal of the Clinical Autonomic Research Society.* 2013; 23(1):49–56. [PubMed: 22975984]
162. Bonnet MH, Arand DL. 24-Hour metabolic rate in insomniacs and matched normal sleepers. *Sleep.* 1995; 18(7):581–8. [PubMed: 8552929]
163. Bonnet MH, Arand DL. Physiological activation in patients with Sleep State Misperception. *Psychosom Med.* 1997; 59(5):533–40. [PubMed: 9316187]
164. Lichstein KL, Johnson RS. Pupillometric discrimination of insomniacs. *Behav Res Ther.* 1994; 32(1):123–9. [PubMed: 8135710]
165. Nofzinger EA, Nissen C, Germain A, Moul D, Hall M, Price JC, et al. Regional cerebral metabolic correlates of WASO during NREM sleep in insomnia. *J Clin Sleep Med.* 2006; 2(3): 316–22. [PubMed: 17561544]
166. Bastien CH, Turcotte I, St-Jean G, Morin CM, Carrier J. Information processing varies between insomnia types: measures of N1 and P2 during the night. *Behav Sleep Med.* 2013; 11(1):56–72. [PubMed: 23347117]
167. Winkelman JW, Buxton OM, Jensen JE, Benson KL, O'Connor SP, Wang W, et al. Reduced brain GABA in primary insomnia: preliminary data from 4T proton magnetic resonance spectroscopy (1H-MRS). *Sleep.* 2008; 31(11):1499–506. [PubMed: 19014069]
168. Laugsand LE, Vatten LJ, Bjorngaard JH, Hveem K, Janszky I. Insomnia and high-sensitivity C-reactive protein: the HUNT study, Norway. *Psychosom Med.* 2012; 74(5):543–53. [PubMed: 22685243]
169. Seidel WF, Ball S, Cohen S, Patterson N, Yost D, Dement WC. Daytime alertness in relation to mood, performance, and nocturnal sleep in chronic insomniacs and noncomplaining sleepers. *Sleep.* 1984; 7(3):230–8. [PubMed: 6484427]
170. Stepanski E, Zorick F, Roehrs T, Young D, Roth T. Daytime alertness in patients with chronic insomnia compared with asymptomatic control subjects. *Sleep.* 1988; 11(1):54–60. [PubMed: 3363270]
171. Sugeran JL, Stern JA, Walsh JK. Daytime alertness in subjective and objective insomnia: some preliminary findings. *Biol Psychiatry.* 1985; 20(7):741–50. [PubMed: 4005333]
172. Vgontzas AN, Fernandez-Mendoza J, Liao D, Bixler EO. Insomnia with objective short sleep duration: the most biologically severe phenotype of the disorder. *Sleep Med Rev.* 2013; 17(4): 241–54. [PubMed: 23419741]
173. Schwartz S, McDowell Anderson W, Cole SR, Cornoni-Huntley J, Hays JC, Blazer D. Insomnia and heart disease: a review of epidemiologic studies. *J Psychosom Res.* 1999; 47(4):313–33. [PubMed: 10616226]
174. Spiegelhalder K, Scholtes C, Riemann D. The association between insomnia and cardiovascular diseases. *Nat Sci Sleep.* 2010; 2:71–8. [PubMed: 23616699]
175. Meng L, Zheng Y, Hui R. The relationship of sleep duration and insomnia to risk of hypertension incidence: a meta-analysis of prospective cohort studies. *Hypertens Res.* 2013; 36(11):985–95. [PubMed: 24005775]
176. Sofi F, Cesari F, Casini A, Macchi C, Abbate R, Gensini GF. Insomnia and risk of cardiovascular disease: a meta-analysis. *European journal of preventive cardiology.* 2014; 21(1):57–64. [PubMed: 22942213]
177. Li M, Zhang XW, Hou WS, Tang ZY. Insomnia and risk of cardiovascular disease: a meta-analysis of cohort studies. *Int J Cardiol.* 2014; 176(3):1044–7. [PubMed: 25156850]

178. Vgontzas AN, Liao D, Bixler EO, Chrousos GP, Vela-Bueno A. Insomnia with objective short sleep duration is associated with a high risk for hypertension. *Sleep*. 2009; 32(4):491–7. [PubMed: 19413143]
179. Vgontzas AN, Liao D, Pejovic S, Calhoun S, Karataraki M, Bixler EO. Insomnia with objective short sleep duration is associated with type 2 diabetes: A population-based study. *Diabetes Care*. 2009; 32(11):1980–5. [PubMed: 19641160]
180. Fernandez-Mendoza J, Vgontzas AN, Liao D, Shaffer ML, Vela-Bueno A, Basta M, et al. Insomnia with objective short sleep duration and incident hypertension: the Penn State Cohort. *Hypertension*. 2012; 60(4):929–35. [PubMed: 22892811]
181. Bathgate CJ, Edinger JD, Wyatt JK, Krystal AD. Objective but Not Subjective Short Sleep Duration Associated with Increased Risk for Hypertension in Individuals with Insomnia. *Sleep*. 2016; 39(5):1037–45. [PubMed: 26951399]
182. Li Y, Vgontzas AN, Fernandez-Mendoza J, Bixler EO, Sun Y, Zhou J, et al. Insomnia with physiological hyperarousal is associated with hypertension. *Hypertension*. 2015; 65(3):644–50. [PubMed: 25624338]
183. Jennings JR, Muldoon MF, Hall M, Buysse DJ, Manuck SB. Self-reported sleep quality is associated with the metabolic syndrome. *Sleep*. 2007; 30(2):219–23. [PubMed: 17326548]
184. Kazman JB, Abraham PA, Zeno SA, Poth M, Deuster PA. Self-reported sleep impairment and the metabolic syndrome among African Americans. *Ethn Dis*. 2012; 22(4):410–5. [PubMed: 23140070]
185. Cronlein T, Langguth B, Busch V, Rupprecht R, Wetter TC. Severe chronic insomnia is not associated with higher body mass index. *J Sleep Res*. 2015; 24(5):514–7. [PubMed: 25776276]
186. Vgontzas AN, Fernandez-Mendoza J, Miksiewicz T, Kritikou I, Shaffer ML, Liao D, et al. Unveiling the longitudinal association between short sleep duration and the incidence of obesity: the Penn State Cohort. *Int J Obes (Lond)*. 2014; 38(6):825–32. [PubMed: 24100421]
187. Rademacher ER, Jacobs DR Jr, Moran A, Steinberger J, Prineas RJ, Sinaiko A. Relation of blood pressure and body mass index during childhood to cardiovascular risk factor levels in young adults. *J Hypertens*. 2009; 27(9):1766–74. [PubMed: 19633567]
188. Taveras EM, Gillman MW, Pena MM, Redline S, Rifas-Shiman SL. Chronic sleep curtailment and adiposity. *Pediatrics*. 2014; 133(6):1013–22. [PubMed: 24843068]
189. Mitchell JA, Rodriguez D, Schmitz KH, Audrain-McGovern J. Sleep duration and adolescent obesity. *Pediatrics*. 2013; 131(5):e1428–34. [PubMed: 23569090]
190. Zhu Y, Li AM, Au CT, Kong AP, Zhang J, Wong CK, et al. Association between sleep architecture and glucose tolerance in children and adolescents. *J Diabetes*. 2015; 7(1):10–5. [PubMed: 25695111]
191. Matthews KA, Dahl RE, Owens JF, Lee L, Hall M. Sleep duration and insulin resistance in healthy black and white adolescents. *Sleep*. 2012; 35(10):1353–8. [PubMed: 23024433]
192. Koren D, Levitt Katz LE, Brar PC, Gallagher PR, Berkowitz RI, Brooks LJ. Sleep architecture and glucose and insulin homeostasis in obese adolescents. *Diabetes Care*. 2011; 34(11):2442–7. [PubMed: 21933909]
193. Navarro-Solera M, Carrasco-Luna J, Pin-Arboledas G, Gonzalez-Carrascosa R, Soriano JM, Codoner-Franch P. Short Sleep Duration Is Related to Emerging Cardiovascular Risk Factors in Obese Children. *J Pediatr Gastroenterol Nutr*. 2015; 61(5):571–6. [PubMed: 25988561]
194. Marcus CL, Brooks LJ, Draper KA, Gozal D, Halbower AC, Jones J, et al. Diagnosis and management of childhood obstructive sleep apnea syndrome. *Pediatrics*. 2012; 130(3):576–84. [PubMed: 22926173]
195. Hartzell K, Avis K, Lozano D, Feig D. Obstructive sleep apnea and periodic limb movement disorder in a population of children with hypertension and/or nocturnal nondipping blood pressures. *J Am Soc Hypertens*. 2016; 10(2):101–7. [PubMed: 26725017]
196. Bixler EO, Vgontzas AN, Lin HM, Liao D, Calhoun S, Fedok F, et al. Blood pressure associated with sleep-disordered breathing in a population sample of children. *Hypertension*. 2008; 52(5):841–6. [PubMed: 18838624]

197. Amin RS, Carroll JL, Jeffries JL, Grone C, Bean JA, Chini B, et al. Twenty-four-hour ambulatory blood pressure in children with sleep-disordered breathing. *Am J Respir Crit Care Med.* 2004; 169(8):950–6. [PubMed: 14764433]
198. Amin RS, Kimball TR, Kalra M, Jeffries JL, Carroll JL, Bean JA, et al. Left ventricular function in children with sleep-disordered breathing. *Am J Cardiol.* 2005; 95(6):801–4. [PubMed: 15757619]
199. Anna Theresa Weber S, Pierri Carvalho R, Ridley G, Williams K, El Dib R. A systematic review and meta-analysis of cohort studies of echocardiographic findings in OSA children after adenotonsillectomy. *Int J Pediatr Otorhinolaryngol.* 2014; 78(10):1571–8. [PubMed: 25108873]
200. Gozal D, Kheirandish-Gozal L, Serpero LD, Sans Capdevila O, Dayyat E. Obstructive sleep apnea and endothelial function in school-aged nonobese children: effect of adenotonsillectomy. *Circulation.* 2007; 116(20):2307–14. [PubMed: 17967978]
201. Baharav A, Kotagal S, Rubin BK, Pratt J, Akselrod S. Autonomic cardiovascular control in children with obstructive sleep apnea. *Clin Auton Res.* 1999; 9(6):345–51. [PubMed: 10638809]
202. Kheirandish-Gozal L, Capdevila OS, Tauman R, Gozal D. Plasma C-reactive protein in nonobese children with obstructive sleep apnea before and after adenotonsillectomy. *J Clin Sleep Med.* 2006; 2(3):301–4. [PubMed: 17410279]
203. Bhushan B, Ayub B, Loghmanee DA, Billings KR. Metabolic alterations in adolescents with obstructive sleep apnea. *Int J Pediatr Otorhinolaryngol.* 2015; 79(12):2368–73. [PubMed: 26581829]
204. Centers for Disease Control and Prevention. CDC Health Disparities and Inequalities Report -- United States, 2011. CDC; Atlanta, GA: 2011.
205. Adler NE, Rehkopf DH. U.S. disparities in health: descriptions, causes, and mechanisms. *Annu Rev Public Health.* 2008; 29:235–52. [PubMed: 18031225]
206. Smedley, BD.; Stith, AY.; Nelson, AR.; Institute of Medicine (U.S.). Unequal treatment : confronting racial and ethnic disparities in health care. Vol. xvi. National Academy Press; Washington, D.C.: 2003. Committee on Understanding and Eliminating Racial and Ethnic Disparities in Health Care.; p. 764
207. Ong KL, Cheung BM, Wong LY, Wat NM, Tan KC, Lam KS. Prevalence, treatment, and control of diagnosed diabetes in the U.S. National Health and Nutrition Examination Survey 1999-2004. *Ann Epidemiol.* 2008; 18(3):222–9. [PubMed: 18201902]
208. Gallo LC, Matthews KA. Understanding the association between socioeconomic status and physical health: do negative emotions play a role? *Psychol Bull.* 2003; 129(1):10–51. [PubMed: 12555793]
209. Jackson, JS.; Hudson, D.; Kershaw, K.; Mezuk, B.; Rafferty, J.; Tuttle, KK. Discrimination, chronic stress, and mortality among black Americans: A life course framework.. In: Rogers, RG.; Crimmins, EM., editors. *International Handbook of Adult Mortality. International Handbooks of Population.* Springer; NY: 2011. p. 311-28.
210. Jackson JS, Knight KM, Rafferty JA. Race and unhealthy behaviors: chronic stress, the HPA axis, and physical and mental health disparities over the life course. *Am J Public Health.* 2010; 100(5): 933–9. [PubMed: 19846689]
211. Finch BK, Phuong Do D, Heron M, Bird C, Seeman T, Lurie N. Neighborhood effects on health: Concentrated advantage and disadvantage. *Health Place.* 2010; 16(5):1058–60. [PubMed: 20627796]
212. Williams DR, Sternthal M, Wright RJ. Social determinants: taking the social context of asthma seriously. *Pediatrics.* 2009; 123(Suppl 3):S174–84. [PubMed: 19221161]
213. Office of Disease Prevention and Health Promotion. *Healthy People 2020 Objective Topic Areas.* US Department of Health and Human Services; Washington, DC: 2011.
214. Ruitter ME, Decoster J, Jacobs L, Lichstein KL. Normal sleep in African-Americans and Caucasian-Americans: A meta-analysis. *Sleep Med.* 2011; 12(3):209–14. [PubMed: 21317037]
215. Hong S, Mills PJ, Loreda JS, Adler KA, Dimsdale JE. The association between interleukin-6, sleep, and demographic characteristics. *Brain Behav Immun.* 2005; 19(2):165–72. [PubMed: 15664789]

216. Mokhlesi B, Pannain S, Ghods F, Knutson KL. Predictors of slow-wave sleep in a clinic-based sample. *J Sleep Res.* 2012; 21(2):170–5. [PubMed: 21955220]
217. Mezick EJ, Matthews KA, Hall M, Strollo PJ Jr. Buysse DJ, Kamarck TW, et al. Influence of race and socioeconomic status on sleep: Pittsburgh SleepSCORE project. *Psychosom Med.* 2008; 70(4):410–6. [PubMed: 18480189]
218. Redline S, Kirchner HL, Quan SF, Gottlieb DJ, Kapur V, Newman A. The effects of age, sex, ethnicity, and sleep-disordered breathing on sleep architecture. *Arch Intern Med.* 2004; 164(4): 406–18. [PubMed: 14980992]
219. Hall MH, Matthews KA, Kravitz HM, Gold EB, Buysse DJ, Bromberger JT, et al. Race and financial strain are independent correlates of sleep in midlife women: the SWAN sleep study. *Sleep.* 2009; 32(1):73–82. [PubMed: 19189781]
220. Whinnery J, Berkowitz Sturgis EK, Staley B, Corbitt C, Gehrman P, Chakravorty S, et al. Sleep architecture differences between good sleepers and patients with primary insomnia. *SLEEP.* 2013; 36:A212. (Abstract Supplement).
- \*221. Grandner MA, Williams NJ, Knutson KL, Roberts D, Jean-Louis G. Sleep disparity, race/ethnicity, and socioeconomic position. *Sleep Med.* 2015 [This review represents an in-depth characterizing of the relationship between race/ethnicity and sleep parameters, covering duration, symptoms, and disorders. It also addresses potential causes of these differences and role in cardiovascular health disparities.]
222. Adenekan B, Pandey A, McKenzie S, Zizi F, Casimir GJ, Jean-Louis G. Sleep in America: role of racial/ethnic differences. *Sleep Med Rev.* 2013; 17(4):255–62. [PubMed: 23348004]
223. Knutson KL. Sociodemographic and cultural determinants of sleep deficiency: implications for cardiometabolic disease risk. *Soc Sci Med.* 2013; 79:7–15. [PubMed: 22682665]
224. Mezick EJ, Hall M, Matthews KA. Sleep duration and ambulatory blood pressure in black and white adolescents. *Hypertension.* 2012; 59(3):747–52. [PubMed: 22275538]
225. Hale L, Do DP. Racial differences in self-reports of sleep duration in a population-based study. *Sleep.* 2007; 30(9):1096–103. [PubMed: 17910381]
226. Ruiter ME, DeCoster J, Jacobs L, Lichstein KL. Sleep disorders in African Americans and Caucasian Americans: a meta-analysis. *Behav Sleep Med.* 2010; 8(4):246–59. [PubMed: 20924837]
227. Grandner MA, Petrov ME, Rattanaumpawan P, Jackson N, Platt A, Patel NP. Sleep symptoms, race/ethnicity, and socioeconomic position. *J Clin Sleep Med.* 2013; 9(9):897–905. A–D. [PubMed: 23997702]
228. Grandner MA, Petrov MER, Rattanaumpawan P, Jackson N, Platt A, Patel NP. Sleep Symptoms, Race/Ethnicity, and Socioeconomic Position. *Journal of Clinical Sleep Medicine.* 2013; 9(9): 897–905. [PubMed: 23997702]
229. Kripke DF, Jean-Louis G, Elliott JA, Klauber MR, Rex KM, Tuunainen A, et al. Ethnicity, sleep, mood, and illumination in postmenopausal women. *BMC Psychiatry.* 2004; 4:8. [PubMed: 15070419]
230. Boss EF, Smith DF, Ishman SL. Racial/ethnic and socioeconomic disparities in the diagnosis and treatment of sleep-disordered breathing in children. *Int J Pediatr Otorhinolaryngol.* 2011; 75(3): 299–307. [PubMed: 21295865]
231. Patel NP, Grandner MA, Xie D, Branas CC, Gooneratne N. “Sleep disparity” in the population: poor sleep quality is strongly associated with poverty and ethnicity. *BMC Public Health.* 2010; 10(1):475. [PubMed: 20701789]
232. Moore M, Kirchner HL, Drotar D, Johnson N, Rosen C, Redline S. Correlates of adolescent sleep time and variability in sleep time: The role of individual and health related characteristics. *Sleep Med.* 2011; 12(3):239–45. [PubMed: 21316300]
233. Hale L, Rivero-Fuentes E. Negative acculturation in sleep duration among Mexican immigrants and Mexican Americans. *J Immigr Minor Health.* 2011; 13(2):402–7. [PubMed: 19728094]
234. Grandner MA, Patel NP, Gehrman PR, Xie D, Sha D, Weaver T, et al. Who gets the best sleep? Ethnic and socioeconomic factors related to sleep disturbance. *Sleep Med.* 2010; 11:470–9. [PubMed: 20388566]

235. Gellis LA, Lichstein KL, Scarinci IC, Durrence HH, Taylor DJ, Bush AJ, et al. Socioeconomic status and insomnia. *J Abnorm Psychol*. 2005; 114(1):111–8. [PubMed: 15709817]
236. Knutson KL, Van Cauter E, Rathouz PJ, Yan LL, Hulley SB, Liu K, et al. Association between sleep and blood pressure in midlife: the CARDIA sleep study. *Arch Intern Med*. 2009; 169(11):1055–61. [PubMed: 19506175]
237. Bastien CH, Vallieres A, Morin CM. Validation of the Insomnia Severity Index as an outcome measure for insomnia research. *Sleep Med*. 2001; 2(4):297–307. [PubMed: 11438246]
238. Moloney ME, Konrad TR, Zimmer CR. The medicalization of sleeplessness: a public health concern. *Am J Public Health*. 2011; 101(8):1429–33. [PubMed: 21680913]
239. Kripke DF. Possibility that certain hypnotics might cause cancer in skin. *J Sleep Res*. 2008; 17(3):245–50. [PubMed: 18844818]
240. Kripke, DF. Evidence that new hypnotics cause cancer. 2008.
241. Kripke DF. Do hypnotics cause death and cancer? The burden of proof. *Sleep Med*. 2009; 10(3):275–6. [PubMed: 19269891]
242. Kripke DF. Chronic hypnotic use: deadly risks, doubtful benefit. REVIEW ARTICLE. *Sleep Med Rev*. 2000; 4(1):5–20. [PubMed: 12531158]
243. Kripke DF, Simons RN, Garfinkel L, Hammond EC. Short and long sleep and sleeping pills. Is increased mortality associated? *Arch Gen Psychiatry*. 1979; 36(1):103–16. [PubMed: 760693]
244. Okajima I, Komada Y, Inoue D. A meta-analysis on the treatment effectiveness of cognitive behavioral therapy for primary insomnia. *Sleep and Biological Rhythms*. 2011; 9(1):24–34.
245. Murtagh DR, Greenwood KM. Identifying effective psychological treatments for insomnia: a meta-analysis. *J Consult Clin Psychol*. 1995; 63(1):79–89. [PubMed: 7896994]
246. Irwin MR, Cole JC, Nicassio PM. Comparative meta-analysis of behavioral interventions for insomnia and their efficacy in middle-aged adults and in older adults 55+ years of age. *Health Psychol*. 2006; 25(1):3–14. [PubMed: 16448292]
247. Morin CM, Culbert JP, Schwartz SM. Nonpharmacological interventions for insomnia: a meta-analysis of treatment efficacy. *Am J Psychiatry*. 1994; 151(8):1172–80. [PubMed: 8037252]
248. Mitchell MD, Gehrman P, Perlis M, Umscheid CA. Comparative effectiveness of cognitive behavioral therapy for insomnia: a systematic review. *BMC Fam Pract*. 2012; 13:40. [PubMed: 22631616]
249. McClusky HY, Milby JB, Switzer PK, Williams V, Wooten V. Efficacy of behavioral versus triazolam treatment in persistent sleep-onset insomnia. *Am J Psychiatry*. 1991; 148(1):121–6. [PubMed: 1888345]
250. Jacobs GD, Pace-Schott EF, Stickgold R, Otto MW. Cognitive behavior therapy and pharmacotherapy for insomnia: a randomized controlled trial and direct comparison. *Arch Intern Med*. 2004; 164(17):1888–96. [PubMed: 15451764]
251. Smith MT, Perlis ML, Park A, Smith MS, Pennington J, Giles DE, et al. Comparative meta-analysis of pharmacotherapy and behavior therapy for persistent insomnia. *Am J Psychiatry*. 2002; 159(1):5–11. [PubMed: 11772681]
252. Morin CM, Colecchi C, Stone J, Sood R, Brink D. Behavioral and pharmacological therapies for late-life insomnia: a randomized controlled trial. *JAMA*. 1999; 281(11):991–9. [PubMed: 10086433]
253. Sivertsen B, Omvik S, Pallesen S, Bjorvatn B, Havik OE, Kvale G, et al. Cognitive behavioral therapy vs zopiclone for treatment of chronic primary insomnia in older adults: a randomized controlled trial. *JAMA*. 2006; 295(24):2851–8. [PubMed: 16804151]
- \*254. Wu JQ, Appleman ER, Salazar RD, Ong JC. Cognitive Behavioral Therapy for Insomnia Comorbid With Psychiatric and Medical Conditions: A Meta-analysis. *JAMA internal medicine*. 2015; 175(9):1461–72. [PubMed: 26147487] [This meta-analysis shows that behavioral interventions for insomnia are effective even if the patient suffers from a comorbidity such as depression or chronic pain. Further, this meta-analysis shows that the effects are about as large as they are for primary insomnia.]
255. Thomas A, Grandner M, Nowakowski S, Nesom G, Corbitt C, Perlis ML. Where are the Behavioral Sleep Medicine Providers and Where are They Needed? A Geographic Assessment. *Behav Sleep Med*. 2016:1–12.

256. Hammond EC. Some Preliminary Findings on Physical Complaints from a Prospective Study of 1,064,004 Men and Women. *Am J Public Health Nations Health*. 1964; 54:11-23.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**KEY POINTS**

- Insufficient sleep duration is prevalent in up to 1/3 of US adults and is associated with weight gain, diabetes, hypertension, inflammation, and mortality.
- Insomnia disorder is prevalent in 5-15% of US adults and is associated with cardiometabolic disease risk, especially when paired with objective short sleep duration.
- Sleep apnea is prevalent among men and women in the US and is a strong predictor of cardiometabolic morbidity and mortality, especially if left untreated.
- Pediatric sleep disorders such as insomnia and sleep apnea are related to cardiovascular disease risk.
- Sleep may play an important and potentially modifiable role in cardiovascular health disparities.