

Northumbria Research Link

Citation: Hajirasouliha, Farzaneh and Zabiegaj, Dominika (2021) Is the shape of air particulate matter important? A study on the interactions of the two shapes of titanium dioxide nanoparticles with a model of pulmonary surfactant. In: UK Fluids Conference 8th-10th September 2021: Book of Abstracts. University of Southampton, Southampton, p. 64.

Published by: University of Southampton

URL: <https://www.southampton.ac.uk/engineering/news/eve...>
<<https://www.southampton.ac.uk/engineering/news/events/2021/09/10-uk-fluids-conference-2021.page?>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/47298/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

Is the Shape of Air Particulate Matter Important?
A study on the Interactions of the Two Shapes of Titanium Dioxide
Nanoparticles with a Model of Pulmonary Surfactant

Farzaneh Hajirasouliha¹, Dominika Zabiegaj¹

¹ Smart Materials and Surfaces Laboratory, Department of Mechanical and Construction Engineering, Faculty of Engineering and Environment, Northumbria University, NE1 8ST, Newcastle Upon Tyne, United Kingdom

Air quality indices are measured based on the concentration and size of the particles. But should the shape of these particles be considered as well? Upon inhalation, Air Particulate Matter with a size of less than or equal to $2.5\mu\text{m}$ (PM_{2.5}) can reach the deepest areas of the respiratory system called alveolar region where the gas exchange with blood circulation happens. In this zone, the walls of almost 500 million tiny alveoli sacs are lined with a tiny layer of fluid which causes a high surface tension and instability in this large air-liquid interface. However, thanks to the lung surfactant, there is no collapse of this region in a healthy human. Therefore, the first barrier that PM_{2.5} encounter before reaching the blood circulation is the interface covered by a monolayer of lung surfactant molecules. The purpose of this research is to study the interfacial properties of these monolayers interacted with titanium (IV) oxide nanoparticles with two different shapes, spherical and irregular ones, and with the same average size of 20 nm. Dipalmitoyl phosphatidylcholine (DPPC) as one of the main constituents of the pulmonary surfactant was used as the synthetic model. Using Profile Analysis Tensiometry (PAT) as an automatic set-up working based on Young-Laplace equation, we measured the interfacial tension and surface dilatational viscoelasticity in the pendant drop mode. The temperature was constant at 37°C. Four different amplitudes, 1%, 2%, 5%, and 10%, for the volume change of drops were used. Moreover, various frequencies, 0.1, 0.125, 0.25, and 0.5 Hz, were applied as the representative of the breathing cycle at different ages for a healthy human. The dependence of the interfacial and mechanical properties of the bespoke monolayers on the shape of the nanoparticles can be considered as an important factor for the environmental authorities.