

Northumbria Research Link

Citation: Mazi, Waleed, Sangal, Vartul, Saeed, Amir, Sandstrom, Gunnar, Weill, François-Xavier, Yu, Jun and Ledebøer, N. A. (2015) Rapid Genotyping of *Shigella sonnei* by Use of Multiplex High-Resolution Melting. *Journal of Clinical Microbiology*, 53 (7). pp. 2389-2391. ISSN 0095-1137

Published by: American Society for Microbiology

URL: <https://doi.org/10.1128/JCM.00874-15> <<https://doi.org/10.1128/JCM.00874-15>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/23342/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



**Northumbria
University**
NEWCASTLE



UniversityLibrary

New-Data Letter: Bacteriology	1
Rapid genotyping of <i>Shigella sonnei</i> using multiplex high resolution melting	2
	3
Running title: Multiplex-HRM typing of <i>Shigella sonnei</i>	4
Waleed Mazi ^{a, b} , Vartul Sangal ^c , Amir Saeed ^a , Gunnar Sandstorm ^a , François-Xavier	5
Weill ^d , Jun Yu ^{b, *}	6
^a Karolinska Institute, Department of Laboratory Medicine, Division of Clinical Microbiology,	7
Karolinska University Hospital, Huddinge, SE- 141 86 Stockholm, Sweden	8
^b Strathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, United	9
Kingdom	10
^c Faculty of Health and Life Sciences, Northumbria University, Newcastle upon Tyne NE1 8ST,	11
United Kingdom	12
^d Institut Pasteur, Unité des Bactéries Pathogènes Entériques, Centre National de Référence des	13
<i>Escherichia coli</i> , <i>Shigella</i> et <i>Salmonella</i> , Paris, France	14
* Correspondence: Jun Yu, Strathclyde Institute of Pharmacy and Biomedical Sciences,	15
University of Strathclyde, 161 Cathedral Street, Glasgow, G4 0RE, United Kingdom. Phone	16
number: 0044 141 548 2137. Email: jun.yu@strath.ac.uk	17
Keywords: <i>Shigella sonnei</i> , high-resolution melting (HRM), melting temperature (<i>T_m</i>), single	18
nucleotide polymorphism (SNP)	19
	20

Shigella sonnei, an emerging global cause of shigellosis, consists of four distinct lineages and the current pandemic involves several geographically associated, multidrug-resistant clones that belong to lineage III (1-3). A typing scheme based on high resolution melting (HRM) of six chromosomal single nucleotide polymorphisms (SNPs) has been described to identify all lineages/sub-lineages (4). HRM detects SNPs in small DNA fragments based on the melting patterns (5, 6). HRM involves minimal labor with rapid turnaround time, and a decreased risk of PCR carryover contamination (7). A sensitive intercalating fluorescent dye, EvaGreen, may yield robust fluorescence intensities and strong sharp melting peaks on melting of the PCR amplicons (8-10). Based on the SNPs summarized in Table 1, we have developed a multiplex-HRM, for identification of lineages I, II, and III using the EvaGreen and a Rotor-Gene 6000 instrument (Qiagen).

Ten strains (2 for each lineage/sublineage) were from a collection held at Institut Pasteur. Only one lineage IV strain is reported so far (1), which is not included in the study. Primers were designed to amplify fragments that contain three SNPs within *kduD*, *deoA* and *emrA* for typing 3 main lineages in a first set run, and to amplify fragments that contain SNPs within *fdx* and *menF* for typing sub-lineages IIIa, IIIb, and IIIc in a second run. Primer sequences are summarized in Table 1.

In the first set run, a 20 µl multiplex real-time PCR reaction was set up, containing 10 µl SensiMix™ with EvaGreen Dye (Bioline), 20 ng of template DNA, and 2.5 µl of each primer of *kduD* (100µM), 1 µl of each primer of *deoA* (100µM) and 1 µl of each primer of *emrA* (50µM). In the second set run, the 20 µl reaction mix was identical, except that the primers used were *fdx* (10 µM) and *menF* (10µM), with a quantity of 2 µl and 1µl, respectively. The PCR reaction was programmed as following: hold at 95°C for 10 min followed by 35 cycles of 95°C for 20s, 60°C for 20s and 72°C for 22s. HRM was performed between 80 and 90°C with a stepwise increase of 0.05°C. The melting temperatures (*T_m*) were determined by the negative derivative of decrease

fluorescence over increased temperature (df/dt) for each SNP using the proprietary software (version 1.7.34).

The first set run clearly separated lineages I, II, and III with distinctive melting curves, and the T_m of each allele was at least half degree apart from that of the other allele (Fig. 1A). The second set run distinguished the sub-lineages IIIa, IIIb, and IIIc with distinctive melting curves (Fig. 1B). Table 2 summarizes T_m for all alleles. To assess the reproducibility, we blindly run all 10 samples of various lineages randomly in same condition and revealed same results with +/- 0.05 different in T_m . Sequencing of five SNPs independently verified reliability of T_m data.

A simple and robust genotyping report is important for rapid identification of the emerging multidrug-resistant clones of lineage III (1). The multiplex-HRM approach we present here can be very useful in this regard.

- References 58
1. Holt KE, Baker S, Weill FX, Holmes EC, Kitchen A, Yu J, Sangal V, Brown DJ, Coia JE, Kim DW, Choi SY, Kim SH, da Silveira WD, Pickard DJ, Farrar JJ, Parkhill J, Dougan G, Thomson NR. 2012. Shigella sonnei genome sequencing and phylogenetic analysis indicate recent global dissemination from Europe. *Nature genetics* **44**:1056-1059. 59
 2. Seol SY, Kim YT, Jeong YS, Oh JY, Kang HY, Moon DC, Kim J, Lee YC, Cho DT, Lee JC. 2006. Molecular characterization of antimicrobial resistance in Shigella sonnei isolates in Korea. *Journal of medical microbiology* **55**:871-877. 60
 3. Wei HL, Wang YW, Li CC, Tung SK, Chiou CS. 2007. Epidemiology and evolution of genotype and antimicrobial resistance of an imported Shigella sonnei clone circulating in central Taiwan. *Diagnostic microbiology and infectious disease* **58**:469-475. 61
 4. Sangal V, Holt KE, Yuan J, Brown DJ, Filliol-Toutain I, Weill FX, Kim DW, da Silveira WD, Pickard D, Thomson NR, Parkhill J, Yu J. 2013. Global phylogeny of Shigella sonnei strains from limited single nucleotide polymorphisms (SNPs) and development of a rapid and cost-effective SNP-typing scheme for strain identification by high-resolution melting analysis. *Journal of clinical microbiology* **51**:303-305. 62
 5. Garritano S, Gemignani F, Voegelé C, Nguyen-Dumont T, Le Calvez-Kelm F, De Silva D, Lesueur F, Landi S, Tavtigian SV. 2009. Determining the effectiveness of High Resolution Melting analysis for SNP genotyping and mutation scanning at the TP53 locus. *BMC genetics* **10**:5. 63
 6. Martino A, Mancuso T, Rossi AM. 2010. Application of high-resolution melting to large-scale, high-throughput SNP genotyping: a comparison with the TaqMan method. *Journal of biomolecular screening* **15**:623-629. 64
 7. Cheng JC, Huang CL, Lin CC, Chen CC, Chang YC, Chang SS, Tseng CP. 2006. Rapid detection and identification of clinically important bacteria by high-resolution melting analysis after broad-range ribosomal RNA real-time PCR. *Clinical chemistry* **52**:1997-2004. 65
 8. Taylor CF. 2009. Mutation scanning using high-resolution melting. *Biochemical Society transactions* **37**:433-437. 66
 9. Erali M, Voelkerding KV, Wittwer CT. 2008. High resolution melting applications for clinical laboratory medicine. *Experimental and molecular pathology* **85**:50-58. 67
 10. Khan SA, Sung K, Nawaz MS. 2011. Detection of aacA-aphD, qacEdelta1, marA, floR, and tetA genes from multidrug-resistant bacteria: Comparative analysis of real-time multiplex PCR assays using EvaGreen((R)) and SYBR((R)) Green I dyes. *Molecular and cellular probes* **25**:78-86. 68

Figure legend

Fig.1. Multiplex-HRM curve of two set runs. A: multiplex-HRM curve of three SNPs in *kduD*, *deoA* and *emrA* respectively (from left to right), which separates lineages I, II and III (strain ID 1263, 55623 and 54184 respectively). B: multiplex-HRM curve of two SNPs in *fdX* (left) and *menF* (right), respectively, which separates sub-lineages IIIa, IIIb, and IIIc (strain ID 54184, 2073, and 6224 respectively).

93

94

95

96

97

98

99

100

Table 1: SNPs and primers used for multiplex-MRM-typing *S. sonnei* in this study:

Gene	SNP position	Ancestor	Derived	Lineage I	Lineage II	Sublineage IIIa	Sublineage IIIb	Sublineage IIIc
<i>kduD</i>	3155111	T	C	C	T	T	T	T
<i>deoA</i>	4803842	G	A	G	A	G	G	G
<i>emrA</i>	2973337	T	C	C	C	T	T	T
<i>fdx</i>	2760031	G	A	G	G	G	A	A
<i>menF</i>	2455693	A	G	A	A	A	A	G
Primers for <i>kduD</i>		forward 5' -CGACGGCGAAACACTTTATC-3' reverse 5' -CGCGTATAAGAAGGCACACG-3'						
Primers for <i>deoAD</i>		forward 5' -GGAGATGCTTATCTCCGGCAAA-3' reverse 5' -AGTCGGTTGGGCCTTT-3'						
Primers for <i>kduD</i>		forward 5' -TGCCACCGAAGTACGTAACG-3' reverse 5' -CATCCACCCACATATTGGTG-3'						
Primers for <i>kduD</i>		forward 5' -CAAAGCCTGGGACTGGA-3' reverse 5' -CATGGTTGATAGTGAACGC-3'						
Primers for <i>kduD</i>		forward 5' -TATTCTCGCGCTGGTTTTA-3' reverse 5' -GCTTTTCTTGGCTCTTCACC-3'						

Note: These SNPs were extracted using *S. sonnei* strain Ss046 as the reference as described by Holt et al., 2012 (1) and Sangal et al., 2013 (4).

Table 2: Summary of T_m for each of the five SNPs used for multiplex-HMR

Main lineages	T_m (+/- 0.05)		
	<i>kduD</i>	<i>deoA</i>	<i>emrA</i>
I	83.0	85.1	87.7
II	82.5	84.8	87.7
III	82.5	85.2	87.3
Sub-lineages	T_m (+/- 0.05)		
	<i>fdX</i>	<i>menF</i>	
IIIa	82.5	85.7	
IIIb	82.5	86.0	
IIIc	82.9	85.7	



