

Northumbria Research Link

Citation: Marrs, Emma, Jones, Amanda, Cummings, Stephen and Perry, John (2014) Demonstration of earlier detection of *Salmonella* species from stool samples by using chromogenic media. In: ESCMID: European Society of Clinical Microbiology and Infectious Diseases, 10-13 May 2014, Barcelona, Spain.

URL:

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/42297/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



**Northumbria
University
NEWCASTLE**



University Library

Demonstration of earlier detection of *Salmonella* species from stool samples by using chromogenic media.

Marrs E.C.L.¹, Jones, A.L.², Cummings, S.P.², & Perry, J.D.^{1*}

¹Freeman Hospital, Newcastle upon Tyne, UK ²Northumbria University, Newcastle upon Tyne, UK

(*john.perry@nuth.nhs.uk)



Introduction

Background: Salmonellosis is a worldwide public health issue and non-typhoid species are one of the most common causative agents of gastroenteritis in the western world.¹ Typhoidal and Paratyphoidal salmonellae cause systemic syndromes characterised by sustained bacteraemia.² Although the number of cases is under reported and therefore the incidence rates are underestimated,³ worldwide up to 1.3 billion non-typhoidal and an estimated 20 million typhoidal cases of *Salmonella* infection are reported annually.^{4,5}

Purpose of the Study: The isolation of salmonellae involves culturing faecal samples directly onto one or more selective culture media, with and without enrichment in Selenite broth. Xylose Lysine Deoxycholate (XLD) agar remains the UK standard selective medium and relies upon the absence of fermentation of lactose and sucrose with or without the production of hydrogen sulphide (H_2S) by salmonellae.⁶ However, this medium is not reliable for some serotypes, is prone to a false positives and overgrowth with commensals e.g. *E. coli*.⁷ In recent years, chromogenic media have been made available for the rapid detection of pathogens in the clinical laboratory.⁸ The aim of the current study was to assess three selective media for the isolation and discrimination of *Salmonella* spp. in stool samples from patients with diarrhoea.

Materials and Methods

Two chromogenic agar media, *Brilliance Salmonella* (Oxoid, Basingstoke, UK) and ASAP medium (bioMérieux / AES, Marcy-l'Étoile, France), were compared with XLD (bioMérieux). The three pre-prepared media were evaluated with 280 stool samples from patients suffering from diarrhoea. Approximately 0.5g or 0.5 ml of sample was suspended in 0.5 ml of saline (0.85 %). The three media were inoculated with 50 µL aliquots of the stool suspension, the remainder of which was inoculated into selenite F broth (bioMérieux). All plates and broths underwent aerobic incubation at $37 \pm 0.5^\circ\text{C}$ for 18 h. After incubation, selenite broths were subcultured onto the same three media, which were incubated in the same conditions for 18 h. All plates were interpreted after 18 h of incubation and any suspect colonies resembling *Salmonella* were identified by Maldi-TOF mass spectrometry (Bruker, Coventry, UK).

References

- Schönenbrücher V, Mallinson ET and Bülte M (2008) A comparison of standard cultural methods for the detection of foodborne *Salmonella* species including three new chromogenic plating media. *International Journal of Food Microbiology*. **123**(1-2): 61-66
- Crawford RW, Reeve KE, and Gunn JS (2010) Flagellated but Not Hyperfimbriated *Salmonella enterica* Serovar Typhimurium Attaches to and Forms Biofilms on Cholesterol-Coated Surfaces. *Journal of Bacteriology*. **192**(12): 2981-2990
- Wong MH and Chen S (2013) First detection of oqxAB in *Salmonella* spp. isolated from food. *Antimicrob. Agents Chemother*. **57**(1):658-60
- Crump JA, Luby SP and Mintz ED (2004) The global burden of typhoid fever. *Bulletin of the World Health Organization*. **82**(5):346-53.
- Coburn B, Grassl GA and Finlay BB (2007) *Salmonella*, the host and disease: a brief review. *Immunology and Cell Biology*. **85**(2):112-8
- UK Standards for Microbiology Investigations - Investigation of Faecal Specimens for Enteric Pathogens. Health Protection Agency. 2013. http://www.hpa.org.uk/webc/hpawebfile/hpaweb_c/1317132856754
- Warburton DW, Bowen B, Konkle A, et al. (1994) A comparison of six different plating media used in the isolation of *Salmonella*. *International Journal of Food Microbiology*. **22**: 277-289
- Perry JD and Freydière AM (2007) The application of chromogenic media in clinical microbiology. *Journal of Applied Microbiology*. **103**(6):2046-55

Results and Conclusions


Salmonella spp. produced purple colonies on *Brilliance Salmonella* and ASAP medium, and black or colourless colonies on XLD. On both chromogenic media commensal bacteria were invariably non-coloured.

Table: The sensitivity and specificity of three culture media for the isolation of *Salmonella* spp.

Medium	Positives	True Positives	False positives	Sensitivity (%)	Specificity (%)
XLD	- Direct	15	9	6	31
	- enrichment	38	28	10	97
BRILLIANCE	- Direct	23	13	10	45
	- enrichment	39	29	10	100
ASAP	- Direct	20	14	6	48
	- enrichment	35	29	6	100

Twenty nine isolates of *Salmonella* were recovered using a combination of all three media and all of these were recovered on *Brilliance Salmonella* and ASAP medium, as opposed to 28 on XLD, following enrichment in selenite broth. Although there was little difference in *Salmonella* recovery after enrichment, there were differences in performance after direct culture. ASAP medium recovered significantly more isolates of *Salmonella* spp. than XLD medium (McNemar's test; $P = 0.04$) whereas ASAP isolated only one additional isolate when compared with *Brilliance Salmonella*. The least number of false positive colonies were recovered on ASAP medium ($n = 12$) compared with *Brilliance Salmonella* ($n = 20$) and XLD ($n = 16$). Data from this study suggests that the use of chromogenic agars can result in earlier detection of *Salmonella* species from some samples, although enrichment in selenite broth remains essential.

Figure: *Salmonella* sp. as black colonies on XLD (a) and purple colonies on *Brilliance Salmonella* (b) and ASAP medium (c) isolated from direct culture.



Acknowledgements

The authors are most grateful to bioMérieux, La-Balme-les-Grottes, France for sponsorship of this study.