

Northumbria Research Link

Citation: Jajja, A., Sutyarjoko, A., Lara Gallegos, Jose, Rennie, K., Brandt, K., Qadir, O. and Siervo, M. (2014) Beetroot supplementation lowers daily systolic blood pressure in older, overweight subjects. *Nutrition Research*, 34 (10). pp. 868-875. ISSN 0271-5317

Published by: Elsevier

URL: <https://doi.org/10.1016/j.nutres.2014.09.007>
<<https://doi.org/10.1016/j.nutres.2014.09.007>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/43882/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

1 Title:
2 **Beetroot supplementation lowers daily systolic blood pressure in**
3 **older, overweight subjects**
4

5 **Authors:**Jajja A^{1#}, Sutayoko A^{1#},Lara J¹, Rennie K², Brandt K³, Qadir O³, Siervo M^{1*}

6 ¹Human Nutrition Research Centre, Institute for Ageing and Health, Newcastle University,
7 Campus for Ageing and Vitality, Newcastle upon Tyne, NE4 5PL, UK

8 ²Centre for Lifespan and Chronic Illness Research, University of Hertfordshire, Hatfield
9 Hertfordshire, AL10 9AB

10 ³Human Nutrition Research Centre, School of Agriculture, Food & Rural Development,
11 Newcastle University,Newcastle upon Tyne, NE1 7RU, UK

12

13 **Running title:** Beetroot juice, overweight and blood pressure

14

15

16 *Corresponding author: Dr Mario Siervo, Human Nutrition Research Centre, Institute for
17 Ageing and Health, Newcastle University, Campus for Ageing and Vitality, Newcastle upon
18 Tyne, NE4 5PL, UK. Email: mario.siervo@ncl.ac.uk

19 #Equal Contribution

20

21 The material presented in this manuscript is original and it has not been submitted for
22 publication to any other journal while under consideration forNutrition Research

23

24

25 Abstract word count: 248

26 Main text word count: 3051

27 References: 26

28 Tables: 1

29 Figures: 4

30 Online Supplementary Material: 1

31 **Abstract**

32 Although inorganic nitrate and beetroot juice supplementation are associated with decreased
33 systolic blood pressure (BP), these results have primarily been obtained from short-term trials
34 that focused on healthy young adults. Therefore, we hypothesized that oral supplementation
35 of beetroot juice concentrate would decrease systolic BP in overweight older participants but
36 that the decline in BP would not be sustained after a 1-week interruption of the beetroot juice
37 supplementation. For 3 weeks, 24 participants were randomized to either the beetroot juice
38 concentrate or blackcurrant juice group, with a 1-week postsupplementation phase (week 4).
39 Changes in systolic and diastolic BP were assessed during the supplementation and
40 postsupplementation phases. Blood pressure was measured using 3 different methods:
41 (1) resting clinic BP, (2) 24-hour ambulatory BP monitoring, and (3) home monitoring of daily
42 resting BP. The first 2 methods were applied at baseline and after weeks 3 and 4. Daily
43 measurements were conducted throughout the study, with 21 subjects completing the study
44 (beetroot/blackcurrant = 10/11; male/female = 12/9; age = 62.0 ± 1.4 years; body mass index
45 =
46 30.1 ± 1.2 kg/m²). After 3 weeks, beetroot juice supplementation was not associated with
47 significant changes in resting clinic BP or 24-hour ABPM. Conversely, beetroot juice
48 concentrate reduced daily systolic BP after 3 weeks (-7.3 ± 5.9 mm Hg, $P = .02$); however, the
49 effect was not maintained after the interruption of the supplementation (week 4, 2.8 ± 6.1 mm
50 Hg, $P = .09$). In overweight older subjects, beetroot juice concentrate supplementation was
51 associated with beneficial effects on daily systolic BP, although the effects were not significant
52 when measured by 24-hour ABPM or resting clinic BP.

53

54 **Trial registration: ISRCTN85926713**

55 **Keywords:** ageing, blood pressure, obesity, beetroot juice concentrate, inorganic nitrate,
56 cardiovascular risk, Human

57

58

59 **1. Introduction**

60 High blood pressure (BP) is responsible for nearly 5% of the global disease burden and an
61 important risk factor for cardiovascular diseases [1–3]. The risk of hypertension is increased
62 among elderly and overweight individuals [4,5]. Results from the 2009 Health Survey for
63 England reported that 47% of women and 51% of men 55 and 64 years of age were classified
64 as hypertensive [6].

65

66 Initiatives that target modifiable risk factors, such as diet, tobacco use, and physical activity
67 levels, are important for the prevention and management of high BP [5]. Recent studies
68 attribute the beneficial BP-lowering effects of dietary patterns, such as the Dietary Approach
69 to Stop Hypertension (DASH) approach, to an increased intake of inorganic nitrate [7]. A
70 recent meta-analysis of 16 trials showed that inorganic nitrate and beetroot juice
71 supplementation were associated with a significant reduction in systolic BP (−4.4 mm Hg),
72 whereas no significant effect was observed for diastolic BP [8]. The trials were generally
73 characterized by a short duration (from 1 to 15 days) and predominantly focused on young
74 subjects. The majority of these studies reported beneficial effects on resting clinic BP.
75 However, no effects have been shown in studies on type 2 diabetic patients evaluating BP
76 with 24-hour ambulatory BP monitoring (ABMP) [9].

77 In this study, we hypothesized that oral supplementation of beetroot juice concentrate would
78 decrease systolic BP in overweight older participants and that the effect on BP would
79 be independent of the BP measurement protocol (clinic, 24-hour ambulatory, daily). We also
80 conjectured that the decline in BP would not be sustained after a 1-week interruption of the
81 beetroot juice supplementation. To test our hypothesis, a 3-week supplementation period with
82 beetroot juice concentrate was implemented among older, overweight participants. We
83 investigated whether potential changes in BP were sustained after the disruption of beetroot

84 supplementation. Three established methods were used for the measurement of BP (ie,
85 resting clinic BP, 24-hour ABPM, and home daily BP monitoring) to minimize the
86 confounding influence of the research environment on BP readings and to assess within-day
87 (24-ABMP) and between-day (daily monitoring) changes in BP.

88 **2. Methods and materials**

89 2.1 Subjects: A total of 24 nonsmoking subjects (13men/11 women), between 55 and 70
90 years of age and with body mass indexes (BMIs) between 25 and 40 kg/m², were recruited
91 between March and December 2013. Participants were excluded if they had medical
92 conditions (ie, diabetes, cancer, systemic inflammatory disorders) or were taking medications
93 (ie, organic nitrates, corticosteroids, diuretics, hormonal therapies, weight loss medications)
94 that could interfere with the nutritional intervention and study outcomes. In addition, subjects
95 were excluded if they were vegetarian, they reported alcohol intake greater than 21 U/wk for
96 men and 14 U/wk for women, or their body weight had changed more than 3 kg within the
97 last month. Written informed consent was obtained from all participants prior to participation
98 in the study. The study was approved by the Newcastle University Medical Ethics
99 Committee.

100

101 2.2 The study was a 28-day, 2-arm, parallel, randomized clinical trial. Participants and
102 researchers were not blind to the study interventions. A block randomization procedure was
103 used to generate the randomization order of the interventions, with 6 participants for each
104 block. RandList (DatInf GmbH, Tübingen, Germany) for Windows was used to perform the
105 randomization. The study was divided into 3 phases: (1) screening, (2) intervention
106 (weeks 1-3), and (3) postsupplementation (week 4).

107 2.3 Study Protocol: telephone screening interview was conducted to ensure eligibility of the
108 participants. Participants, in fasting conditions, visited the NU-Food Research Facilities at

109 Newcastle University for a 2-hour assessment visit. Anthropometric measurements (weight,
110 height, and waist circumference) were obtained following standard procedures, and BMI was
111 calculated as weight in kilograms divided by the squared height in meters. Participants were
112 then randomized to one of two interventions (beetroot or blackcurrant juice) and underwent
113 baseline assessment of resting BP, collection of urine and saliva samples, measurement of
114 body composition via leg-to-leg bioelectrical impedance analysis [10] (Tanita BC420 MA;
115 Tanita Corporation, Tokyo, Japan), and calculation of physical activity using the International
116 Physical Activity Questionnaire [11]. At the end of the visit, participants were fitted with a
117 24-hour AMBP monitor to continuously assess BP over the next 24-hour period. For 28 days,
118 participants were also instructed to measure their daily resting BP at home. The intervention
119 phase started immediately after the completion of the 24-hour BP monitoring period and
120 lasted for 21 days. During this phase, each participant was expected to comply with the
121 assigned nutritional intervention and low-nitrate dietary plan. On day 7, participants were
122 instructed to collect urine samples for the assessment of nitrate concentration. At the end of
123 the 3 weeks, participants returned to the research unit to repeat the set of measurements
124 performed at baseline. At the end of the second visit, participants were fitted again with a 24-
125 hour BP monitor and reminded that they were not required to drink either blackcurrant or
126 beetroot juice over the following 7 days (post supplementation phase) but were still required
127 to perform the daily resting BP measurements and follow the low-nitrate dietary plan.
128 On the last day of the study (28th day), participants were invited to attend the research
129 facility for their final visit and assessment. At the end of the visit, participants were fitted
130 with a 24-hour BP monitor. A summary of the study design is shown in Fig. S1 of the Online
131 Supplementary Material.

132 2.4 Nutritional Supplementation: Participants randomized to the intervention group were
133 asked to consume 70 mL of concentrated beetroot juice for 21 days (Beet-It Sport Shot;

134 James White Company Ltd, Ipswich, UK; 71 kcal). This dose provided approximately 300 to
135 400 mg of nitrate per bottle, according to the information reported on the label. A similar
136 dose of inorganic nitrate was also used in another 2-week trial testing the effects of beetroot
137 juice on BP and insulin sensitivity in older obese subjects [9]. However, after the study, we
138 analyzed the contents of 3 remaining bottles and found a nitrate concentration of 165 ± 2
139 mg per bottle; these results need to be interpreted with caution in consideration of the small
140 number of beetroot samples analyzed, and future studies are therefore required to confirm
141 these findings.

142 Participants randomized to the control group were asked to consume 200 mL of blackcurrant
143 juice (Capri-Sun Blackcurrant Juice, Rudolf Wild GmbH & Co. KG, Eppelheim,
144 Germany, 100 kcal), containing 2.7 ± 0.1 mg of nitrate per bottle. The choice of the
145 blackcurrant juice as a neutral source of inorganic nitrate was confirmed by a previous study
146 conducted by Vanhatalo et al [12]. Participants of both groups were instructed to consume the
147 drinks in the morning and were required to follow a low-nitrate diet as a means to
148 standardize nitrate intake during the study period. A low-nitrate diet limits the consumption
149 of processed/ cured meats and patés, mature cheese, and green leafy vegetables (eg, rocket,
150 spinach, cabbage, beetroot). The nitrate intake estimated by following this dietary scheme
151 was approximately 50 to 100 mg/d, which is similar to the typical dietary nitrate intake of the
152 British population [13]. During the study period, participants were also requested to not
153 change their daily physical activity patterns, to avoid the use of mouthwash during the study,
154 to limit alcohol and caffeine consumption, and to avoid consuming different sources/types of
155 water.

156 **2.5 Resting Blood Pressure Measurements:** Resting BP was measured in triplicate using an
157 automated BP monitor (Omron M2 Basic, Omron Healthcare, UK) with the patient seated
158 comfortably for 15 min prior to measurement and the arm supported at the level of the heart.

159 The final value was calculated as the mean of the lowest two measurements. A large cuff was
160 used for obese subjects.

161

162 2.6 24-hrABPM: A validated device approved by British Hypertension Society was used to
163 monitor 24-hr systolic and diastolic BP (Mobil-O-Graph NG, I.E.M. GmbH). All participants
164 were instructed on the use of the device. During monitoring BP was measured every 30
165 minutes between 8.00am and 22.00 pm, and every 60 minutes between 22.00 pm and 8.00
166 am. Patients were advised to continue their normal activity during the monitoring period.

167 Over 70% of BP measurements were considered valid and were included in final analyses.

168 All of the valid recordings were analyzed to obtain average 24-hour systolic and diastolic BP.

169

170 2.7 Daily Home Blood Pressure Monitoring: An automated BP monitor (Omron M2 Basic,
171 Omron Healthcare, UK) was provided to each participant for the measurements of daily
172 resting BP at home. Participants were asked to conduct duplicate measurements in seated
173 position in the morning before drinking the juice and in the evening before going to bed.

174 Participants were trained on the correct use of the monitor, with emphasis on the position of
175 the cuff. In addition, written instructions on the measurement of BP were provided. A form
176 was given to participants to record the BP readings (systolic, diastolic and HR) and time of
177 each measurement. Agreement was verified against the BP recordings obtained from the 24-
178 hr ABPM at baseline and end of intervention phase (Day 21). The average of the morning
179 and evening daily measurements was calculated; the average of the values for each week
180 (week 1, week 2, week 3, week 4) was calculated to evaluate differences in BP between the
181 two interventions.

182

183 2.8 Nitrate Concentrations: A modified version of the method proposed by Tsikas et al [14]

184 was used to determine nitrate concentrations in urine and saliva samples using gas
185 chromatography–mass spectrometry (GC-MS). The validation and protocol of the modified
186 GC-MS method are described elsewhere [15]. This method showed a good repeatability, as
187 coefficients of variation for replicate analyses of samples were 7.8% and 8.6% in saliva and
188 urine samples, respectively. The analyses were performed by a single operator who was blind
189 to the treatment allocation.

190 2.9 Sample Size: A total sample size of 24 participants (i.e, 12 per group) was required to
191 detect a significant difference of 5.0 mmHg (SD: 4.0mmHg) in systolic BP between the two
192 groups with a power and significance levels set at 0.80 and 0.05, respectively. These
193 calculations are based on a t-test for independent samples performed using G-Power 3 for
194 Windows.

195

196 2.10 Statistical Analysis: All statistical analyses were completed using SPSS for Windows
197 (SPSS, version 17.0; SPSS Inc, Chicago, Ill, USA). Summary data are presented as mean (SD
198 or SEM). Data were checked for normality of distribution using Q-Q plots and appropriate
199 transformations were applied to skewed distributions. T-test for independent samples was
200 used to compare between-groups differences on baseline measurements and changes in
201 salivary and urinary nitrate concentrations at each phase. A general linear model was used to
202 test differences in resting clinic, 24-hr ABMP and home daily BP readings between the two
203 nutritional interventions. Analyses testing differences between the two groups during the
204 intervention phase were adjusted for baseline values of the selected outcome. Analyses
205 testing differences between the two groups during the post-supplementation phase (Day 28)
206 were adjusted for values measured at Day 21 for each selected outcome. P values < 0.05 (2-
207 tailed) were considered as statistically significant.

208

209 **3. Results**

210 3.1 Recruitment and Baseline: The flowchart describing the recruitment process is described
211 in **Figure 1**. A total of 12 males and 9 females completed the study. **Table 1** shows the mean
212 and range of baseline characteristics of the subjects. Mean age was 62.7 ± 1.5 years in the
213 beetroot group and 61.4 ± 1.3 years in the blackcurrant group. BMI was 30.5 ± 1.3 and 29.4 ± 4.4
214 kg/m^2 in the beetroot group and blackcurrant group, respectively. There were no significant
215 differences in BP readings between the two intervention groups. Only 24-hr ABPM HR was
216 slightly higher in the blackcurrant group ($p=0.03$).

217

218 3.2 Safety and Compliance: Both interventions were well tolerated. The most commonly self-
219 reported side effect with beetroot juice was red urine (beeturia); in addition, one subject
220 experienced mild, temporary abdominal discomfort (beetroot juice concentrate group) and
221 another one reported lower urinary tract symptoms (blackcurrant juice group). Both subjects
222 completed the study. Urinary and salivary nitrate concentrations at baseline were similar
223 between the two groups. However, the beetroot juice group was associated with a significant
224 rise in urinary and salivary nitrate concentrations during the intervention phase ($p<0.001$),
225 which returned to baseline levels at the end of the post-supplementation phase (**Figure 2A**
226 **and 2B**). Body weight and physical activity level did not change during the study (**Figure S2**
227 **of the Online Supplementary Material**).

228

229 3.3 Resting Clinic BP: Resting clinic systolic BP showed a tendency to decline during the
230 study but differences between the two groups were not significant at the end of the
231 intervention ($p=0.31$) and post-supplementation ($p=0.82$) phases. Similarly, changes in
232 resting clinic diastolic BP were not significant between the two groups at the end of both
233 study phases (intervention, $p=0.87$; post-supplementation, $p=0.64$) (**Figure 3A and Table S1**

234 **of the Online Supplementary Material).**

235

236 3.4 24-hr ABPM: Beetroot juice supplementation was not associated with significant changes
237 in systolic ($p=0.87$) and diastolic ($p=0.91$) compared to blackcurrant juice. After the
238 interruption of the supplementation, 24-hr ABPM readings remained essentially unchanged
239 for both systolic ($p=0.64$) and diastolic ($p=0.74$) BP (**Figure 3B and Table S1 of the Online**
240 **Supplementary Material).**

241

242 3.5 Daily Home BP Monitoring: A progressive decrease in systolic BP during the
243 intervention phase was observed in the beetroot juice group. Systolic BP was not significantly
244 lower compared to blackcurrant juice during week 1 (-3.3 ± 6.5 mmHg, $p=0.56$) or week 2 ($-$
245 4.9 ± 5.9 mmHg, $p=0.27$), but was significant during the final week of the intervention (week
246 3, -7.3 ± 5.9 mmHg, $p=0.02$). After the interruption of the intervention, there was a progressive
247 increase in systolic BP in the beetroot juice group and values returned to baseline levels on
248 the last day of the trial ($p=0.09$). No significant changes in daily diastolic BP readings were
249 observed in either group throughout the study (**Figure 4**).

250

251 **4. Discussion**

252 This is the first study to use 3 different methods (clinic, daily, 24-hour ABPM) to assess
253 changes in BP during beetroot supplementation in older overweight and obese subjects.
254 Furthermore, for the first time, we also tested whether the BP-lowering effect of beetroot
255 juice was sustained after the interruption of the supplementation. Our results support our
256 hypothesis stating that a 3-week beetroot juice supplementation would lead to a progressive
257 decline in daily systolic BP. In addition, the 1-week interruption of beetroot supplementation
258 resulted in a progressive return of the daily systolic BP to baseline levels. We also found a

259 nonsignificant effect of beetroot juice concentrate on clinic and 24-hour ambulatory BP.

260 Choosing the most accurate and precise method for measuring BP is a complicated matter in

261 clinical and research settings for monitoring the effectiveness of specific dietary, lifestyle,

262 and pharmacological treatments [16]. Although the use of clinic resting BP is the most

263 commonly used method, its poor reliability is universally recognized due to measurement

264 bias associated with white-coat syndrome, standardization of protocol, and operator bias [17].

265 Some of these issues are resolved by the recommended adoption of 24-hour ABPM and daily

266 home monitoring for a more objective assessment of BP [18]. However, the 24-hour ABPM

267 method may be prone to measurement bias (lack of precision), mostly related to the lack

268 of standardization of the measurement conditions that include differences in posture (seated,

269 standing), physical activity, and emotional status [19]. These factors may have contributed to

270 the between-method differences. Home BP readings are shown to provide more reproducible

271 results due to controlled conditions [20]. Furthermore, our findings of the significant BP-

272 lowering effects observed with the home measurements but not with 24-hour ABPM might

273 be related to transient effects of inorganic nitrate supplementation on resting BP that were not

274 preserved during times of movement or activity. However, further research is needed to

275 investigate this topic. Previous trials have shown that beetroot juice is associated with acute

276 positive effects on seated clinic systolic BP in healthy, young volunteers [21–25]. Our study

277 did not find an effect on clinic and 24-hour ABPM, but we demonstrated a significant drop in

278 systolic daily BP. It is possible that a lack of effect on 24-hour ambulatory BP may relate to a

279 lower nitrate dose as compared to studies reporting beneficial effects of beetroot juice on BP.

280 However, a single bottle of beetroot juice was considered sufficient to increase nitrate intake

281 over the 3-week period in view of the reported nitrate content of the bottle and similar

282 amount of nitrate supplementation in another study [9]. The underlying mechanism is most

283 likely due to an increase in nitric oxide (NO) bioavailability as a result of an increased

284 nonenzymatic reduction of nitrate into nitrite and NO[26,27]. The association of inorganic
285 nitrate with BP is shown by the rise of systolic BP within 24 hours after interrupting the
286 supplementation and the return to baseline after 3 days. This seems to coincide with the
287 pharmacokinetic effect of inorganic nitrate in humans ($t_{1/2}$: ~8 hours; ~60% of ingested
288 nitrate is excreted in the urine within 48 hours) [28]. These results suggest that continuous
289 inorganic nitrate supplementation may be necessary to sustain beneficial cardiovascular
290 effects. Although the results on clinic BP are not aligned to the current evidence, the lack of
291 an effect on 24-hour ABPM was also reported by Gilchrist et al [9], who found a
292 nonsignificant effect of 2-week beetroot juice supplementation on 24-ABMP in older
293 obese diabetic patients. The only other study that recruited older subjects tested the effects of
294 a 3-day beetroot supplementation on resting BP and observed a significant reduction in both
295 systolic and diastolic BP [29]. No other study has tested the effects of either inorganic nitrate
296 or beetroot juice supplementation on daily BP. The study is characterized by several
297 limitations. First, the small sample size and the discontinuation of 3 participants may have
298 reduced the power of the study to detect significant differences in BP. However, 24-hour
299 ABMP and clinic BP were essentially unchanged; and therefore, it is unlikely that the
300 lack of power influenced the lack of a significant effect of the beetroot intervention on these 2
301 outcomes. Second, our study was not a blinded trial; and measurements may have been
302 unconsciously influenced by the allocation to the interventions. Third, participants were
303 asked to manually record their daily BP readings, which may introduce a bias in the recording
304 of BP measurements. We found a high level of compliance to the measurements and daily
305 home monitoring, with the 24-hour BP readings taken at the same times and thus indicating
306 reliable BP reporting. Similarly, participants were not asked to record their physical activity
307 during the 24-hour ABPM recording period. However, participants received detailed
308 instructions to avoid strenuous physical activities during this period. We only provided 1

309 bottle of beetroot juice, whereas previous studies commonly used higher doses. In
310 addition, our estimation of nitrate concentrations in the beetroot juice consumed in this study
311 showed a lower content than that reported by the manufacturer. However, the
312 administration of the beetroot juice concentrate determined increases greater than 100% in
313 both urinary and salivary nitrate concentrations. Therefore, careful interpretation of
314 these results is recommended; and the results should be verified in future studies. In addition,
315 the blackcurrant and beetroot juice interventions had different volumes (200 vs 70 mL).
316 However, it is unlikely that this marginal volume difference influenced the results
317 considering that we did not look at the acute effects on BP or control for the overall fluid
318 intake during the study. Whether this may have been a factor in determining the
319 nonsignificant decline in clinic and 24-hour- ABPM is currently not known, but the
320 measurement of urinary and salivary nitrate indicates a marked increase in inorganic nitrate
321 intake. Lastly, we did not obtain blood samples in this study to assess plasma nitrite levels or
322 biomarkers of NO bioactivity, such as cyclic guanosine monophosphate, arginine, or
323 methylated arginines. In older, overweight subjects, beetroot juice supplementation
324 was associated with beneficial effects on daily systolic BP; but the effects were not
325 significant when measured by 24-hour ABPM. The ability to detect an effect of beetroot juice
326 concentrate supplementation in older, overweight subjects appeared to be influenced by the
327 method that was used to measure BP. The daily monitoring of BP could represent a
328 sensitive method to evaluate the efficacy of nutritional interventions on BP. Future research is
329 needed to investigate the role of vascular aging in modulating the responsiveness of
330 BP to nutritional interventions targeting the NO pathway.

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346 **Acknowledgements**

347 We thank Lisa Gilder for her assistance during the trial. The corresponding author (MS) is the
348 guarantor for the manuscript and had full access to all of the data in the study and takes
349 responsibility for the integrity of the data and the accuracy of the data analysis. All authors
350 read and approved the final version of the paper. The study was intramural support of core
351 budget. All authors have no conflicts of interest to declare.

352

353

354 **Conflicts of Interests:** All authors have no conflicts of interest to declare

355 **References**

- 356 [1] Staessen JA, Wang J, Bianchi G, Birkenhager WH. Essential hypertension. *Lancet*.
357 2003;361:1629-41.
- 358 [2] Kaplan NM, Shanthi; Poulter, Neil; Whitworth, Judith. 2003 World Health
359 Organization (WHO)/International Society of Hypertension (ISH) statement on
360 management of hypertension. UK: Lippincott Williams & Wilkins; 2003.
- 361 [3] Centre NCG. Hypertension - NICE Clinical Guidelines. UK: NHS; 2011.
- 362 [4] Gelber RP, Gaziano JM, Manson JE, Buring JE, Sesso HD. A prospective study of
363 body mass index and the risk of developing hypertension in men. *American Journal of*
364 *Hypertension*.20:370-7.
- 365 [5] Whelton PK, He J, Appel LJ, Cutler JA, Havas S, Kotchen TA, et al. Primary
366 prevention of hypertension: clinical and public health advisory from The National High
367 Blood Pressure Education Program. *JAMA*.288:1882-8.
- 368 [6] HSE E. Health and lifestyles. 2009.
- 369 [7] Hord NG, Tang Y, Bryan NS. Food sources of nitrates and nitrites: the physiologic
370 context for potential health benefits. *American Journal of Clinical Nutrition*.90:1-10.
- 371 [8] Siervo M, Lara J, Ogbonmwan I, Mathers JC. Inorganic nitrate and beetroot juice
372 supplementation reduces blood pressure in adults: a systematic review and meta-
373 analysis. *The Journal of nutrition*. 2013;143:818-26.
- 374 [9] Gilchrist M, Winyard PG, Aizawa K, Anning C, Shore A, Benjamin N. Effect of
375 dietary nitrate on blood pressure, endothelial function, and insulin sensitivity in type 2
376 diabetes. *Free radical biology & medicine*. 2013;60:89-97.
- 377 [10] Jebb SA, Siervo M, Murgatroyd PR, Evans S, Fruhbeck G, Prentice AM. Validity
378 of the leg-to-leg bioimpedance to estimate changes in body fat during weight loss and
379 regain in overweight women: a comparison with multi-compartment models. *Int J Obes*
380 *(Lond)* 2007;31:756–62.
- 381 [11] Craig CL, Marshall AL, Sjöström M, Bauman AE, Booth ML, Ainsworth BE, et al.
382 International Physical Activity Questionnaire: 12-country reliability and validity. *Med*
383 *Sci Sports Exerc* 2003;35:1381–95.
- 384 [12] Vanhatalo A, Bailey SJ, Blackwell JR, DiMenna FJ, Pavey TG, Wilkerson DP, et
385 al. Acute and chronic effects of dietary nitrate supplementation on blood pressure and
386 the physiological responses to moderate-intensity and incremental exercise. *Am J Physiol*
387 *Regul Integr Comp Physiol* 2010;299:R1121–31.
- 388 [13] Knight T, Forman D, Al-Dabbagh S, Doll R. Estimation of dietary intake of nitrate

389 and nitrite in Great Britain. *Food Chem Toxicol* 1987;25:277–85.

390 [14] Tsikas D, Schwarz A, Stichtenoth DO. Simultaneous measurement of [15N]nitrate
391 and [15N]nitrite enrichment and concentration in urine by gas chromatography mass
392 spectrometry as pentafluorobenzyl derivatives. *Anal Chem* 2010;82:2585–7.

393 [15] Qadir O, Teh J, Siervo M, Seal C, Brandt K. Method using gas chromatography
394 mass spectrometry (GC-MS) for analysis of nitrate and nitrite in vegetables. In:
395 D’Haene K, editor. *NUTRIHORT nutrient management, innovative techniques
396 and nutrient legislation in intensive horticulture for an improved water quality
397 September 16-18, 2013, Ghent, Belgium. September 16-18, 2013. Ghent: Institute for
398 Agricultural and Fisheries Research; 2013.*

399 [16] Pickering TG, Hall JE, Appel LJ, Falkner BE, Graves J, Hill MN, et al.
400 **Recommendations for blood pressure measurement in humans and experimental
401 animals: part 1: blood pressure measurement in humans: a statement for professionals
402 from the Subcommittee of Professional and Public Education of the American Heart
403 Association Council on High Blood Pressure Research. *Hypertension* 2005;45:142–61.**

404 [17] Beevers G, Lip GYH, O’Brien E. Blood pressure measurement. *BMJ*
405 **2001;322:981–5.**

406 [18] O’Brien E, Asmar R, Beilin L, Imai Y, Mancia G, Mengden T, et al. Practice
407 guidelines of the European Society of Hypertension for clinic, ambulatory and self blood
408 pressure measurement. *J Hypertens* 2005;23:697–701.

409 [19] O’Brien E, Parati G, Stergiou G, Asmar R, Beilin L, Bilo G, et al. European Society
410 of Hypertension position paper on ambulatory blood pressure monitoring. *J Hypertens*
411 **2013;31: 1731–68. <http://dx.doi.org/10.1097/HJH.0b013e328363e964>.**

412 [20] Parati G, Stergiou GS, Asmar R, Bilo G, de Leeuw P, Imai Y, et al. European
413 Society of Hypertension practice guidelines for home blood pressure monitoring. *J Hum
414 Hypertens* 2010;24:779–85.

415 [21] Webb AJ, Patel N, Loukogeorgakis S, Okorie M, Aboud Z, Misra S, et al. Acute
416 blood pressure lowering, vasoprotective, and antiplatelet properties of dietary nitrate
417 via bioconversion to nitrite. *Hypertension* 2008;51:784–90.

418 [22] Bahra M, Kapil V, Pearl V, Ghosh S, Ahluwalia A. Inorganic nitrate ingestion
419 improves vascular compliance but does not alter flow-mediated dilatation in healthy
420 volunteers. *Nitric Oxide* 2012;26:197–202.

421 [23] Hobbs DA, Kaffa N, George TW, Methven L, Lovegrove JA. Blood pressure-
422 lowering effects of beetroot juice and novel beetroot-enriched bread products in

423 normotensive male subjects. *Br J Nutr* 2012;108:2066–74.

424 [24] Kapil V, Milsom AB, Okorie M, Maleki-Toyserkani S, Akram F, Rehman F, et al.

425 Inorganic nitrate supplementation lowers blood pressure in humans: role for nitrite-

426 derived NO. *Hypertension* 2010;56:274–81.

427 [25] Larsen FJ, Ekblom B, Sahlin K, Lundberg JO, Weitzberg E. Effects of dietary

428 nitrate on blood pressure in healthy volunteers. *N Engl J Med* 2006;355:2792–3.

429 [26] Butler AR, Feelisch M. Therapeutic uses of inorganic nitrite and nitrate: from the

430 past to the future. *Circulation* 2008;117:2151–9.

431 [27] Lundberg JO, Weitzberg E, Gladwin MT. The nitrate-nitrite-nitric oxide pathway

432 in physiology and therapeutics. *Nat Rev Drug Discov* 2008;7:156–67.

433 [28] L'Hirondel J. Nitrate and man: toxic, harmless or beneficial? Wallingford, UK:

434 CABI; 2002.

435 [29] Kelly J, Fulford J, Vanhatalo A, Blackwell JR, French O, Bailey SJ, et al. Effects of

436 short-term dietary nitrate supplementation on blood pressure, O₂ uptake kinetics, and

437 muscle and cognitive function in older adults. *Am J Physiol Regul Integr Comp*

438 *Physiol* 2013;304:R73–83.

439

440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466

Figure Legends

Figure 1: Participant flow through the trial

Figure 2: Urinary (Fig 1A) and salivary (Fig 1B) nitrate concentrations stratified by type of intervention (beetroot juice; blackcurrant juice). Nitrate concentrations were measured by gas-chromatography mass spectrometry. T-test for independent samples was used to compare the two groups. Mean values are reported. Error bars are 1SE.

Figure 3: Effects of beetroot (n=10) and blackcurrant juice (n=11) on resting (Figure 3A) and 24-hr ambulatory (ABPM) (Fig 3B) systolic and diastolic blood pressure (BP). Changes in BP were not significant between groups using both methods. Mean values are reported. Error bars are 1SE.

Figure 4: Effects of beetroot (n=10) and blackcurrant juice (n=11) on home daily monitored systolic and diastolic blood pressure (BP). BP values were averaged for each week and changes compared between groups at each week after adjustment for baseline values. At week 4, adjustment was performed for BP values measured on day 21. Weekly values in systolic and diastolic BP for the beetroot (BT) and blackcurrant (BC) interventions are:


Systolic BP ($\bar{X} \pm SD$): Baseline BT = 129.8±19.1mmHg, Baseline BC = 131.5±14.6mmHg; Week 1 BT = 127.6±16.6mmHg, Week 1 BC = 131.0±13.4mmHg; Week 2 BT = 126.6±14.7mmHg, Week 2 BC = 131.6±12.3mmHg; Week 3 BT = 125.2±16.1mmHg, Week 3 BC = 132.5±10.6mmHg; Week 4 BT = 128.6±15.8mmHg, Week 4 BC =


467 131.4±11.9mmHg. Diastolic BP ($\bar{X} \pm SD$): Baseline BT = 77.1±15.4mmHg, Baseline BC =
468 75.5±7.1mmHg; Week 1 BT = 76.0±12.4mmHg, Week 1 BC = 76.6±7.1mmHg; Week 2 BT
469 = 75.5±10.7mmHg, Week 2 BC = 76.4±7.3mmHg; Week 3 BT = 74.5 ±10.9mmHg, Week 3
470 BC = 76.3±7.4mmHg; Week 4 BT = 75.9±11.2mmHg, Week 4 BC = 76.1±6.7mmHg.

Table 1: Baseline characteristics of participants randomised to the two nutritional interventions (beetroot concentrate; blackcurrant juice)


| | Beetroot | Blackcurrant | p |
|-----------------------------------|------------|--------------|------|
| N | 10 | 11 | |
| M/F | 7/3 | 5/6 | 0.38 |
| Age (years) | 62.7±1.5 | 61.4±1.3 | 0.54 |
| Height (m) | 174.4±0.03 | 168.8±0.03 | 0.28 |
| Weight (kg) | 92.5±4.8 | 84.2±4.4 | 0.21 |
| BMI (kg/m ²) | 30.5±1.3 | 29.4±1.2 | 0.55 |
| WC (cm) | 103.8±4.0 | 100.7±3.0 | 0.55 |
| FM (kg) | 31.8±3.5 | 27.3±2.5 | 0.30 |
| FFM (kg) | 60.7±3.4 | 56.8±4.0 | 0.48 |
| Clinic SBP (mmHg) | 135.1±4.7 | 131.1±4.4 | 0.55 |
| Clinic DBP (mmHg) | 77.4±3.0 | 76.1±3.3 | 0.78 |
| Clinic HR (bpm) | 60.0±2.1 | 66.1±1.8 | 0.06 |
| 24-hour ABPM SBP (mmHg) | 125.8±5.2 | 130.9±3.9 | 0.44 |
| 24-hour ABPM DBP (mmHg) | 81.0±3.8 | 82.0±2.2 | 0.83 |
| 24-hour ABPM HR (bpm) | 65.9±1.7 | 74.4±3.0 | 0.03 |
| HM SBP (mmHg) | 129.8±6.1 | 131.6±4.4 | 0.61 |
| HM DBP (mmHg) | 77.2±4.9 | 75.6±2.3 | 0.68 |
| HM HR (bpm) | 61.5±2.4 | 66±1.9 | 0.08 |
| Urinary NO ₃ (mmol/L) | 0.95±0.2 | 0.68±0.09 | 0.34 |
| Salivary NO ₃ (mmol/L) | 0.85±0.23 | 0.97±0.30 | 0.76 |

Data are presented as mean±SD; N= number of subjects; M/F= male/female; BMI= body mass index; WC= waist circumference; FM= fat mass; FFM= fat free mass; SBP= systolic blood pressure; DBP= diastolic blood pressure; HR= heart rate; ABPM= ambulatory blood pressure monitoring; HM= home monitoring; NO₃= nitrate. T test for independent samples was used to compare the two groups.





A



B

