

# Northumbria Research Link

Citation: Hill, Matthew J., Greaves, Helen M., Sayer, Carl, Hassall, Christopher, Milin, Melanie, Milner, Victoria S., Marazzi, Luca, Hall, Ruth, Harper, Lynsey R., Thornhill, Ian, Walton, Richard, Biggs, Jeremy, Ewald, Naomi, Law, Alan, Willby, Nigel, White, James C., Briers, Robert A., Mathers, Kate L., Jeffries, Michael J. and Wood, Paul J. (2021) Pond ecology and conservation: research priorities and knowledge gaps. *Ecosphere*, 12 (12). e03853. ISSN 2150-8925

Published by: Ecological Society of America

URL: <https://doi.org/10.1002/ecs2.3853> <<https://doi.org/10.1002/ecs2.3853>>

This version was downloaded from Northumbria Research Link:  
<http://nrl.northumbria.ac.uk/id/eprint/46856/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



**Northumbria  
University**  
NEWCASTLE



**UniversityLibrary**

# **Pond ecology and conservation: research priorities and knowledge gaps**

Running head: Pond ecology and conservation

Type of paper: Innovative Viewpoints

Matthew J Hill<sup>1</sup>, Helen M Greaves<sup>2</sup>, Carl D Sayer<sup>2</sup>, Christopher Hassall<sup>3</sup>, Mélanie Milin<sup>1</sup>, Victoria S Milner<sup>1</sup>, Luca Marazzi<sup>4</sup>, Ruth Hall<sup>5</sup>, Lynsey R Harper<sup>6</sup>, Ian Thornhill<sup>7</sup>, Richard Walton<sup>8</sup>, Jeremy Biggs<sup>9</sup>, Naomi Ewald<sup>9</sup>, Alan Law<sup>10</sup>, Nigel Willby<sup>10</sup>, James C White<sup>11</sup>, Robert A Briers<sup>12</sup>, Kate L Mathers<sup>13,14</sup>, Michael J Jeffries<sup>15</sup>, and Paul J Wood<sup>14</sup>.

<sup>1</sup>School of Applied Sciences, University of Huddersfield, Queensgate, Huddersfield, HD1 3DH, UK.

Email: [m.hill@hud.ac.uk](mailto:m.hill@hud.ac.uk)

<sup>2</sup>Pond Restoration Group, Environmental Change Research Centre, Department of Geography, University College London, Gower Street, London WC1E 6BT, UK.

<sup>3</sup>School of Biology, Faculty of Biological Sciences, University of Leeds, Woodhouse Lane, Leeds, LS2 9JT, UK.

<sup>4</sup>Institute of Environment, Florida International University, Miami, FL 33199, USA.

<sup>5</sup>Natural England Mail Hub, Natural England, Worcester County Hall, Spetchley Road, Worcester, WR5 2NP, UK.

<sup>6</sup>School of Biological and Environmental Sciences, Liverpool John Moores University, Liverpool, L3 3AF, UK.

<sup>7</sup>School of Sciences, Bath Spa University, Newton St. Loe, Bath, BA2 9BN, UK.

<sup>8</sup>School of Geography, Politics and Sociology Newcastle University, King's Gate, Newcastle upon Tyne, NE1 7RU, UK.

<sup>9</sup>Freshwater Habitats Trust, Bury Knowle House, Headington, Oxford, OX3 9HY, UK.

<sup>10</sup>Biological and Environmental Sciences, University of Stirling, Stirling, FK9 4LA, UK.

<sup>11</sup>River Restoration Centre, Cranfield University, Cranfield, Bedfordshire MK43 0AL, UK.

25   <sup>12</sup>School of Applied Sciences, Edinburgh Napier University, Edinburgh, EH11 4BN, UK.

26   <sup>13</sup>Eawag, Department of Surface Waters Research and Management, 6047 Kastanienbaum,  
27   Switzerland.

28   <sup>14</sup>Centre for Hydrological and Ecosystem Science, Department of Geography, Loughborough  
29   University, Loughborough, Leicestershire, LE11 3TU, UK.

30   <sup>15</sup>Department of Geography and Environmental Sciences, Northumbria University, Newcastle upon  
31   Tyne, NE1 8ST, UK.

32

33   **Author for correspondence**

34   Matthew J Hill

35   School of Applied Science

36   University of Huddersfield

37   Queensgate

38   Huddersfield

39   HD1 3DH, UK.

40   Email: [m.hill@hud.ac.uk](mailto:m.hill@hud.ac.uk)

## **Abstract**

Ponds are amongst the most biodiverse and ecologically important freshwater habitats globally and may provide a significant opportunity to mitigate anthropogenic pressures and reverse the decline of aquatic biodiversity. Ponds also provide important contributions to society through the provision of ecosystem services. Despite the ecological and societal importance of ponds, freshwater research, policy and conservation have historically focussed on larger waterbodies, with significant gaps remaining in our understanding and conservation of pond ecosystems. In May 2019, pond researchers and practitioners participated in a workshop to tackle several pond ecology, conservation and management issues. Nine research themes and 30 research questions were identified during and following the workshop to address knowledge gaps around: (i) pond habitat definition; (ii) global and long-term data availability; (iii) anthropogenic stressors; (iv) pond monitoring and technological advances; (v) aquatic-terrestrial interactions; (vi) succession and disturbance; (vii) freshwater connectivity; (viii) socio-economic factors; and (ix) conservation, management and policy. Key areas for the future inclusion of ponds in environmental and conservation policy were also discussed. Addressing gaps in our fundamental understanding of pond ecosystems will facilitate more effective research-led conservation and management of ponds, their inclusion in environmental policy, the provision of ecosystem services, and help address many of the global threats driving the decline in freshwater biodiversity.

## **Keywords**

Aquatic-terrestrial linkages, biodiversity, connectivity, ecosystem services, management, policy, small lentic waterbodies

## Introduction

Biodiversity is declining worldwide with significant reductions particularly in freshwater fauna and flora (Grooten and Almond 2018). With anthropogenic pressures such as agricultural intensification, urbanisation, hydrological alterations, habitat fragmentation, climate change and pollution likely to continue and intensify, biodiversity loss and species extinctions within freshwater environments are expected to increase in the future (Jantz et al. 2015). Ponds, defined here as lentic waterbodies between 1 m<sup>2</sup> and <2 ha in area (Williams et al. 2010), and pondsapes (a network of ponds and their surrounding terrestrial matrix) are a significant freshwater resource in the landscape. There are an estimated 547 million – 3.19 billion ponds globally, although significant uncertainty surrounds pond numbers at national and international scales in many regions (Holgerson and Raymond 2016). As such, the management and conservation of ponds offer considerable opportunities to mitigate some threats faced by freshwater habitats and reverse the decline of freshwater biodiversity globally.

Cumulatively, ponds often support a greater biodiversity than other freshwater habitats (e.g., lakes and rivers), sustain many rare and endangered aquatic taxa, and act as important ‘refuges’ in heavily modified landscapes (Davies et al. 2008). Alongside aquatic species, many terrestrial species, including insect pollinators, birds, bats and other mammals rely on ponds for water, food and habitat (e.g., Nummi et al. 2011; Lewis-Phillips et al. 2020). Ponds also play a significant role in the provision of ecosystem services to society including water purification, flood alleviation, irrigation, watering livestock, fish production, support for pollinators and climate change mitigation (Pereira Souza et al. 2019; Lundy and Wade, 2011; Coutts et al. 2012; Stewart et al. 2017; Vico et al. 2020). Ponds also have significant amenity and educational value (Bastien, Arthur and McLoughlin, 2012), and can be used to raise awareness of biodiversity and nature conservation as well as providing a space for physical activity and relaxation (Higgins et al. 2019). Despite their biodiversity and societal value, ponds have historically been dismissed as unimportant and have been drained and removed as a result of changes in agricultural practices and urban development (Wood et al. 2003), or because they are seen as breeding grounds for disease vectors (e.g., mosquitos; Dos Reis et al. 2015).

90

91 In recent decades, there has been a growing interest amongst scientific and non-scientific  
92 communities in pond biodiversity, their conservation and contribution to ecosystem services (Biggs et  
93 al. 2017), reflected in the rapid rise of publications between 2000 and 2019 (Figure 1). Despite this  
94 growing recognition of the ecological and societal importance of ponds, freshwater scientific research,  
95 policy and conservation remains disproportionately focussed on rivers and larger lakes (Figure 1),  
96 leaving significant gaps in our understanding of these valuable ecosystems. In addition, ponds are  
97 threatened by anthropogenic activities but remain a low priority of national and international  
98 conservation and environmental legislation in most countries (Hill et al. 2018). International  
99 collaboration and commitment is urgently needed to increase fundamental understanding of pond  
100 ecology, develop more effective practical conservation and management strategies, and for the  
101 implementation of detailed national and international conservation policies (Hill et al. 2018).

102

103 In May 2019, leading pond researchers, regulators and practitioners across the United Kingdom  
104 (including Natural England, the Environment Agency and the Freshwater Habitats Trust) came  
105 together at a workshop (*Pond Ecology and Conservation in the Anthropocene*), to discuss the current  
106 status and future directions of key pond ecology, conservation and management issues. The  
107 workshops objectives were to: (i) identify critical knowledge gaps where research is required to  
108 advance understanding of pond ecosystems; (ii) develop and / or facilitate more effective conservation  
109 and management strategies; and (iii) establish the required evidence base to support the future  
110 inclusion of pond ecosystems in wider environmental and conservation policy initiatives. Initially,  
111 attendees individually and independently listed the ‘key challenges’ and ‘research priorities’ for pond  
112 ecosystems, based on theoretical (e.g., fundamental understanding) and applied (e.g., management,  
113 conservation and wider societal functions) topics. Detailed group discussions followed, which reduced  
114 individual lists (over >30 research priorities in total) to nine research themes: (i) definition of pond  
115 habitats; (ii) global and long-term data; (iii) anthropogenic stressors; (iv); aquatic-terrestrial  
116 interactions; (v) succession and disturbance; (vi) freshwater connectivity; (vii) pond monitoring and

technological advances (viii) socio-economic factors; and (xi) conservation management and policy (Figure 2; Table 1). Subsequently, workshop organisers collated research questions from attendees, based on the research themes, and refined them into 30 questions for future pond research, building upon previous studies by Calhoun et al. (2017) and Biggs et al. (2017). The research themes and questions are outlined below by theme and are not ordered by importance or relevance.

## **Research Priorities**

### **Definition of pond habitats**

Since the 19<sup>th</sup> century there have been numerous attempts to define ponds (see Appendix 1 in Biggs et al. 2005). The distinction between smaller lakes and ponds has proved particularly difficult, as well as between ponds and virtually all other small standing freshwaters into which they merge (Biggs and Williams 2020), although several factors can be used to distinguish larger lakes from smaller lakes and ponds (e.g., littoral zone size, catchment size, wind protection, and environmental heterogeneity; Sondergaard, et al. 2005). Despite this, distinguishing between small lakes and large ponds can still be problematic as they share many characteristics (Biggs et al. 2017). This discrepancy in definition is likely to have implications for our understanding of ecological processes in small waterbodies, and the development of pond management and conservation strategies.

In operational terms, ponds have been defined based on their size as waterbodies between 1 m<sup>2</sup> and 2 ha in the UK (Williams et al. 2010), although the upper size limit used to define a pond varies considerably among countries. Values between 1 ha and 6 ha are often employed (Ilg and Oertli 2017), and the Ramsar Convention on Wetlands proposed an area of 8 ha to distinguish between ponds and lakes (Ramsar Convention Secretariat 2013). To provide a consistent, scientifically derived definition, research is required to examine the significance of depth, wind action, nutrient transport and light penetration across a range of waterbody sizes, as these factors may have a large influence on biological communities. Recent research has found that small waterbodies may experience strong

diurnal stratification and mixing (due to convection), and seasonal stratification (Sayer et al. 2013), a process which can influence pond environmental conditions, patterns in freshwater biodiversity and the performance of organisms, particularly sessile taxa (Anderson et al. 2016; Martinsen et al. 2019). A threshold in water body size where there is a change in mixing processes may provide a suitable characteristic to distinguish between ponds and lakes, although this currently remains inadequately quantified. The wide range of pond definitions partly reflects the wide range of perennial and temporary pond types that exist internationally, all of which demonstrate significant variability in environmental conditions and origin (Biggs et al. 2017). Given the complexity of this task, simple size-based definitions will probably continue to play an important role in practical identification and management of small standing waters. But clearly an overarching, process-based definition of a pond is required to allow researchers to undertake more targeted and comparable research, and for practitioners to develop effective conservation and management strategies and policies.

#### *Key research question*

- Are there critical thresholds in physical, biological and chemical processes, and ecosystem function, which reflect changing lentic waterbody size?

#### **Global and long-term data**

While global-scale research has been undertaken for lotic (e.g., Tiegs et al. 2019) and lake ecosystems (e.g., Alahuhta et al. 2017), there has been no equivalent research examining global- and international-scale patterns in pond diversity and functioning, despite the threats facing freshwater habitats worldwide (but see Downing et al. 2008). Some studies have examined pond communities at a national-scale (e.g., Hill et al. 2017), but it is not clear if similar ecological patterns and processes occur at continental or global-scales. To enable more effective international management and conservation of the biodiversity and associated ecosystem services of ponds, there is a need for the systematic consolidation of existing datasets to examine global-scale biodiversity patterns and the



community assembly processes at multiple spatial scales. Key to these studies will be placing ponds in the context of other freshwater habitats (e.g., lakes, rivers) across the aquatic landscape (Sayer 2014). The development of an open access, global pond database of biotic and environmental data will facilitate significant global-scale research and increase our understanding of these important waterbodies; although this database does not currently exist. New large-scale collaborative studies will be critical to understand how global pressures (e.g., land-use change and pollution) are affecting pond diversity (e.g., total biodiversity and rare species distributions) and ecosystem service provision (e.g., stormwater collection, water purification and human wellbeing).

A further scaling issue that inhibits conservation efforts and the development of effective policy for ponds is the lack of long-term datasets. With a few exceptions (see Williams et al. 2020; Jeffries 2011), most pond research has been dominated by studies covering a single season or year (Hill et al. 2016), with studies covering timescales longer than 10 years largely missing. Consequently, there are significant knowledge gaps regarding the long-term dynamics and patterns in pond environments, biological diversity and functioning. As for larger lentic waterbodies, palaeoecological approaches could provide critical information for inferring long-term (c.100s-1000s years) changes in pond ecology in response to environmental change, and, in turn, inform restoration strategies (Walton et al. 2021). This is of particular importance for demonstrating the scope for returning communities to those recorded prior to major human impacts. As well as encouraging long-term data collection, the comparison of contemporary samples with older records (Hassall et al. 2012) and mesocosm experimentation may elucidate long-term processes (Yvon-Durocher et al. 2017).

#### *Key research questions*

- What are the abiotic and biotic drivers of pond environmental and biological patterns at different spatial and temporal scales?

- How do biological communities within ponds respond to global environmental change along spatial and temporal gradients?

## **Anthropogenic stressors**

Ponds are increasingly being subjected to multiple anthropogenic stressors which can affect the diversity and resilience of their communities (Ryan et al. 2014). The cocktail of anthropogenic stressors may include chemical (e.g., pollution), biological (e.g., invasive species), and physical (e.g., infilling, management, and climatic warming) pressures. In addition, stressors may originate from anthropocentric perceptions of ponds, such as the high values placed upon neatness, carefully managed plant communities (removal of ‘weedy’ and aesthetically unpleasing species), and large areas of open water, perceptions which often do not reflect the natural functioning of ponds (Nassauer 2004). Many stressors remain poorly understood, and in this section, we focus on the impacts of pollution, non-native species and climate change on pond communities, and identify knowledge gaps and key areas for future research.

### *Pollution*

One of the most pervasive threats to pond communities is pollution. Nutrient enrichment, which is known to decrease species diversity at a landscape-scale (Rosset et al. 2014), is one of the better understood threats to pond communities in agricultural and heavily populated landscapes. In urban landscapes, run off from impermeable surfaces can result in an increase in heavy metals, nutrients, road salts and chloride in ponds (Moss 2017). However, many natural and anthropogenic ponds in urban landscapes are designed and managed to hold stormwater runoff and are thus intended to be polluted as a means to protect water quality in the catchment downstream (Gold et al. 2017). As such there is a paradox in urban landscapes, where ponds have high biodiversity value but are being designed to hold pollutants to improve down-stream water quality of more highly valued (but not actually more biodiversity rich) freshwater habitats, such as larger lakes and rivers. Given urban areas

are predicted to increase in size across the globe (Seto et al. 2012), more ponds are likely to experience pollution, but the impacts on urban pond biodiversity are not fully understood (Hintz and Relyea 2019). Higher numbers of polluted ponds also poses potential risks from increasing the number of water bodies which are producing large quantities of climate heating gases (Peacock et al. 2019; Rosentreter et al. 2021).

The process of suburbanization also poses a major threat to freshwater biodiversity (Van Acker et al. 2019). Pond habitats frequently survive the conversion from natural to suburban land cover, but they are often subject to wastewater runoff (chemical loading), physical changes (e.g., decreased canopy cover and increased temperature) and isolation which can have large effects on food web dynamics, species health and taxa richness (Van Acker et al. 2019; Holgerson et al. 2018; Homan et al. 2004). Pesticide pollution in freshwaters also represents a long-term persistent issue (e.g., Ito et al. 2020), yet quantifying concentrations remains challenging due to high spatiotemporal variability, limitations of instrument availability and difficulty in identifying new / emerging chemicals (Lorenz et al. 2017). In particular, neonicotinoid pesticides (Raby et al. 2018) are contaminants of freshwaters which have been given little attention in pond environments thus far. Moreover, despite a large body of literature in lotic systems (Calabrese et al. 2020), there has been little research examining interactive effects of multiple stressors on pond communities, which could be additive, subtractive, antagonistic or synergistic (Hassall 2014).

#### *Non-native species*

Non-native species are one of the greatest threats to many freshwater systems, but are probably less common in ponds than other highly connected freshwater environments associated with their hydrological isolation (Williams et al. 2010), although ponds in urban areas are highly vulnerable to intentional introductions (Patoka, et al. 2017).

Where non-native species become highly invasive after successfully colonising and establishing in ponds, impacts are invariably detrimental. For example, the effects of New Zealand Pigmywort (*Crassula helmsii*) on pond communities in Europe are widely acknowledged (e.g., loss of plant biomass and abundance, plant germination suppression; Langdon et al. 2004; Smith and Buckley 2020) and the introduction of the Topmouth Gudgeon (*Pseudorasbora parva*) into European pond habitats has been shown to shift the trophic position of fishes (Britton et al. 2010). However, there are a wide range of non-native invasive species that are expanding their habitat range and of which we know little about their impacts for pond ecosystem functioning and biodiversity (e.g., *Dikerogammarus villosus* and *Pacifastacus leniusculus*).

While it is clear that the establishment of non-native invasive species can be detrimental to native communities, not all introductions of non-native taxa are negative. For example, in Oregon, USA, the non-native Reed Canary grass (*Phalaris arundinacea*) has been beneficial to breeding amphibians in many ponds, providing suitable sites for oviposition, increasing tadpole survival and the abundance of adult amphibians (Holzer and Lawler 2015) and in Washington State, USA Holgerson et al. (2019) found that non-native bullfrogs (*Lithobates catesbeianus*) did not impact pond amphibian occupancy (but non-native fish did). Should ponds be less widely affected than other freshwaters they may represent refugia for some taxa otherwise threatened by non-native invasive species in shallow lakes and rivers (Sayer et al. 2011). Historically, the effects of non-native species on trophic interactions within ponds have been poorly studied compared to their lotic counterparts and as such further consideration is required to identify the implications of predator-prey interactions between non-native and native species, as well as how non-native species may be driving shifts in pond ecosystem functioning. Research is clearly required to examine the full range of ecological effects (negative, positive and negligible) that non-native taxa may have on native pond communities, and how they can be managed to facilitate conservation of native species.

*Climate change*

As small but abundant habitats, ponds are sensitive to the effects of environmental change and are therefore essential to consider within climate feedbacks. Agricultural impoundment ponds are among the most significant habitats for carbon sequestration due to their high nutrient loads and associated high primary productivity (Downing et al. 2008). Even accounting for their relatively small size, research has demonstrated that carbon burial rates in small ponds are at least equivalent to woodland or grassland habitats (Taylor et al. 2019). However, other studies have suggested that small waterbodies can switch between carbon sources and sinks, while boreal ponds and small agricultural reservoirs may even act as significant sources of greenhouse gas emissions (Holgerson and Raymond 2016; Webb et al. 2019). Given recent drives towards afforestation for climate mitigation (Brown 2020), determining the net value of ponds within the carbon cycle may reveal their potential, helping diversify climate mitigation beyond forestry alone.

Understanding the different dimensions of climate change – gradual warming, heat waves, droughts, and floods – and how they interact and affect pond communities is vital to the future conservation of pond environments. Extreme climatic events may result in homogenised pond communities, reducing the biodiversity value of pondscapes (Bertoncin, et al. 2019). Drought is likely to increase the proportion of ponds that experience hydrological intermittency resulting in changing community structures of existing temporary ponds, extirpation of species unable to adapt to increasing ephemerality (Ryan et al. 2014; Abney et al. 2019), and potentially influence greenhouse gas emissions from ponds (Holgerson and Raymond 2016). These climate shifts typically favour freshwater communities from warmer regions that can take advantage of higher temperatures to circumvent the increasingly ephemeral nature of the ponds. However, a better understanding of the interaction between short- and long-term climate effects on ponds to predict future effects is required.

Climate change has been shown to interact with other stressors for example by exacerbating the effects of pesticides (Janssens and Stoks, 2013) and sedimentation (Piggott et al. 2012) due to climatic changes to rainfall and runoff regimes altering the delivery of fine sediment and its associated

contaminants. In contrast, some studies suggest that climate change may be antagonistic to biological invasions, where non-native species increase their functional capacity to the same level as native species under warming (Kenna et al. 2017), but climate change has also been found to intensify the effects of non-native introductions (Bellard et al. 2012). While some progress has been made in understanding how co-occurring stressors interact (if at all), a synthetic approach to gathering evidence across spatial and temporal scales could yield further insights across ponds and the wider freshwater network and inform new research.

Climate change also potentially mediates interactions among species through driving spatio-temporal patterns of occurrence. While phenological change has been described for some freshwater taxa (Hassall 2015), the ecological and fitness implications of phenological changes in pond communities have received little attention. The existing focus has largely been on birds (Radchuk et al. 2019), and changes in synchrony of interacting taxa have mostly been studied in larger waterbodies (e.g., Winder and Schindler 2004). Given the importance of links between aquatic and terrestrial food webs (Stenroth et al. 2015), more research examining the effect of climate change on aquatic-terrestrial networks and the flow of nutrients and services between these habitats is required (Soininen et al. 2015).

#### *Key research questions*

- What are the interactive effects of multiple-stressors on pond ecosystem resilience?
- How do pollutants impact pond ecological functioning and resilience?
- How widely are ponds affected by non-native species and what are the dynamics of their spread?
- How do non-native species introductions affect trophic interactions in pond ecosystems?
- What are the potential ecological implications for pond communities associated with different climate change scenarios?

- How will climate change affect the interactions between aquatic and terrestrial food webs?

### **Aquatic-terrestrial interactions**

The importance of freshwater habitats is not limited to the waterbody itself. Interactions between aquatic and terrestrial ecosystems can be significant, demonstrated by the importance of terrestrial vegetation shading in structuring freshwater invertebrate assemblages (Suh and Samways 2005), the negative effects of land-use change on terrestrial insectivores reliant on aquatic insects, and the effect of predation (e.g., dragonflies preying on bees) on aquatic-terrestrial interactions (Knight et al. 2005; Stenroth et al. 2015). There is a growing recognition of the importance of interactions between ponds and terrestrial species, especially farmland birds (Lewis-Phillips et al. 2020) and insect pollinators (Stewart et al. 2017). The reliance of terrestrial organisms on ponds may be largely due to trophic interactions that bridge the aquatic-terrestrial divide. Farmland ponds managed by scrub and sediment removal (to re-establish macrophyte-dominance) have been shown to increase avian diversity and abundance due to enhanced emerging invertebrate food sources (Lewis-Phillips et al. 2020) and support distinct diurnal pollinator communities (Walton et al. 2020). An increased abundance of local pollinators enhances pollination services in neighbouring crop fields (Stewart et al. 2017), while increased numbers of spiders and beetles adjacent to some ponds have been linked to emerging insect prey from ponds (McCafferey and Eby 2016) and might promote natural pest control over wider scales.

Of particular note, ponds can act as important components of the food web and energy flow between aquatic and terrestrial organisms. For example, birds feeding on aquatic invertebrates have crucial access to higher levels of unsaturated omega-3 fatty acids needed for improved physical and cognitive development than birds relying primarily on terrestrial invertebrates (Twining et al. 2016). However, it is unclear how pollution and other anthropogenic stressors in ponds may influence the nutritional value of aquatic macroinvertebrates and therefore their value to terrestrial predators. Conversely,

nutrients that have moved from aquatic to terrestrial systems can eventually be transported back when semi-terrestrial organisms such as amphibians return to ponds to lay eggs (Regehr et al. 2006; Capps et al. 2015). Amphibian larvae often also form part of pond fish diets which can act as a limiting factor on amphibian richness and abundance within ponds (Hecnar and M'Closkey 1997; Hartel et al. 2007). Furthermore, terrestrial leaf litter and other detritus that falls into ponds provides nutrients (Fey et al. 2015) as well as food for invertebrate communities (Holgerson et al. 2016). These examples highlight the significance of ponds as food and energy links between the aquatic and terrestrial realms.

Despite this emerging knowledge, there is still a lack of understanding regarding the role that ponds play in aquatic-terrestrial interactions. For example, bats and other small mammals may benefit from pond resources including emerging aquatic insects and the provision of drinking water (Nummi et al. 2011), while vertebrate herbivores, including waterfowl, beaver, elk and moose are attracted by shallow water and extensive palatable vegetation (Law et al. 2014). Furthermore, terrestrial vertebrate visitors to ponds may provide important ecosystem processes, including trampling, grazing and poaching which could arrest successional processes, stimulate seedbank emergence, disperse propagules (Cieminski and Flake 1997), and enhance fine-scale spatial heterogeneity (Willby et al. 2019). Such dynamics are considered critical in maintaining populations of many endangered plant species. More research on pond aquatic-terrestrial interactions and food web structures that transcends traditional aquatic-terrestrial boundaries is clearly needed.

#### *Key research questions*

- Does the nutritional value of aquatic macroinvertebrates for terrestrial predators reflect pond water quality?
- How important are terrestrial vertebrates for the dispersal of pond biota and for maintaining local environmental heterogeneity and does this vary systematically with pond setting?



- How do aquatic–terrestrial interactions in predation influence pond community structure and vice versa?

### **Succession and disturbance**

Due to their small size and shallow depth, hydrosere succession may operate faster in ponds compared to larger waterbodies. However, little is known regarding the timescales or pathways of succession across ponds, particularly in relation to their origin, climatic and geological settings (Biggs and Williams 2020). Disturbance, both natural and anthropogenically-driven, is an important influence on succession, but remains poorly studied among pond habitats. Natural disturbances, including beaver damming, natural tree fall, wetting and drying cycles, floods (resulting in meander cut-off and channel avulsion), and the activities of large herbivores (both domestic and wild), may act to create ponds, or delay and reset succession trajectories. Equally, human-induced disturbances can create new ponds, for example, war time bomb craters supporting high diversity ponds in Central Europe (e.g., Vad et al. 2017). In ponds where natural disturbances occur, it is generally accepted that ponds will be at different stages of succession, which may enhance both beta and gamma diversity (Hill et al. 2017). In contrast, in landscapes where natural disturbances have been reduced, uninterrupted succession can result in areas entirely dominated by late-succession ponds, thus reducing biodiversity (Sayer et al. 2012). The influence of succession and disturbance regimes (natural and anthropogenic) on landscape-scale species distribution and diversity patterns remains poorly understood, in part due to a lack of long-term monitoring, but could provide critical information for the development of landscape-scale conservation and management strategies.

Many pond species possess life cycles and adaptive strategies enabling them to persist across the landscape in the face of disturbance, such as dormant and resistant propagules of aquatic invertebrates and plants (Alderton et al. 2016; Williams et al. 1997), and relatively long lifespans for amphibians and some invertebrates. Efficient dispersal mechanisms by many invertebrates allow rapid recolonization when favourable conditions return including endozoochory (Kleyheeg and van

Leeuwen 2015), aerial dispersal (Bilton et al. 2001) and ‘hitch-hiking’ (Okamura et al. 2019). Research focused on the life history strategies that enable species and communities to persist after a disturbance will be especially valuable to conservation managers as manipulating and facilitating disturbance via active management or passively by rewilding may be key to the survival and protection of many pond species.

#### *Key research questions*

- What role do natural disturbances play in structuring aquatic and terrestrial biodiversity in pond habitats across landscapes minimally impacted by human influences?
- What are the best management strategies to reduce detrimental disturbances and increase positive disturbance in anthropogenically dominated landscapes?
- What are the principal trajectories, timescales and outcomes of pond succession across different pond types?

#### **Freshwater connectivity**

Connectivity is important because it facilitates the movement of energy, materials, organisms and genetic resources within and between habitats in a landscape or more widely. Published studies examining connectivity have accelerated greatly during the 21st century alongside the growing debate on the relative roles of dispersal limitation and local species sorting across all ecosystem types (Heino et al. 2017). A growing number of studies have reported the importance of spatial factors (Juračka et al. 2019), or biological traits linked to dispersal ability (De Bie et al. 2012) in determining assemblage structure, indicating that dispersal limitation is likely to be an important influence on pond communities. Contrasting active and passive dispersers, both within and across taxon groups, has confirmed expectations that dispersal limitation among ponds is stronger among less mobile taxa (De Bie et al. 2012; Hill et al. 2017a). The transferability of these patterns across a range of landscape types and pond characteristics now needs to be determined as most studies on connectivity in

freshwater systems focus on longitudinal connectivity in rivers and tend to examine freshwater habitats in isolation. In the case of ponds, two-way connectivity both across the aquatic-terrestrial interface and with other freshwater habitats in the wider landscape are likely to be especially important.

Connectivity is complex to quantify and is typically estimated using simple indicators, including Euclidean distances to similar habitats (Juračka et al. 2019), hydrological pathway lengths or percentage of surrounding freshwater with a buffer zone (Law et al. 2019), waterbird migration flyways (Viana, et al. 2016) and human population densities or proximity to recreational facilities (Chapman et al. 2020). Measures of geneflow or similarity in assemblage composition are often used to infer realised connectivity (Bilton et al. 2001). Metrics based on the structural properties of spatial networks (e.g., centrality and percolation thresholds) or species-specific dispersal distances and costs to crossing different habitats, have also been suggested as possible solutions (Thornhill et al. 2018; Hunter-Ayad and Hassall 2020). Rapid increases in the quality and availability of national biological recording datasets are improving understanding of the role of connectivity on the large-scale distribution of freshwater taxa. However, among ponds, the lack of high-resolution mapping may confound such attempts. It is also important to acknowledge that connectivity is not a static property and varies temporally for example, with hydroperiod or hydrological events, or seasonality linked to water bird migration, and can be fundamentally altered by anthropogenic activities or ecosystem engineers (Bilton et al. 2001). Moreover, rare or long-distance colonisation events may be important for connectivity (Jordano 2017), but accurately modelling or predicting these events will be complex when related to local physicochemical variables. Freshwater habitats are not discrete entities but often exist in networks (e.g., an interconnected system of ponds, rivers, streams and lakes; Sayer 2014), and ubiquitous species are recorded across these freshwater habitats (Davies et al. 2008). As such, it will be misleading to study ponds in isolation and more holistic research is required to understand the ecological processes operating across a range of connected freshwater habitats (the ‘waterscape’),

which will likely provide more accurate information for future landscape-scale conservation initiatives (Heino et al. 2021).

Management strategies to increase habitat patch connectedness is a common response to fragmentation however, the relative isolation of ponds may be an asset in insulating them from common stressors such as diffuse nutrient loading, pathogens or invasive species that are easily transmitted between more connected water body types. Understanding the aspects of connectivity that confer an advantage in terms of enhanced resilience, and those where it represents a risk in terms of accelerated transfer of stressors, is therefore critical in terms of future restoration or the design of pondscapes and wider waterscape.

#### *Key research questions*

- How does connectivity between pond habitats influence trophic interactions that bridge aquatic-terrestrial divides?
- How do species move between ponds and other freshwaters and what are the dominant mechanisms?
- How are pond networks (pondscapes) best designed or managed to ensure that rarer and less mobile species benefit?
- What is the role of spatial processes in assemblage structure and does pond context or landscape create regional differences?

#### **Pond monitoring and technological advances**

Recent developments in pond monitoring techniques and biostatistics can advance our understanding of the ecology and conservation of pondscapes. There are several challenges facing pond monitoring including; (1) often being located difficult to access and remote landscape settings, (2) identifying representative sites due to their high abundance, (3) their high environmental heterogeneity, resulting

in multiple ponds needing to be monitored to capture abiotic and biotic diversity, and (4) the availability of taxonomic specialists to identify the highly diverse floral and faunal taxa recorded within ponds. However, the utilisation of new technologies may help to overcome some of these challenges. In this section, we first outline the contribution that molecular tools have made to pond monitoring (Biggs et al. 2015; Deiner et al. 2017), and then discuss the opportunities that recent technological advances in remote sensing, un-manned aerial vehicles and biostatistics have provided for pond ecology and conservation.

#### *Pond monitoring using molecular tools*

The emergence of environmental DNA (eDNA) analysis has the potential to transform freshwater biodiversity assessment. eDNA is genetic material released by organisms into their environment, which can be sampled and analysed to target specific species or passively screen entire communities (Harper et al. 2019a). Targeted eDNA analysis can be used to assess the distribution and range of threatened, rare or non-native pond species (Biggs et al. 2015; Mauvisseau et al. 2018), and estimate relative abundance or biomass, and detection probability (Buxton, et al. 2017). Similarly, eDNA metabarcoding could be employed to assess multi-species distribution, reveal species interactions (Harper et al. 2019b), and characterise genetic diversity (Parsons et al. 2018), all of which are only beginning to be considered for pond ecosystems. Community DNA is distinct from eDNA samples, being sourced from biological material such as invertebrate blood meals (invertebrate-derived DNA: iDNA), faeces, and collected specimens (Deiner et al. 2017). iDNA analysis uses DNA that was ingested by invertebrates, such as leeches, to detect biodiversity within freshwater habitats (Abrams et al. 2019). iDNA metabarcoding could identify vertebrate biodiversity and enable multi-species occupancy modelling (Abrams et al. 2019) while faecal metabarcoding could be used to assess diets of threatened or invasive species and construct pond food-webs (Kaunisto et al. 2017). Although bulk tissue DNA and eDNA metabarcoding have been used for macroinvertebrate assessment in other freshwater ecosystems (Elbrecht et al. 2017), these techniques have rarely been applied to ponds. Using bulk tissue DNA and eDNA metabarcoding in pond research may provide more holistic

estimates of alpha and beta-diversity (Harper et al. 2020). Similarly, stable isotope analysis can complement metabarcoding to determine trophic relationships among pond taxa (Compson et al. 2019). These tools could more accurately quantify target species distribution and ranges, and determine the interactive effects of anthropogenic stressors, such as invasive species on communities and food webs.

#### *Technological advances in remote sensing, unmanned aerial vehicles and biostatistics*

Conventional data collection methods for some environmental variables within ponds (e.g., visual estimation of macrophyte coverage) are typically subjective or at best semi-quantitative and / or lack sufficient detail of wider environmental conditions. Remote sensing has been widely used in water quality assessment and resource management within lakes and rivers (Gholizadeh et al. 2016). Remote sensing can accurately measure certain catchment characteristics (e.g., land cover, productivity), physical properties (e.g., surface area, turbidity) and biological properties (e.g., aquatic macrophyte coverage) of freshwater environments across large spatiotemporal scales in a cost-effective and standardised manner (Giardino et al. 2010). Remote sensing could provide an efficient means to collect large-scale environmental and spatial metadata (e.g., pond numbers, connectivity, pond spatial structure and physical barriers) for pond research and assistance in the development of effective monitoring strategies, particularly for remote ponds and pond networks (Rose et al. 2015). Recent research, using remote sensing, was able to determine the number (>1000 ponds) and distribution of ponds across the Greater Kuala Lumpur region (~2,950 km<sup>2</sup>) in Malaysia, and quantify particular environmental conditions of each pond including the surface area, shape, connectivity and surrounding land-use (Teo et al. In review). Despite this, it remains a largely unused tool in pond research. However, given the spatial resolution of remotely sensed data it is unclear how remote sensing could be used to record small (<10m<sup>2</sup>) or intermittent ponds (during the dry phase), and those located under forest canopy (Gallant 2015; Kissel et al. 2020).

In lotic settings, the recent use of small Unmanned Aerial Vehicles (UAV) and structure-from motion (SfM) photogrammetric processing has provided a significant improvement in the objectivity, accuracy and efficiency of physical habitat data collection (Woodget et al. 2015). The parallels between streams and ponds suggest that similar advancements could be made for surveying surface area (particularly of small ponds), substrate composition, aquatic and riparian vegetation structure, habitat complexity, pollution events and the spatial structure of ponds and pondscape. UAVs can generate fine spatial resolutions (<10cm/pixel; Lucieer et al. 2014) and as such, have the potential to significantly increase the accuracy and consistency of pondscape data, elucidating the processes impacting their ecology and providing detailed site information to underpin effective monitoring strategies. However, no empirical studies have thus far examined the use of UAVs to characterise pondscape.

Recent developments in biostatistics may also provide considerable advances to our understanding of pond ecology and their conservation. Research on the drivers of species richness and community composition among lentic systems has focussed almost exclusively on local environmental and spatial factors, and has largely ignored the potential influence of biotic interactions (Heino et al. 2015). Recently, new statistical analysis has enabled the influence of biotic interactions, environmental conditions and spatial factors to be considered together providing a more realistic understanding of the factors that govern the spatial and temporal patterns in pond biodiversity (Garcia-Giron et al. 2020). Similarly, Local Contributions to Beta Diversity (LCBD; Legendre 2014) analyses goes beyond traditional measures of beta-diversity (a single measure of dissimilarity across the landscape) and calculates the contribution to overall beta-diversity by individual sites (provides a measure of ecological uniqueness for individual sites; Heino and Grönroos 2017). LCBD may contribute to pond conservation by identifying ponds with high numbers of unique species (that may be missed by traditional conservation practices that focus on taxonomic richness), whose protection can increase the number of species that are conserved at a landscape-scale. Recent research by Hill et al. (2021), found 70%–97% of the regional species pool was protected when ecologically unique (sites with high

LCBD values) and high taxonomic diversity sites (sites with >50 taxa) were considered together compared to 54%–94% when only sites with high taxonomic diversity were considered. Applying these new statistical techniques to different ecological and geographical settings will facilitate a greater understanding of pond ecology and more effective and targeted conservation strategies.

#### *Key research questions*

- Can molecular tools be used to assess the distribution of conservation priority and invasive species as well as community diversity at the pondscape scale?
- Are molecular tools and remote sensing able to identify the effects of anthropogenic stressors and climate change on ponds and improve management strategies to mitigate stressors?
- Can UAV-based data collection record physicochemical and spatial characteristics of ponds and pondsapes more accurately than conventional data collection?
- How can the development of new statistical analyses in biodiversity assessment contribute to more effective pond conservation planning?

#### **Socio-economic factors**

Ponds form an intrinsic component of urban and rural landscapes and many were historically created for a range of purposes, including industrial processes such as mineral extraction, provision of food and water, irrigation, watering livestock and as ornamental features (Gledhill and James 2012). In much of the world, many of the historical purposes of ponds are now redundant and today ponds are often managed as amenity features (e.g., angling) or have been abandoned. Angling ponds have been demonstrated to support limited faunal diversity (Wood et al. 2001), but there remains a paucity of research considering social elements associated with angling ponds. Research is needed to assess the personal, social and educational contributions made by angling ponds, and to better understand the practical and emotional governance conflicts that exist between anglers and other users of ponds to increase opportunities for sustainable management (Arlinghaus 2005). Given that the origin of many



ponds is a by-product from industrial processes, their presence in rural areas is unwanted by some landowners (Wood et al. 2003). As a result, there are still numerous barriers to the creation of new ponds and the management of existing ponds, despite recent evidence of biodiversity gains (Williams et al. 2020). Studies of the human dimensions (including social, cultural, institutional, emotional, communicative, governance and lifestyle tolerance factors) of stakeholder engagement with pond creation and management are currently largely absent but are required to ensure the success of any local or landscape-scale pond initiatives.

There is emerging evidence of the importance of blue space for the health and wellbeing of individuals by promoting psychological restoration, and providing spaces for physical activity, recreation and social interaction (Gascon et al. 2017; Foley and Kistemann 2015). Blue space regeneration could also help foster a sense of civic pride and ownership (Higgins et al. 2019). However, existing research has primarily focussed on coastal areas or lotic ecosystems, and there remain considerable gaps in our knowledge of the contribution of ponds to human health and wellbeing (Foley and Kistemann 2015). The immersive benefits of ponds in terms of their contribution to physical health, imaginative, emotional and therapeutic aspects, and the range of meanings for individuals and groups of pond ecosystems all require a greater understanding. In addition, research is needed to explore the relationships between ponds, wellbeing, access and habitat quality, particularly in urban landscapes (Higgins et al. 2019).

With increased urban living, the incorporation of ponds as blue spaces into the aesthetic design of urban areas represents an opportunity to engage the public with freshwater ecosystems (de Bell et al. 2017). Online packs of pond dipping materials and citizen science pond initiatives can reconnect individuals with nature, whilst engaging non-professionals in scientific research and practical freshwater conservation (Dickinson et al. 2012). Ponds also provide opportunities for education as the presence of ponds within schools may provide an important resource for educational study and can encourage an initial interaction and familiarisation with ponds and wildlife from an early age (Braund

1997). Research that considers the complexities of engagement, the educational value of ponds, barriers to environmental education, and the use of pond ecosystems for the development of an environmental conscience in individuals and society will be particularly beneficial in addressing freshwater environmental and ecological degradation and increase opportunities for pond conservation.

#### *Key research questions*

- How do ponds contribute to human physical and mental health and wellbeing within urban and rural populations?
- What are the barriers (including social, cultural, institutional, emotional, communicative, and governance) in stakeholder-pond conservation interactions, and how might these be addressed?
- What are the short- and long-term effects of environmental education for pond conservation?

#### **Pond conservation, management and policy**

While lake and riverine habitats have overwhelmingly dominated historic freshwater policy, conservation, and management research, more recently there has been a growing focus on ponds. Research on strategies for pond creation, pond and pondscape design, and where best to locate new ponds to maximise aquatic conservation benefits has developed significantly (Thiere et al. 2009). In addition, there is a growing body of research and expertise centred on the successful restoration and management of ponds in a variety of landscape settings, in particular European farmland and semi-natural habitats (Sayer et al. 2012; Sayer and Greaves 2020), Mediterranean coastal plains (Sebastian-Gonzalez et al. 2014) and the prairies of Canada and the USA (Bortolotti et al. 2016). Pond creation and restoration studies are typically short-term in duration, and studies assessing the medium to long-term success of measures alongside natural dynamics are needed (Seabloom and van der Valk 2003). These studies will ultimately determine the need (or not) for subsequent policy and management

activities to maintain conservation benefits (Sayer et al. 2012). Further, research is also required on the mechanisms that affect pond creation and restoration success, covering key issues such as water quality, grazing regimes, hydrology, connectivity and invasive species. In this respect, studies that compare and combine pond restoration with creation at the landscape-scale will be important to inform pond conservation planning and prioritisation.

Policy on the conservation and management of ponds has generally suffered from the assumption that small waterbodies were not important due to their size (Biggs and Williams 2020). This long-standing assumption, noted since the 1950s, has generally led to the absence of small waterbodies from environmental and conservation policy globally (Hill et al. 2018). For example, the European Union's Water Framework Directive, although intended to protect 'all' waterbodies specifically excludes those less than 50 ha from monitoring schemes, thereby excluding millions of ponds (although some are specially selected for nature conservation under the EU Habitats Directive). Similarly, in North America and Asia, ponds are generally not directly considered in environmental policy or legislation despite the establishment of the Clean Water Rule clarifying freshwater protection under the Clean Water Act in the USA (Department of Army, Corps of Engineers and US Environmental Protection Agency 2015) and various national-scale environmental legislation across Asia (e.g., Environment Protection Act 1986 and the Wildlife Protection Act 1972 in India). However, the policy failure relating to small waterbodies is slowly being reversed as awareness of the importance of these waterbodies appears to be increasing. For example, the Intergovernmental Science Policy Platform on Biodiversity and Ecosystem Services (IPBES) included temporary and perennial ponds in its classification of freshwater habitats in 2018 (IPBES 2018).

The future inclusion of ponds in environmental and conservation policy should focus on four areas: (1) stopping further deterioration, especially of the most vulnerable high-quality sites. Although evidence is scarce, the quality of the most protected ponds continues to decline (Williams 2019); (2) protecting species and communities that are a special feature of ponds. Increasing knowledge of pond communities

and the distribution of freshwater species will refine our understanding of the role of ponds play in protecting endangered freshwater species and inform policy; (3) encouraging the creation of new clean water ponds (clean water ponds reflect pond water chemistry and biology that is typical for a given area in the absence of human activity; Williams et al. 2010) and ensuring the effective restoration and management of existing ponds to maximise opportunities for maintaining and increasing species diversity at the landscape-scale, and; (4) creating / restoring ponds for ecosystem service measures for land and water management. Emerging evidence suggests that this could be one of the most effective means of enhancing landscape-scale freshwater biodiversity, but to maintain their long-term contribution, management interventions may be required (Williams et al. 2020). However, it is important to recognise, however, that ponds for ecosystem service purposes such as sediment retention or to attenuate flood flows, may not always provide suitable conditions for biodiversity provision (Williams et al. 2020).

#### *Key research questions*

- Where are the most ecologically important ponds at national and international scales?
- How does pond creation and management affect biotic communities at local and landscape-scales in different environmental settings?
- How can we best conserve pondscapes in anthropogenically dominated landscapes?
- What are the mechanisms required to better incorporate ponds into national and international environmental policy and legislation?

#### **Conclusion**

Large knowledge gaps remain in our understanding of pond ecosystems. Yet, it is clear that ponds can benefit society and wildlife by providing habitats that support significant freshwater and terrestrial biodiversity across a range of landscapes, whilst also providing ecosystem services required by society. Although ponds have received less research attention than other freshwater habitats to date, there is an

increasing community of researchers and practitioners interested in pond ecology and conservation, and a rapidly increasing awareness of the importance of ponds by society. This paper has highlighted some of the major themes and provides key questions for future pond research, which aim to address existing knowledge gaps and increase fundamental and practical understanding of pond ecology. However, to continue to progress interest, knowledge and awareness of pond ecosystems, international collaboration and commitment among researchers and end users is required. A better understanding of pond ecosystems will benefit society and wildlife by enabling more effective research-led conservation and management of pondscapes, facilitating their inclusion in environmental policy more clearly whilst simultaneously addressing many of the threats driving the decline in global freshwater biodiversity.

## **Acknowledgements**

The authors gratefully acknowledge the support of The British Ecological Society for funding and supporting the workshop. MJH would also like to thank to the University of Huddersfield for supporting the workshop. The authors would like to thank the reviewers of the draft manuscript for their valuable comments and input which have greatly improved this publication.

## **References**

- Abney, C.R., Balzer, S.W., Dueckman, A., Baylis, A. and Clements, D.R. 2019. Early spring and early vanishing wetlands as harbingers of the future? The climate change trap for ephemeral pond-breeding frogs. *Northwest Science* 93:52-65.
- Abrams, J.F., Hörig, L.A., Brozovic, R., Axtner, J., Crampton-Platt, A., Mohamed, A., Wong, S.T., Sollmann, R., Yu, D.W. and Wilting, A. 2019. Shifting up a gear with iDNA: from mammal detection events to standardized surveys. *Journal of Applied Ecology* 56:1637-1648.
- Alahuhta, J., et al. 2017. Global variation in the beta diversity of lake macrophytes is driven by environmental heterogeneity rather than latitude. *Journal of Biogeography* 44:1758-1769.

720 Alderton, E., Sayer, C.D., Davies, R., Lambert, S.J. and Axmacher, J.C. 2017. Buried alive: Aquatic  
 721 plants survive in 'ghost ponds' under agricultural fields. *Biological Conservation* 212:105-110.

722 Andersen, M.R., Sand-Jensen, K., Woolway, R.I. and Jones, I.D., (2017). Profound daily vertical  
 723 stratification and mixing in a small, shallow, wind-exposed lake with submerged macrophytes.  
 724 *Aquatic Sciences* 79:395-406.

725 Arlinghaus, R. 2005. A conceptual framework to identify and understand conflicts in recreational  
 726 fisheries systems, with implications for sustainable management. *Aquatic Resources, Culture and*  
 727 *Development* 1:145-174.

728 Bastien, N.R.P., Arthur, S. and McLoughlin, M.J. 2012. Valuing amenity: public perceptions of  
 729 sustainable drainage systems ponds. *Water and Environment Journal* 26:19-29.

730 Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W. and Courchamp, F., 2012. Impacts of climate  
 731 change on the future of biodiversity. *Ecology Letters* 15:365-377.

732 Bertoncin, A.P.d.S., Pinha, G.D., Baumgartner, M.T and Mormul, R.P. 2019. Extreme drought events  
 733 can promote homogenization of benthic macroinvertebrate assemblages in a floodplain pond in Brazil.  
 734 *Hydrobiologia* 826:379-393.

735 Biggs, J. and Williams, P. 2021. *Ponds, Pools and Puddles*. Harper Collins, London.

736 Biggs, J., et al. 2015. Using eDNA to develop a national citizen science-based monitoring programme  
 737 for the great crested newt (*Triturus cristatus*). *Biological Conservation* 183:19-28.

738 Biggs, J., Williams, P., Whitfield, M., Nicolet, P. and Weatherby, A. 2005. 15 years of pond  
 739 assessment in Britain: results and lessons learned from the work of Pond Conservation. *Aquatic*  
 740 *Conservation: Marine and Freshwater Ecosystems* 15:693-714.

741 Biggs, J., Von Fumetti, S. and Kelly-Quinn, M. 2017. The importance of small waterbodies for  
 742 biodiversity and ecosystem services: implications for policy makers. *Hydrobiologia* 793:3-39.

743 Bilton, D.T., Freeland, J.R. and Okamura, B. 2001. Dispersal in freshwater invertebrates. *Annual*  
 744 *Review of Ecology and Systematics* 32:159-181.

745 Bortolotti, L.E., Vinebrooke, R.D. and St Louis, V.L. 2016. Prairie wetland communities recover at  
746 different rates following hydrological restoration. *Freshwater Biology* 61:1874-1890.

747 Braund, M.R. 1997. School Ponds: Their Current Status and Likely Contribution to Education,  
748 Conservation and Local Environmental Enhancement. ERIC.

749 Brown, I. 2020. Challenges in delivering climate change policy through land use targets for  
750 afforestation and peatland restoration. *Environmental Science and Policy* 107:36-45.

751 Buxton, A.S., Groombridge, J.J., Zakaria, N.B. and Griffiths, R.A. 2017. Seasonal variation in  
752 environmental DNA in relation to population size and environmental factors. *Scientific Reports*  
753 7:46294.

754 Calabrese, S., Mezzanotte, V., Marazzi, F., Canobbio, S. and Fornaroli, R. 2020. The influence of  
755 multiple stressors on macroinvertebrate communities and ecosystem attributes in Northern Italy pre-  
756 Alpine rivers and streams. *Ecological Indicators* 115106408.

757 Calhoun, A.J.K., Muschet, D.M., Bell, K.P., Boix, D., Fitzsimons, J.A. and Isselin-Nondedeu, F.  
758 2017. Temporary wetlands: challenges and solutions to conserving a ‘disappearing’ ecosystem.  
759 *Biological Conservation* 211:3-11.

760 Capps, K. A., Berven, K. A., and Tiegs, S. D. 2015. Modelling nutrient transport and transformation  
761 by pool-breeding amphibians in forested landscapes using a 21-year dataset. *Freshwater Biology*  
762 60:500–511.

763 Chapman, D.S., Gunn, I.D., Pringle, H.E., Siriwardena, G.M., Taylor, P., Thackeray, S.J., Willby,  
764 N.J. and Carvalho, L., 2020. Invasion of freshwater ecosystems is promoted by network connectivity  
765 to hotspots of human activity. *Global Ecology and Biogeography*, 29:645-655.

766 Cieminski, K.L. and Flake, L.D. 1997. Mule deer and pronghorn use of wastewater ponds in a cold  
767 desert. *The Great Basin Naturalist* 57:327-337.

768 Compson, Z.G., et al. 2019. Network-based biomonitoring: exploring freshwater food webs with  
769 stable isotope analysis and DNA metabarcoding. *Frontiers in Ecology and Evolution* 7:395.

770 Coutts, A. M., Tapper, N. J., Beringer, J., Loughnan, M., and Demuzere, M. 2012. Watering our  
 771 cities: the capacity for water sensitive urban design to support urban cooling and improve human  
 772 thermal comfort in the Australian context. *Progress in Physical Geography* 37:2-28.

773 de Bell, S., Graham, H., Jarvis, S. and White, P. 2017. The importance of nature in mediating social  
 774 and psychological benefits associated with visits to freshwater blue space. *Landscape and Urban*  
 775 *Planning* 167:118-127.

776 Dickinson, J.L., Shirk, J., Bonter, D., Bonney, R., Crain, R.L., Martin, J., Phillips, T. and Purcell, K.,  
 777 2012. The current state of citizen science as a tool for ecological research and public engagement.  
 778 *Frontiers in Ecology and the Environment* 10:291-297.

779 Davies, B., Biggs, J., Williams, P., Whitfield, M., Nicolet, P., Sear, D., Bray, S. and Maund, S., 2008.  
 780 Comparative biodiversity of aquatic habitats in the European agricultural landscape. *Agriculture,*  
 781 *Ecosystems and Environment* 125:1-8.

782 De Bie, T., et al. 2012. Body size and dispersal mode as key traits determining metacommunity  
 783 structure of aquatic organisms. *Ecology Letters* 15:740-747.

784 Deiner, K., et al. 2017. Environmental DNA metabarcoding: Transforming how we survey animal and  
 785 plant communities. *Molecular Ecology* 26:5872-5895.

786 Department of Army, Corps of Engineers and US Environmental Protection Agency. Clean Water  
 787 Rule: Definition of “Waters of the United States”. 2015. Available from  
 788 [https://www.federalregister.gov/documents/2015/06/29/2015-13435/clean-water-rule-definition-of-](https://www.federalregister.gov/documents/2015/06/29/2015-13435/clean-water-rule-definition-of-waters-of-the-united-states)  
 789 [waters-of-the-united-states](https://www.federalregister.gov/documents/2015/06/29/2015-13435/clean-water-rule-definition-of-waters-of-the-united-states). Accessed 26/03/2021.

790 Dos Reis, I.C., Honório, N.A., de Barros, F.S.M., Barcellos, C., Kitron, U., Camara, D.C.P., Pereira,  
 791 G.R., Keppeler, E.C., da Silva-Nunes, M. and Codeço, C.T., 2015. Epidemic and endemic malaria  
 792 transmission related to fish farming ponds in the Amazon frontier. *PLoS One* 10:p.e0137521.

793 Downing, J.A., Cole, J.J., Middelburg, J.J., Striegl, R.G., Duarte, C.M., Kortelainen, P., Prairie, Y.T.  
 794 and Laube, K.A., 2008. Sediment organic carbon burial in agriculturally eutrophic impoundments  
 795 over the last century. *Global Biogeochemical Cycles* 22:1-10.



796 Elbrecht, V., Vamos, E.E., Meissner, K., Aroviita, J. and Leese, F. 2017. Assessing strengths and  
797 weaknesses of DNA metabarcoding-based macroinvertebrate identification for routine stream  
798 monitoring. *Methods in Ecology and Evolution* 8:1265–1275.

799 Fey, S. B., Mertens, A. N., and Cottingham, K. L. 2015. Autumn leaf subsidies influence spring  
800 dynamics of freshwater plankton communities. *Oecologia*, 178:875–885.

801 Foley, R. and Kistemann, T. 2015. Blue space geographies: Enabling health in place. *Health and Place*  
802 35:157-165.

803 Gallant, A.L., (2015). The challenges of remote monitoring of wetlands. *Remote Sensing* 7:10938-  
804 10950.

805 García-Girón, J., Heino, J., García-Criado, F., Fernández-Aláez, C. and Alahuhta, J., 2020. Biotic  
806 interactions hold the key to understanding metacommunity organisation. *Ecography* 43:1180-1190.

807 Gascon, M., Zijlema, W., Vert, C., White, M.P. and Nieuwenhuijsen, M.J. 2017. Outdoor blue spaces,  
808 human health and well-being: a systematic review of quantitative studies. *International Journal of*  
809 *Hygiene and Environmental Health* 220:1207-1221.

810 Gholizadeh, M.H., Melesse, A.M. and Reddi, L. 2016. A comprehensive review on water quality  
811 parameters estimation using remote sensing techniques. *Sensors* 16:1298.

812 Giardino, C., Bresciani, M., Villa, P. and Martinelli, A. 2010. Application of Remote Sensing in  
813 Water Resource Management: The Case Study of Lake Trasimeno, Italy. *Water Resources*  
814 *Management* 24:3885–3899.

815 Gledhill D.G., and James, P. 2012. Socio-economic variables as indicators of pond conservation in an  
816 urban landscape. *Urban Ecosystems* 15:849–861

817 Grooten, M. and Almond, R.E.A. 2018. Living Planet Report 2018: Aiming higher. World Wildlife  
818 Fund, Gland, Switzerland.

819 Hanski, I., 1998. Metapopulation dynamics. *Nature* 396:41-49.

820 Harper, L.R., et al. 2019a. Prospects and challenges of environmental DNA (eDNA) monitoring in  
821 freshwater ponds. *Hydrobiologia* 826:25-41.

822 Harper, L.R., Griffiths, N.P., Lawson Handley, L., Sayer, C.D., Read, D.S., Harper, K.J., Blackman,  
823 R.C., Li, J. and Hänfling, B., 2019b. Development and application of environmental DNA  
824 surveillance for the threatened crucian carp (*Carassius carassius*). *Freshwater Biology* 64:93-107.

825 Harper, L.R., Lawson Handley, L., Sayer, C.D., Read, D.S., Benucci, M., Blackman, R.C., Hill, M.J.  
826 and Hänfling, B. 2020. Assessing the impact of the threatened crucian carp (*Carassius carassius*) on  
827 pond invertebrate diversity: A comparison of conventional and molecular tools. *Molecular Ecology*  
828 00:1-18.

829 Hartel, T., Nemes, S., Cogălniceanu, D., Öllerer, K., Schweiger, O., Moga, C. I., and Demeter, L.  
830 2007. The effect of fish and aquatic habitat complexity on amphibians. *Hydrobiologia*, 583:173.

831 Hassall C. 2014. The ecology and biodiversity of urban ponds. *Wiley Interdisciplinary Reviews:*  
832 *Water* 1:187–206.

833 Hassall, C. 2015. Odonata as candidate macroecological barometers for global climate change.  
834 *Freshwater Science* 34:1040-1049

835 Hassall, C., Hollinshead, J. and Hull, A. 2012. Temporal dynamics of aquatic communities and  
836 implications for pond conservation. *Biodiversity and Conservation* 21:829-852.

837 Heino, J., et al. 2017. Integrating dispersal proxies in ecological and environmental research in the  
838 freshwater realm. *Environmental Reviews* 25:334-349.

839 Heino, J., et al. 2021. Lakes in the era of global change: moving beyond single-lake thinking in  
840 maintaining biodiversity and ecosystem services. *Biological Reviews* 96:89-106.

841 Heino, J. and Grönroos, M., 2017. Exploring species and site contributions to beta diversity in stream  
842 insect assemblages. *Oecologia* 183:151-160.

843 Heino, J., Melo, A.S., Siqueira, T., Soininen, J., Valanko, S. and Bini, L.M., 2015. Metacommunity  
844 organisation, spatial extent and dispersal in aquatic systems: patterns, processes and prospects.  
845 *Freshwater Biology* 60:845-869.

846 Hecnar, S. J., and M'Closkey, R. T. 1997. The effects of predatory fish on amphibian species richness  
847 and distribution. *Biological Conservation* 79:123–131.

848 Higgins, S.L., Thomas, F., Goldsmith, B., Brooks, S.J., Hassall, C., Harlow, J., Stone, D., Völker, S.  
849 and White, P., 2019. Urban freshwaters, biodiversity, and human health and well-being: Setting an  
850 interdisciplinary research agenda. *Wiley Interdisciplinary Reviews: Water* 6: p.e1339.

851 Hill, M.J., et al. 2018. New policy directions for global pond conservation. *Conservation Letters* 11:  
852 p.e12447.

853 Hill, M.J., Sayer, C.D. and Wood, P.J. 2016. When is the best time to sample aquatic  
854 macroinvertebrates in ponds for biodiversity assessment? *Environmental Monitoring and Assessment*  
855 188:194.

856 Hill, M.J., Biggs, J., Thornhill, I., Briers, R.A., Gledhill, D.G., White, J.C., Wood, P.J. and Hassall,  
857 C., 2017. Urban ponds as an aquatic biodiversity resource in modified landscapes. *Global Change*  
858 *Biology* 23:986-999.

859 Hill, M.J., Heino, J., Thornhill, I., Ryves, D.B. and Wood, P.J. 2017a. Effects of dispersal mode on  
860 the environmental and spatial correlates of nestedness and species turnover in pond communities.  
861 *Oikos* 126:1575-1585.

862 Hill, M.J., White, J.C., Biggs, J., Briers, R.A., Gledhill, D., Ledger, M.E., Thornhill, I., Wood, P.J.  
863 and Hassall, C., 2021. Local contributions to beta diversity in urban pond networks: Implications for  
864 biodiversity conservation and management. *Diversity and Distributions*.

865 Hintz, W.D. and Relyea, R.A. 2019. A review of the species, community, and ecosystem impacts of  
866 road salt salinisation in fresh waters. *Freshwater Biology* 64:1081–1097.

867 Holgerson, M.A., Lambert, M.R., Freidenburg, L.K. and Skelly, D.K. 2018. Suburbanization alters  
868 small pond ecosystems: shifts in nitrogen and food web dynamics. *Canadian Journal of Fisheries and*  
869 *Aquatic Sciences* 75:641-652.

870 Holgerson, M. A., Post, D. M., and Skelly, D. K. 2016. Reconciling the role of terrestrial leaves in  
871 pond food webs: A whole-ecosystem experiment. *Ecology* 97:1771–1782.

872 Holgerson, M.A. and Raymond, P.A. 2016. Large contribution to inland water CO<sub>2</sub> and CH<sub>4</sub>  
873 emissions from very small ponds. *Nature Geoscience* 9:222.

874 Holgerson, M.A., Duarte, A., Hayes, M.P., Adams, M.J., Tyson, J.A., Douville, K.A. and Strecker,  
875 A.L. 2019. Floodplains provide important amphibian habitat despite multiple ecological threats.  
876 *Ecosphere* 10:p.e02853.

877 Homan, R.N., Windmiller, B.S. and Reed, J.M. 2004. Critical thresholds associated with habitat loss  
878 for two vernal pool-breeding amphibians. *Ecological Applications* 14:547-1553.

879 Holzer, K.A. and Lawler, S.P. 2015. Introduced reed canary grass attracts and supports a common  
880 native amphibian. *The Journal of Wildlife Management* 79:1081-1090.

881 Hunter-Ayad, J. and Hassall, C. 2020. An empirical, cross-taxon evaluation of landscape-scale  
882 connectivity. *Biodiversity and Conservation* 29:1339-1359.

883 IPBES, 2018. The IPBES regional assessment report on biodiversity and ecosystem services for  
884 Europe and Central Asia. Rounsevell, M., Fischer, M., Torre-Marín Rando, A. and Mader, A. (eds).  
885 Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem  
886 Services, Bonn, Germany. 892 pages.

887 Ilg, C. and Oertli, B. 2017. Effectiveness of amphibians as biodiversity surrogates in pond  
888 conservation. *Conservation Biology* 31:437-445.

889 Ito, H.C., Shiraishi, H., Nakagawa, M. and Takamura, N., 2020. Combined impact of pesticides and  
890 other environmental stressors on animal diversity in irrigation ponds. *PloS One* 15: p.e0229052.

891 Janssens, L. and Stoks. R. 2013. Fitness effects of Chlorpyrifos in the damselfly *Enallagma*  
892 *cyathigerum* strongly depend upon temperature and food level and can bridge metamorphosis. PloS  
893 One 8:e68107-e68107.

894 Jantz, S.M., Barker, B., Brooks, T.M., Chini, L.P., Huang, Q., Moore, R.M., Noel, J. and Hurtt, G.C.,  
895 2015. Future habitat loss and extinctions driven by land-use change in biodiversity hotspots under  
896 four scenarios of climate-change mitigation. Conservation Biology 29:1122-1131.

897 Jeffries, M.J. 2011. The temporal dynamics of temporary pond macroinvertebrate communities over a  
898 10-year period. Hydrobiologia 661:391-405.

899 Jordano, P. 2017. What is long-distance dispersal? And a taxonomy of dispersal events. Journal of  
900 Ecology 105:75-84.

901 Juračka, P.J., Dobiáš, J., Boukal, D.S., Šorf, M., Beran, L., Černý, M. and Petrusek, A., 2019. Spatial  
902 context strongly affects community composition of both passively and actively dispersing pool  
903 invertebrates in a highly heterogeneous landscape. Freshwater Biology 64:2093-2106.

904 Kaunisto, K.M., Roslin, T., Sääksjärvi, I.E. and Vesterinen, E.J. 2017. Pellets of proof: First glimpse  
905 of the dietary composition of adult odonates as revealed by metabarcoding of feces. Ecology and  
906 Evolution 7:8588–8598.

907 Kissel, A.M., Halabisky, M., Scherer, R.D., Ryan, M.E. and Hansen, E.C., 2020. Expanding wetland  
908 hydroperiod data via satellite imagery for ecological applications. Frontiers in Ecology and the  
909 Environment, 18: 432-438.

910 Kleyheeg, E. and van Leeuwen, C.H.A. 2015. Regurgitation by waterfowl: An overlooked mechanism  
911 for long-distance dispersal of wetland plant seeds. Aquatic Botany 127:1-5.

912 Knight, T. M., McCoy, M. W., Chase, J. M., McCoy, K. A., and Holt, R. D. 2005. Trophic cascades  
913 across ecosystems. Nature 437:880–883.

914 Langdon, S.J., Marrs, R.H., Hosie, C.A., McAllister, H.A., Norris, K.M. and Potter, J.A. 2004.  
 915 *Crassula helmsii* in UK Ponds: Effects on Plant Biodiversity and Implications for Newt  
 916 Conservation1. Weed Technology 18:1349-1352.

917 Law, A., Jones, K.C. and Willby, N.J., 2014. Medium vs. short-term effects of herbivory by Eurasian  
 918 beaver on aquatic vegetation. Aquatic Botany 116:27-34.

919 Law, A., Baker, A., Sayer, C., Foster, G., Gunn, I.D., Taylor, P., Pattison, Z., Blaikie, J. and Willby,  
 920 N.J., 2019. The effectiveness of aquatic plants as surrogates for wider biodiversity in standing fresh  
 921 waters. Freshwater Biology 64:1664-1675.

922 Leibold, M.A., Holyoak, M., Mouquet, N., Amarasekare, P., Chase, J.M., Hoopes, M.F., Holt, R.D.,  
 923 Shurin, J.B., Law, R., Tilman, D. and Loreau, M. 2004. The metacommunity concept: a framework  
 924 for multi-scale community ecology. Ecology Letters 7:601-613.

925 Legendre, P. 2014. Interpreting the replacement and richness difference components of beta diversity.  
 926 Global Ecology and Biogeography 23:1324-1334.

927 Lewis-Phillips, J., Brooks, S.J., Sayer, C.D., Patmore, I.R., Hilton, G.M., Harrison, A., Robson, H.  
 928 and Axmacher, J.C. 2020. Ponds as insect chimneys: Restoring overgrown farmland ponds benefits  
 929 birds through elevated productivity of emerging aquatic insects. Biological Conservation 241:108253.

930 Lorenz, S., Rasmussen, J.J., Süß, A., Kalettka, T., Golla, B., Horney, P., Stähler, M., Hommel, B. and  
 931 Schäfer, R.B., 2017. Specifics and challenges of assessing exposure and effects of pesticides in small  
 932 water bodies. Hydrobiologia 793:213-224.

933 Lucieer, A., Turner, D., King, D.H. and Robinson, S.A., 2014. Using an Unmanned Aerial Vehicle  
 934 (UAV) to capture micro-topography of Antarctic moss beds. International Journal of Applied Earth  
 935 Observation and Geoinformation 27:53-62.

936 Lundy, L. and Wade, R. 2011. Integrating sciences to sustain urban ecosystem services. Progress in  
 937 Physical Geography 35:653-669.

938 Martinsen, K.T., Andersen, M.R. and Sand-Jensen, K., 2019. Water temperature dynamics and the  
 939 prevalence of daytime stratification in small temperate shallow lakes. *Hydrobiologia* 826:247-262.

940 Mauvisseau, Q., Coignet, A., Delaunay, C., Pinet, F., Bouchon, D. and Souty-Grosset, C. 2018.  
 941 Environmental DNA as an efficient tool for detecting invasive crayfishes in freshwater ponds.  
 942 *Hydrobiologia* 805:163–175.

943 McCaffery, M. and Eby, L., 2016. Beaver activity increases aquatic subsidies to terrestrial consumers.  
 944 *Freshwater Biology* 61:518-532.

945 Moss, B. 2017. Ponds and small lakes. Exeter, UK: Pelagic Publishing.

946 Nassauer, J., 2004. Monitoring the success of metropolitan wetland restorations: Cultural  
 947 sustainability and ecological function. *Wetlands* 24:756-765.

948 Nummi, P., Kattainen, S., Ulander, P., and Hahtola, A., 2011. Bats benefit from beavers: a facilitative  
 949 link between aquatic and terrestrial food webs. *Biodiversity and Conservation* 20:851-859.

950 Okamura, B., Hartikainen, H. and Trew, J. 2019. Zoochory and freshwater biodiversity: general  
 951 insights from bryozoans. *Frontiers in Ecology and Evolution* 7:29.

952 Parsons, K.M., Everett, M., Dahlheim, M. and Park, L. 2018. Water, water everywhere:  
 953 environmental DNA can unlock population structure in elusive marine species. *Royal Society Open*  
 954 *Science* 5:180537.

955 Patoka, J., Bláha, M., Kalous, L., and Kouba, A. 2017. Irresponsible vendors: Non-native, invasive  
 956 and threatened animals offered for garden pond stocking. *Aquatic Conservation: Marine and*  
 957 *Freshwater Ecosystems* 27:692–697.

958 Pavelková, R., Frajer, J., Havlíček, M., Netopil, P., Rozkošný, M., David, V., Dzuráková, M. and  
 959 Šarapatka, B., 2016. Historical ponds of the Czech Republic: an example of the interpretation of  
 960 historic maps. *Journal of Maps* 12:551-559.

961 Peacock, M., Audet, J., Jordan, S., Smeds, J. and Wallin, M.B. 2019. Greenhouse gas emissions from  
 962 urban ponds are driven by nutrient status and hydrology. *Ecosphere* 10:p.e02643.

963 Pereira Souza, F., Leite Costa, M.E. and Koide, S., 2019. Hydrological modelling and evaluation of  
 964 detention ponds to improve urban drainage system and water quality. *Water* 11:1547.

965 Piggott, J.J., Lange, K. Townsend, C.R. and Matthaei, C.D. 2012. Multiple Stressors in Agricultural  
 966 Streams: A Mesocosm Study of Interactions among Raised Water Temperature, Sediment Addition  
 967 and Nutrient Enrichment. *Plos One* 7:e49873

968 Raby, M., Nowierski, M., Perlov, D., Zhao, X., Hao, C., Poirier, D.G., and Sibley, P.K. 2018. Acute  
 969 toxicity of 6 neonicotinoid insecticides to freshwater invertebrates. *Environmental Toxicology and*  
 970 *Chemistry* 37:1430–1445.

971 Radchuk, V. et al., 2019. Adaptive responses of animals to climate change are most likely insufficient.  
 972 *Nature Communications* 10:1-14.

973 Ramsar Convention Secretariat, 2013. The Ramsar convention manual: a guide to the convention on  
 974 wetlands (Ramsar, Iran, 1971), 6th ed. Ramsar Convention Secretariat, Gland.

975 Regester, K. J., Lips, K. R., and Whiles, M. R. 2006. Energy flow and subsidies associated with the  
 976 complex life cycle of ambystomatid salamanders in ponds and adjacent forest in southern Illinois.  
 977 *Oecologia* 147:303–314.

978 Rose, R.A., et al. 2015. Ten ways remote sensing can contribute to conservation. *Conservation*  
 979 *Biology* 29:350-359.

980 Rosentreter, J.A., Borges, A.V., Deemer, B.R., Holgerson, M.A., Liu, S., Song, C., Melack, J.,  
 981 Raymond, P.A., Duarte, C.M., Allen, G.H. and Olefeldt, D., 2021. Half of global methane emissions  
 982 come from highly variable aquatic ecosystem sources. *Nature Geoscience* 1-6.

983 Rosset, V., Angélibert, S., Arthaud, F., Bornette, G., Robin, J., Wezel, A., Vallod, D. and Oertli, B.,  
 984 2014. Is eutrophication really a major impairment for small waterbody biodiversity? *Journal of*  
 985 *Applied Ecology* 51:415-425.



986 Ryan, M.E., Palen, W.J., Adams, M.J. and Rochefort, R.M., 2014. Amphibians in the climate vise:  
987 loss and restoration of resilience of montane wetland ecosystems in the western US. *Frontiers in*  
988 *Ecology and the Environment* 12:232-240.

989 Sayer, C.D., Copp, G.H., Emson, D., Godard, M.J., Zieba, G., and Wesley, K.J. 2011. Towards the  
990 conservation of crucian carp *Carassius carassius*: understanding the extent and causes of decline  
991 within part of its native English range. *Journal of Fish Biology* 79:1608-1624.

992 Sayer, C., Andrews, K., Shilland, E., Edmonds, N., Edmonds-Brown, R., Patmore, I., Emson, D. and  
993 Axmacher, J., 2012. The role of pond management for biodiversity conservation in an agricultural  
994 landscape. *Aquatic Conservation: Marine and Freshwater Ecosystems* 22:626-638.

995 Sayer, C.D., Shilland, E.M., Greaves, H., Dawson, B., Patmore, I., Emson, D., Alderton, E.,  
996 Robinson, P., Andrews, K., Axmacher, J.A., Wiik, E. 2013. Managing Britain's ponds - conservation  
997 lessons from a Norfolk farm. *British Wildlife* 25:21-28.

998 Sayer, C.D. 2014. Conservation of aquatic landscapes: ponds, lakes, and rivers as integrated systems.  
999 *Wiley Interdisciplinary Reviews: Water* 1: 573-585.

1000 Sayer, C.D. and Greaves, H. 2020. Making an impact on UK farmland pond conservation. *Aquatic*  
1001 *Conservation: Marine and Freshwater Ecosystems* 30:1821-1828.

1002 Seto, K.C., Güneralp, B. and Hutrya, L.R. 2012. Global forecasts of urban expansion to 2030 and  
1003 direct impacts on biodiversity and carbon pools. *Proceedings of the National Academy of Sciences*  
1004 109: 16083-16088.

1005 Seabloom, E.W. and van der Valk, A.G. 2003. Plant diversity, composition, and invasion of restored  
1006 and natural prairie pothole wetlands: Implications for restoration. *Wetlands* 23:1-12.

1007 Søndergaard, M., Jeppesen, E. and Jensen, J.P. 2005. Pond or lake: does it make any difference?  
1008 *Archiv für Hydrobiologie*, 162:143-165.

1009 Sebastián-González, E. and Green, A.J., 2014. Habitat use by waterbirds in relation to pond size,  
 1010 water depth, and isolation: lessons from a restoration in southern Spain. *Restoration Ecology* 22:311-  
 1011 318.

1012 Smith, T. and Buckley, P. 2020. Biological Flora of the British Isles: *Crassula helmsii*. *Journal of*  
 1013 *Ecology* 108:97-813.

1014 Soininen, J., Bartels, P.I.A., Heino, J., Luoto, M. and Hillebrand, H., 2015. Toward more integrated  
 1015 ecosystem research in aquatic and terrestrial environments. *BioScience* 65:174-182.

1016 Stenroth, K., Polvi, L.E., Fälström, E., and Jonsson, M. 2015. Land-use effects on terrestrial  
 1017 consumers through changed size structure of aquatic insects. *Freshwater Biology* 60:136-149.

1018 Stewart, R.I.A., Andersson, G.K.S., Brönmark, C., Klatt, B.K., Hansson, L-A., Zülsdorff, V. and  
 1019 Smith, H.G. 2017. Ecosystem services across the aquatic-terrestrial boundary: linking ponds to  
 1020 pollination. *Basic and Applied Ecology* 18:13–20.

1021 Suh, A. and Samways, M. 2005. Significance of temporal changes when designing a reservoir for  
 1022 conservation of dragonfly diversity. *Biodiversity and Conservation* 14:165–178.

1023 Taylor, S., Gilbert, P.J., Cooke, D.A., Deary, M.E. and Jeffries, M.J. 2019. High carbon burial rates  
 1024 by small ponds in the landscape. *Frontiers in Ecology and the Environment* 17:25-31.

1025 Teo, H. C., Hill, M. J., Lechner, A., Yenn, T. F. and Gibbins, C., In review. Landscape-scale remote  
 1026 sensing and classification of pond and lake habitat diversity in a rapidly expanding tropical city:  
 1027 Greater Kuala Lumpur, Malaysia. *Remote Sensing of Environment*.

1028 Thiere, G., Milenkovski, S., Lindgren, P-E., Sahlén, G., Berglund, O. and Weisner, S.E.B. 2009.  
 1029 Wetland creation in agricultural landscapes: Biodiversity benefits on local and regional scales.  
 1030 *Biological Conservation* 142:964-973.

1031 Thornhill, I., Batty, L., Hewitt, M., Friberg, N.R. and Ledger, M.E., 2018. The application of graph  
 1032 theory and percolation analysis for assessing change in the spatial configuration of pond networks.  
 1033 *Urban Ecosystems* 21:213-225.

1034 Tiegs, S.D., Costello, D.M., Isken, M.W., Woodward, G., McIntyre, P.B., Gessner, M.O., Chauvet,  
 1035 E., Griffiths, N.A., Flecker, A.S., Acuña, V. and Albariño, R., 2019. Global patterns and drivers of  
 1036 ecosystem functioning in rivers and riparian zones. *Science Advances* 5:p.eaav0486.

1037 Twining, C.W., Brenna, J.T., Hairston, N.G., and Flecker, A.S., 2016. Highly unsaturated fatty acids  
 1038 in nature: what we know and what we need to learn. *Oikos* 125:749-760.

1039 Vad, et al. 2017. Wartime scars or reservoirs of biodiversity? The value of bomb crater ponds in  
 1040 aquatic conservation. *Biological Conservation* 209:253-262.

1041 VanAcker, M.C., Lambert, M.R., Schmitz, O.J. and Skelly, D.K. 2019. Suburbanization increases  
 1042 Echinostome infection in green frogs and snails. *EcoHealth* 16:235-247.

1043 Viana, D.S., Santamaría, L. and Figuerola, J. 2016. Migratory birds as global dispersal vectors.  
 1044 *Trends in Ecology and Evolution* 31:763-775.

1045 Vico, G., Tamburino, L. and Rigby, J.R., 2020. Designing on-farm irrigation ponds for high and  
 1046 stable yield for different climates and risk-coping attitudes. *Journal of Hydrology* 584:p.124634.

1047 Walton, R.E., Sayer, C.D., Bennion, H. and Axmacher, J.C. 2021. Once a pond in time: employing  
 1048 palaeoecology to inform farmland pond restoration. *Restoration Ecology* 29:p.e13301.

1049 Walton, R.E., Sayer, C.D., Bennion, H. and Axmacher, J.C. 2020. Nocturnal pollinators strongly  
 1050 contribute to pollen transport of wild flowers in an agricultural landscape. *Biology Letters*  
 1051 16:p.20190877.

1052 Webb, J.R., Leavitt, P.R., Simpson, G.L., Baulch, H.M., Haig, H.A., Hodder, K.R. and Finlay, K.  
 1053 2019. Regulation of carbon dioxide and methane in small agricultural reservoirs: optimizing potential  
 1054 for greenhouse gas uptake. *Biogeosciences* 16:4211-4227.

1055 Willby, N.J., Law, A., Levanoni, O., Foster, G. and Ecke, F. 2019. Rewilding wetlands: beaver as  
 1056 agents of within-habitat heterogeneity and the responses of contrasting biota. *Philosophical*  
 1057 *Transactions of the Royal Society B: Biological Sciences* 373:20170444.

1058 Williams, D.D., 1997. Temporary ponds and their invertebrate communities. *Aquatic Conservation:*  
1059 *Marine and Freshwater Ecosystems* 7:105-117.

1060 Williams, P. 2019. What's happening to the quality of our best ponds? A re-survey of National Pond  
1061 Survey sites after 24 years. Freshwater Habitats Trust.

1062 Williams, P., Biggs, J. and Nicolet, P. 2010. Comment: new clean-water ponds—a way to protect  
1063 freshwater biodiversity. *British Wildlife* 22:77.

1064 Williams, P., Biggs, J., Stoate, C., Szczur, J., Brown, C. and Bonney, S. 2020. Nature based measures  
1065 increase freshwater biodiversity in agricultural catchments. *Biological Conservation* 244:108515.

1066 Williams, P.J., Biggs, J., Crowe, A., Murphy, J., Nicolet, P., Weatherby, A., and Dunbar, M. 2010. CS  
1067 Technical Report No. 7/07 Countryside Survey: Ponds Report from 2007. Lancaster.

1068 Winder, M. and Schindler, D.E. 2004. Climate change uncouples trophic interactions in an aquatic  
1069 ecosystem. *Ecology* 85:2100-2106.

1070 Wood, P.J, Greenwood, M.T., and Agnew, M.D. 2003. Pond biodiversity and habitat loss in the UK.  
1071 *Area* 35:206–216.

1072 Wood, P.J., Greenwood, M.T., Barker, S.A. and Gunn, J. 2001. The effects of amenity management  
1073 for angling on the conservation value of aquatic invertebrate communities in old industrial ponds.  
1074 *Biological Conservation*, 102:17-29.

1075 Woodget, A.S., Carbonneau, P.E., Visser, F. and Maddock, I.P. 2015. Quantifying submerged fluvial  
1076 topography using hyperspatial resolution UAS imagery and structure from motion photogrammetry.  
1077 *Earth Surface Processes and Landforms* 40:47-64.

1078 Yvon-Durocher, G., Hulatt, C.J., Woodward, G. and Trimmer, M. 2017. Long-term warming  
1079 amplifies shifts in the carbon cycle of experimental ponds. *Nature Climate Change* 7:209-213.

1080 **Tables**

1081 Table 1 –Key research themes, research questions and future research identified in this paper to advance understanding of pond ecology and increase  
1082 opportunities for pond conservation.

Key research theme	Context	Future research	Key Research question(s)
Definition of pond habitats	A longstanding lack of congruency of how pond habitats are identified and defined.	Development of an overarching, process-based global definition of a pond.	1. Are there critical thresholds in physical, biological and chemical processes, and ecosystem function, which reflect changing waterbody size?
Global and long-term data	Minimal research examining global- and international-scale patterns in pond numbers, diversity and functioning. Pond studies dominated by single season or single year data collection.	Systematic consolidation of existing datasets and international collaborative studies to examine global-scale biodiversity patterns. Collection of long-term data to examine temporal trends in response to global environmental change.	1. What are the abiotic and biotic drivers of pond environmental and biological patterns at different spatial and temporal scales? 2. How do biological communities within ponds respond to global environmental change along spatial and temporal gradients?
Anthropogenic stressors	Little research examining the effect of the interaction/additive effect of multiple stressors (pollution, invasive species and climate change) on pond habitats.	Increased understanding of water chemistry standards for ponds. Examine the effects of neonicotinoid and microplastics pollution, invasive species colonisation and changes in pondscape connectivity on pond biodiversity. Examine the interaction between short- and long-term climate effects in ponds. Greater research focus on the value of small water bodies as a scalable solution to carbon sequestration.	1. What are the interactive effects of multiple-stressors on pond ecosystem resilience? 2. How do pollutants impact pond ecological functioning and resilience? 3. How widely are ponds affected by non-native species and what are the dynamics of their spread? 4. How do non-native species introductions affect trophic interactions in pond ecosystems? 5. What are the potential ecological implications for pond communities associated with different climate change scenarios? 6. How will climate change affect the interactions between aquatic and terrestrial food webs?
Aquatic-terrestrial interactions	A lack of understanding on the role that pond communities play in metaecosystem dynamics, trophic and food web structure of terrestrial communities and the role of predation for aquatic-terrestrial communities.	Examine the interaction between freshwater ecological health and terrestrial ecological health. Identify the role of aquatic biota in structuring terrestrial communities and food webs.	1. Does the nutritional value of aquatic macroinvertebrates for terrestrial predators reflect pond water quality? 2. How important are terrestrial vertebrates for the dispersal of pond biota and for maintaining local environmental heterogeneity and does this vary systematically with pond setting? 3. How do aquatic–terrestrial interactions in predation influence pond community structure and vice versa?

Succession and disturbance	The effect of disturbance on pond communities (at local and landscape scales) is poorly understood, in part due to the lack of long-term data and anthropogenic location of many ponds.	Increase fundamental understanding of how disturbance structures pond communities spatially and temporally across different land-use types. Examine the life history strategies and mechanisms that species and communities employ to survive disturbance events.	<ol style="list-style-type: none"> <li>1. What role do natural disturbances play in structuring aquatic and terrestrial biodiversity in pond habitats across landscapes minimally impacted by human influences?</li> <li>2. What are the best management strategies to reduce detrimental disturbances and increase positive disturbance in anthropogenically dominated landscapes?</li> <li>3. What are the principal trajectories, timescales and outcomes of pond succession across different pond types?</li> </ol>
Freshwater connectivity	Research is focussed on lotic habitats and typically examines freshwater habitats in isolation. The lack of high-resolution pond mapping is inhibiting research understanding the role of connectivity in the large-scale patterns of pond communities.	<p>Examine how spatial processes influence the patterns of actively and passively dispersing pond taxa across landscape types.</p> <p>Examine spatial processes operating across the watery landscape by incorporating multiple freshwater habitats.</p> <p>Characterise where connectivity is an advantage and where it represents a risk to pond biodiversity.</p>	<ol style="list-style-type: none"> <li>1. How does connectivity between pond habitats influence trophic interactions that bridge aquatic-terrestrial divides?</li> <li>2. How do species move between ponds and other freshwaters and what are the dominant mechanisms?</li> <li>3. How are pond networks (pondscapes) best designed or managed to ensure that rarer and less mobile species benefit?</li> <li>4. What is the role of spatial processes in assemblage structure and does pond context or landscape create regional differences?</li> </ol>
Pond monitoring and technological advances	eDNA, remotely sensed and drone-based data have been poorly applied to pond monitoring and multi-taxon assessments. Advances in biostatistics may increase effectiveness of pond conservation.	The use of remote sensing and drone technologies to identify the distribution and physicochemical properties of pond environments. Using multiple technologies (eDNA, UAV) effectively to answer fundamental, multitrophic ecological questions.	<ol style="list-style-type: none"> <li>1. Can molecular tools be used to assess the distribution of conservation priority and invasive species as well as community diversity at the pondscape scale?</li> <li>2. Are molecular tools and remote sensing able to identify the effects of anthropogenic stressors and climate change on ponds and improve management strategies to mitigate stressors?</li> <li>3. Can UAV-based data collection record physico-chemical and spatial characteristics of ponds and pondscapes more accurately than conventional data collection?</li> <li>4. How can the development of new statistical analyses in biodiversity assessment contribute to more effective pond conservation planning?</li> </ol>
Socio economic factors	Limited understanding of the personal, social and educational contributions made by ponds, and governance conflicts that exist between users of ponds. There remain gaps in our knowledge of the contribution of ponds to human health and wellbeing.	<p>Studies of the human dimensions of stakeholder engagement in pond creation and management are needed.</p> <p>Explore the relationships between ponds, wellbeing, access, and habitat quality.</p> <p>Assess the complexities of engagement, the, educational value of ponds, and the use of ponds in fostering an environmental conscience.</p>	<ol style="list-style-type: none"> <li>1. How do ponds contribute to human physical and mental health and wellbeing within urban and rural populations?</li> <li>2. What are the barriers (including social, cultural, institutional, emotional, communicative, and governance) in stakeholder-pond conservation interactions, and how might these be addressed?</li> <li>3. What are the short- and long-term effects of environmental education for pond conservation?</li> </ol>

Conservation, management and policy	There is a need to increase fundamental understanding of pond ecosystems to inform practical pond conservation and management. Ponds are largely excluded from environmental policy and legislation.	Examine the mechanisms that affect pond creation and restoration success. A need for studies to assess medium to long-term pond conservation and management measures. Identify opportunities for the inclusion of ponds in environmental and conservation policy. Quantify the distribution of rare species in ponds, and high-quality pond sites for conservation.	<ol style="list-style-type: none"> <li>1. Where are the most ecologically important ponds at national and international scales?</li> <li>2. How does pond creation and management affect biotic communities at local and landscape-scales in different environmental settings?</li> <li>3. How can we best conserve ponds in anthropogenically dominated landscapes?</li> <li>4. What are the mechanisms required to better incorporate ponds into national and international environmental policy and legislation?</li> </ol>
---	--	--	--

1083

1084 **Figure Captions**

1085 **Figure 1** - Total number of peer reviewed publications based on the search topic (a) “biodiversity” or  
1086 (b) “conservation”, with “pond”, “stream”, “lake” or “river” between 2000-2019, using the Scopus  
1087 database.

1088 **Figure 2** – The nine priority research themes and their contribution to the conservation and  
1089 management of pond ecosystems.

1090



1091 **Figure 1**



