

Northumbria Research Link

Citation: Ersek, Vasile, Sharples, Jamie and Williams, Thomas (2022) Stable hydrogen and oxygen isotope abundance of major bottled water brands sold in the United Kingdom. *Isotopes in Environmental and Health Studies*, 58 (2). pp. 113-120. ISSN 1025-6016

Published by: Taylor & Francis

URL: <https://doi.org/10.1080/10256016.2021.2005041>
<<https://doi.org/10.1080/10256016.2021.2005041>>

This version was downloaded from Northumbria Research Link:
<https://nrl.northumbria.ac.uk/id/eprint/47683/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>


This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



**Northumbria
University**
NEWCASTLE



UniversityLibrary



Stable hydrogen and oxygen isotope abundance of major bottled water brands sold in the United Kingdom

Journal:	<i>Isotopes in Environmental & Health Studies</i>
Manuscript ID	GIEH-2021-0044.R1
Manuscript Type:	Short Communication
Date Submitted by the Author:	24-Sep-2021
Complete List of Authors:	Ersek, Vasile; Northumbria University, Department of Geography and Environmental Sciences Sharples, Jamie; Northumbria University, Department of Geography and Environmental Sciences William, Thomas; Northumbria University, Department of Geography and Environmental Sciences
Keywords:	stable isotopes, bottled water, oxygen-18, hydrogen-2, groundwater < isotope hydrology, United Kingdom, water cycle < isotope hydrology, isotope geochemistry < Isotope geology, water resources, drinking water

SCHOLARONE™
Manuscripts

1
2
3 **Stable hydrogen and oxygen isotope abundance of major bottled water**
4 **brands sold in the United Kingdom**
5
6

7
8 Vasile Ersek, Jamie Sharples and William Thomas
9

10
11 *Department of Geography and Environmental Sciences, Ellison Building, Northumbria*
12 *University, Newcastle upon Tyne, UK*
13
14

15 Contact Vasile Ersek Email: vasile.ersek@northumbria.ac.uk
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Stable hydrogen and oxygen isotope abundance of major bottled water brands sold in the United Kingdom

Bottled water in the UK has a ~20 % share of the soft drinks market with a sales value of >£1.5 billion. Bottled water is susceptible to fraud and it is important to characterise the chemical signature of aquifers used by the bottled water industry. Measuring $^{18}\text{O}/^{16}\text{O}$ and $^2\text{H}/^1\text{H}$ ratios in bottled water is one important step in fraud prevention and aquifer characterisation as these ratios in groundwater tend to be stable or change very slowly through time. Here we characterise the isotopic signature of 30 brands of bottled water sold in the UK. The average $\delta^{18}\text{O}$ of bottled waters is -7.4 and -48.4 for $\delta^2\text{H}$. This isotopic composition is closely related to that of the annual rainfall and follows latitudinal and longitudinal gradients which combine to explain 77 % of the $\delta^{18}\text{O}$ variance.

Keywords: bottled water; geochemistry; hydrogen-2; isotope geology; isotope hydrology; oxygen-18; United Kingdom

1. Introduction

Bottled water (defined as single-use packaged water) has become increasingly important in the human diet with rises in consumption driven by complex factors, including the perceived higher purity of bottled water, rapid population growth, pollution of natural resources, and climate change [1,2]. During the last decade, bottled water consumption has increased by 175 % in many low and middle income countries, while in developed countries there has been an increase of 26 % over the same period [3]. This raises significant concerns about the environmental effects of the bottled water industry [4]. In addition, a major challenge associated with bottled water consumption is their authentication [5] which is required due to the potential for product misrepresentation and fraud in the form of mislabelling source location. Oxygen and hydrogen stable isotopes have also been used as useful tracers in detecting the authenticity and origin of other beverages and food products, including wines [6], ciders

1
2
3 [7] or roasted coffee [8].
4

5 Stable isotopes in bottled water can be used for fraud prevention [9], as the
6 water reflects the environment from which it originates [10]. Most bottled water
7 isotopic variation occurs as a result of differences in source location [11], with the main
8 spatial patterns occurring as a result of latitudinal temperature gradients. There have
9 been many studies of isotopes in drinking water [10,12–15] which have identified
10 spatial correlations between the isotopic ratios of bottled water, tap water and
11 precipitation. Studies have found that the stable isotope composition of bottled water is
12 similar to that of locally available water sources and precipitation [16,17].
13
14
15
16
17
18
19
20
21
22
23

24 The aim of this study is to investigate the isotopic composition of some of the
25 most popular brands of bottled water for sale in the United Kingdom (UK) and, to our
26 knowledge, this is the first study of this type in this country. In the UK, bottled water
27 has a share of 20.6 % of soft drink sales with a total volume of sales in 2019 of
28 2.811 billion litres and a value of 1.6 billion British pounds [18]. There are three types
29 of bottled waters sold in the UK market: natural mineral water, spring water and bottled
30 drinking water. Governmental regulations are different for the four nations of the UK.
31 The Department for Environment, Food and Rural Affairs regulates bottled waters in
32 England, the Food Standards Scotland is responsible for regulations in Scotland, and the
33 Food Standards Agency is responsible for the regulation of this sector in Wales and
34 Northern Ireland. In broad terms, they all require bottled waters to be free from disease-
35 causing bacteria and parasites and meet limits for chemical, microbial and radioactive
36 substances. Natural mineral waters and spring waters must come from an underground
37 source, be bottled at the source, and be protected from pollution. Natural mineral waters
38 have the additional requirements that the chemical composition must be labelled, be
39 stable over a lengthy period of time, and must not change from source to bottle.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

2. Materials and methods

The acquisition of bottled water samples was completed by purchasing those manufactured and commercially available in the UK between June 2018 and April 2021. A total of 30 different bottled water brands were purchased with volumes ranging from 330 ml to 2 L. The brands purchased in this study are identified as the most popular bottled water brands sourced in the UK [19]. Purchases were made from local shops in NE England with three bottles for each brand being acquired separately. The purchase of bottled water samples was not limited to still water, with both still and sparkling samples being collected from the same brand (where possible). The origin of each sample (Figure 1, Table 1) was obtained from the brand labelling where the bottles clearly state the contents were bottled at source. All waters were packaged in polyethylene terephthalate (PET) bottles, except for Waitrose Royal Deeside and M&S Mountain Water Speyside which used glass bottles. Geographical coordinates for each location were obtained from Google Earth, and the water source is assumed to be within 15 km of the location indicated on the bottle. Samples remained unopened and in their original bottles from the time of purchase until analysis and were stored in a cool, dark environment. Secondary data containing the stable isotope composition of precipitation across the UK was obtained from the GNIP database [20]. We constructed the meteoric water line for the UK based on the GNIP stations at Altnabreac (1981–1982, 13 samples), Armagh Observatory (2012–2019, 87 samples), Fleam Dyke (1980–1983, 48 samples), Inchnadamph (2003–2005, 22 samples; [21]), Keyworth (1985–1996, 118 samples) and Wallingford (1979–2015, 409 samples). Secondary bottled water $\delta^{18}\text{O}$ and $\delta^2\text{H}$ values were obtained from Bowen et al. [16] for two brands to increase the stable isotope dataset. In order to compare the consistency of isotopic composition of bottled water through time, we have also analysed bottled water from

1
2
3 Blackford, Scotland, which was also included in [16].
4

5 The waters were analysed on a Los Gatos Research off-axis integrated cavity
6 output laser absorption spectrometer (TWIA-45EP) to simultaneously determine $^2\text{H}/^1\text{H}$
7 and $^{18}\text{O}/^{16}\text{O}$ stable isotope ratios in liquid water [22]. Based on the expected isotopic
8 range of our samples, we used standards provided by Los Gatos Research, Inc. (San
9 Jose, CA, USA) which were calibrated to the VSMOW2-SLAP2 scale. The standards
10 have the following $\delta^2\text{H}$ and $\delta^{18}\text{O}$ values, respectively (expressed relative to VSMOW):
11 lgr1e: -165.7 and -21.28 ‰, lgr2e: -123.8 and -16.71 ‰, lgr3e: -79.6 and -11.04 ‰,
12 and lgr5e: -9.9 and -2.99 ‰. In addition, for each set of measurements, the standard
13 lgr4e (-51.00 and -7.69 ‰) was run as internal control as it was the most similar to the
14 expected isotopic range of our samples. Each measurement included two preparatory
15 injections (to help mitigate any memory effects prior to measured injections) followed
16 by six measured injections. The isotopic value of each sample or standard was
17 determined based on the mean of the last four injections, with the first two injections
18 ignored in order to remove any residual memory effects left after the preparatory
19 injections. For each batch of samples, a full set of standards were run at the beginning
20 and end of the batch, and a standard was measured after every fifth sample. Quality
21 control and data processing was performed via the LWIA Post Analysis software
22 (version 4.5.0), developed by Los Gatos Research. The analytical precision for the
23 reported values is better than 0.2 ‰ for $\delta^{18}\text{O}$ and 0.8 ‰ for $\delta^2\text{H}$.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50 **3. Results and discussion**

51
52 The average value of all bottled waters is -7.5 ± 0.9 ‰ for $\delta^{18}\text{O}$ and -48.4 ± 6.8 ‰ for
53 $\delta^2\text{H}$. Three of the waters were available in both still and sparkling form, and it is
54
55
56
57
58
59
60 apparent that the process of adding CO_2 to bottled waters does not have a significant

1
2
3 impact on the isotopic signature of sparkling waters (Table 1). The lowest isotopic
4
5 values were found at Glenlivet Estate in Scotland ($\delta^{18}\text{O} = -9.6 \pm 0.18 \text{ ‰}$ and
6
7 $\delta^2\text{H} = -61 \pm 0.52 \text{ ‰}$) and the highest in SW Wales ($\delta^{18}\text{O} = -5.5 \pm 0.1 \text{ ‰}$ and
8
9 $\delta^2\text{H} = -32.1 \pm 0.2 \text{ ‰}$) (Table 1). The stable isotope ratios of water bottled in Blackford,
10
11 Scotland were reported in 2005 as $-8.2 \pm 0.07 \text{ ‰}$ for $\delta^{18}\text{O}$ and $-56 \pm 0.6 \text{ ‰}$ for $\delta^2\text{H}$
12
13 [16], which is similar to our findings here (Table 1). This suggests that the isotopic
14
15 signature of this water source remains constant over a long time as is required by UK's
16
17 bottled water regulations.
18
19
20
21

22 The water bottling sites span a latitudinal range of 5.7° , from Brecon Beacons,
23
24 Wales in the south to Glenlivet Estate, Scotland in the north, and extend over 4.7°
25
26 longitude, from Ballymena in the west to Morpeth in the east (Figure 1). In order to
27
28 evaluate the role of the geographical position in controlling the isotopic ratios of bottled
29
30 waters, we performed linear regressions of $\delta^{18}\text{O}$ vs latitude and longitude. There is a
31
32 significant correlation between latitude and $\delta^{18}\text{O}$ (p -value = 0.0003), with latitude
33
34 explaining 58 % of $\delta^{18}\text{O}$ variance (Figure 2). The correlation of $\delta^{18}\text{O}$ with longitude is
35
36 much weaker than with latitude ($r^2 = 0.16$, p -value < 0.005). However, using a multiple
37
38 regression, the latitude and longitude combined explain 77 % of $\delta^{18}\text{O}$ variance
39
40 ($\delta^{18}\text{O} = 13.997 + (-0.418) \cdot \text{Latitude} + (-0.354) \cdot \text{Longitude}$). The spatial pattern of
41
42 isotopic composition of bottled water therefore follows latitudinal and longitudinal
43
44 gradients similar to those observed in precipitation and groundwater in the UK [23].
45
46 The latitudinal control is primarily due to changes in mean annual temperature and
47
48 degree of rainout, while the longitudinal effect is due to the Rayleigh fractionation of
49
50 airmasses affecting the area [24], which are dominated by southwesterly flow, and their
51
52 interaction with local topography. Since we do not have information about the recharge
53
54 rate, the age of groundwaters used in bottled waters in this study, and detailed
55
56
57
58
59
60

1
2
3 hydrogeological characterisation of each well, a comparison with variables such as
4
5 temperature, precipitation amount or altitude is not feasible here.
6
7

8 The plot of $\delta^{18}\text{O}$ vs $\delta^2\text{H}$ for bottled waters has a slope of 7.4, which is similar to
9
10 the slope of 7.2 for the UK meteoric water line, and is close to the slope of 8 for the
11
12 global meteoric water line (Figure 3). We can therefore infer that the bottled
13
14 groundwaters are relatively young and reflect the isotopic composition of modern
15
16 precipitation. The $\delta^{18}\text{O}$ vs $\delta^2\text{H}$ slope in precipitation can vary significantly both on a
17
18 seasonal basis and also from year to year.
19
20
21
22

23 **4. Conclusions**

24
25 We characterised the isotopic composition of 30 brands of water sourced and bottled in
26
27 the UK. The isotopic composition of bottled water is similar to that of rainfall, which
28
29 suggests limited kinetic fractionation in the aquifers used for commercial operations.
30
31 Latitude and longitude have an important influence on the isotopic signature of
32
33 groundwaters and together they explain 77 % of $\delta^{18}\text{O}$ variance.
34
35
36
37
38

39 **Disclosure statement**

40
41 No potential conflict of interest was reported by the authors.
42
43

44 **References**

- 45
46 [1] Debbeler LJ, Gamp M, Blumenschein M, et al. Polarized but illusory beliefs
47
48 about tap and bottled water: A product- and consumer-oriented survey and blind
49
50 tasting experiment. *Sci Total Environ.* 2018;643:1400–1410.
51
52 [2] Rosinger AY, Herrick KA, Wutich AY, et al. Disparities in plain, tap and
53
54 bottled water consumption among US adults: National Health and Nutrition
55
56 Examination Survey (NHANES) 2007–2014. *Public Health Nutr.*
57
58 2018;21(8):1455–1464.
59
60

- 1
2
3 [3] Cohen A, Ray I. The global risks of increasing reliance on bottled water. *Nat*
4 *Sustain.* 2018;1(7):327–329.
5
6 [4] Horowitz N, Frago J, Mu D. Life cycle assessment of bottled water: A case
7 study of Green2O products. *Waste Manage.* 2018;76:734–743.
8
9 [5] John Dennis M. Recent developments in food authentication. *Analyst.*
10 1998;123(9):151R–156R.
11
12 [6] Gremaud G, Quaile S, Piantini U, et al. Characterization of Swiss vineyards
13 using isotopic data in combination with trace elements and classical parameters.
14 *Eur Food Res Technol.* 2004;219(1):97–104.
15
16 [7] Carter JF, Yates HSA, Tinggi U. Stable isotope and chemical compositions of
17 European and Australasian ciders as a guide to authenticity. *J Agric Food Chem.*
18 2015;63(3):975–982.
19
20 [8] Carter JF, Yates HSA, Tinggi U. Isotopic and elemental composition of roasted
21 coffee as a guide to authenticity and origin. *J Agric Food Chem.*
22 2015;63(24):5771–5779.
23
24 [9] Raco B, Dotsika E, Cerrina Feroni A, et al. Stable isotope composition of Italian
25 bottled waters. *J Geochem Explor.* 2013;124:203–211.
26
27 [10] Brenčič M, Vreča P. The use of a finite mixture distribution model in bottled
28 water characterisation and authentication with stable hydrogen, oxygen and
29 carbon isotopes — Case study from Slovenia. *J Geochem Explor.*
30 2010;107(3):391–399.
31
32 [11] Rangarajan R, Ghosh P. Tracing the source of bottled water using stable isotope
33 techniques. *Rapid Commun Mass Spectrom.* 2011;25(21):3323–3330.
34
35 [12] Bowen GJ, Ehleringer JR, Chesson LA, et al. Stable isotope ratios of tap water
36 in the contiguous United States. *Water Resour Res.* 2007;43(3):W03419.
37
38 [13] Bong Y-S, Ryu J-S, Lee K-S. Characterizing the origins of bottled water on the
39 South Korean market using chemical and isotopic compositions. *Anal Chim*
40 *Acta.* 2009;631(2):189–195.
41
42 [14] Dotsika E, Poutoukis D, Raco B, et al. Stable isotope composition of Hellenic
43 bottled waters. *J Geochem Explor.* 2010;107(3):299–304.
44
45 [15] Godoy JM, Godoy mlDP, Neto A. Direct determination of $\delta(D)$ and $\delta(^{18}O)$ in
46 water samples using cavity ring down spectrometry: Application to bottled
47 mineral water. *J Geochem Explor.* 2012;119–120:1–5.
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 [16] Bowen GJ, Winter DA, Spero HJ, et al. Stable hydrogen and oxygen isotope
4 ratios of bottled waters of the world. *Rapid Commun Mass Spectrom*.
5 2005;19(23):3442–3450.
6
7
8 [17] Chesson LA, Bowen GJ, Ehleringer JR. Analysis of the hydrogen and oxygen
9 stable isotope ratios of beverage waters without prior water extraction using
10 isotope ratio infrared spectroscopy. *Rapid Commun Mass Spectrom*.
11 2010;24(21):3205–3213.
12
13 [18] British Soft Drink Association Annual Report 2020 [Internet]. London: BSDA;
14 [cited 19/1/2021]. Available from:
15 [https://www.britishsoftdrinks.com/write/MediaUploads/BSDA_Annual_Report_](https://www.britishsoftdrinks.com/write/MediaUploads/BSDA_Annual_Report_2020.pdf)
16 [2020.pdf](https://www.britishsoftdrinks.com/write/MediaUploads/BSDA_Annual_Report_2020.pdf)
17
18 [19] Brands of bottled mineral water ranked by number of users in the United
19 Kingdom (UK) in 2017 [Internet]. New York, NY: Statista; 2018 [cited
20 19/1/2021]. Available from: [https://www.statista.com/statistics/308644/leading-](https://www.statista.com/statistics/308644/leading-brands-of-bottled-mineral-water-in-the-uk/)
21 [brands-of-bottled-mineral-water-in-the-uk/](https://www.statista.com/statistics/308644/leading-brands-of-bottled-mineral-water-in-the-uk/).
22
23 [20] The GNIP Database [Internet]. Global Network of Isotopes in Precipitation;
24 2021 [cited 14/5/2021]. Accessible at <https://nucleus.iaea.org/wiser>
25
26 [21] Fuller L, Baker A, Fairchild IJ, et al. Isotope hydrology of dripwaters in a
27 Scottish cave and implications for stalagmite palaeoclimate research. *Hydrol*
28 *Earth Syst Sci*. 2008;12(4):1065–1074.
29
30 [22] Berman ESF, Levin NE, Landais A, et al. Measurement of $\delta^{18}\text{O}$, $\delta^{17}\text{O}$, and ^{17}O -
31 excess in water by off-axis integrated cavity output spectroscopy and isotope
32 ratio mass spectrometry. *Anal Chem*. 2013;85(21):10392–10398.
33
34 [23] Darling WG, Bath AH, Talbot JC. The O and H stable isotope composition of
35 freshwaters in the British Isles. 2. Surface waters and groundwater. *Hydrol Earth*
36 *Syst Sci*. 2003;7(2):183–195.
37
38 [24] Craig H. Isotopic variations in meteoric waters. *Science*. 1961;133(3465):1702–
39 1703.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Figure and table captions**
4

5
6 **Figure 1.** Source locations for bottled waters used in this study.
7

8
9 **Figure 2.** The influence of latitude and longitude on the $\delta^{18}\text{O}$ signature of UK bottled
10
11 waters.
12

13
14 **Figure 3.** Bottled water line (blue) compared with the UK meteoric water line (black)
15
16 and the global meteoric water line (dotted). The two red circles are bottled waters values
17
18 published in [16] from Trofarth Farm and Bethania.
19
20

21
22 **Table 1.** Mean $\delta^{18}\text{O}$ and $\delta^2\text{H}$ values for each bottled water brand. The isotopic values
23
24 from Trofarth Farm and Bethania are from [16].
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. Source locations for bottled waters used in this study.

543x393mm (600 x 600 DPI)




Figure 2. The influence of latitude and longitude on the $\delta^{18}\text{O}$ signature of UK bottled waters.

357x234mm (600 x 600 DPI)




Figure 3. Bottled water line (blue) compared with the UK meteoric water line (black) and the global meteoric water line (dotted). The two red circles are bottled waters values published in [16] from Trofath Farm and Bethania.

357x234mm (600 x 600 DPI)

Brand	$\delta^{18}\text{O}$ (‰ VSMOW)	St. dev.	$\delta^2\text{H}$ (‰ VSMOW)	St. dev.2	Source	Latitude	Longitude	Type
Buxton	-7.9	0.05	-49.2	0.15	Buxton	53.27611	-1.88269	natural mineral
Nestle Pure Life	-7.5	0.09	-48.4	0.01	Buxton	53.27611	-1.88269	spring
Harrogate	-7.9	0.05	-51.8	0.03	Harrogate	53.98752	-1.56378	spring
H2GO	-7.3	0.11	-49.9	0.08	Churchstoke	52.53732	-3.06382	natural mineral
Belu	-7.8	0.37	-52.0	0.27	Churchstoke	52.53732	-3.06382	natural mineral
Coop Still Water Churchstoke	-7.5	0.43	-49.9	0.07	Churchstoke	52.53732	-3.06382	natural mineral
Coop Still Water Chrucl Stretton	-7.7	0.29	-49.7	0.19	Church Stretton	52.538535	-2.780155	natural mineral
Actiph	-7.1	0.35	-48.2	0.16	Church Stretton	52.538535	-2.780155	spring
Morrisons Yorkshire Vale	-7.3	0.07	-49.3	0.63	Shepley	53.58572	-1.69939	spring
Morrisons Yorkshire Vale (Sparkling)	-7.5	0.27	-48.0	0.15	Shepley	53.58572	-1.69939	spring
ASDA Still	-7.4	0.14	-52.4	0.07	Armathwaite	54.76941	-2.77232	natural mineral
Tesco Ashbeck	-7.8	0.07	-52.9	0.11	Armathwaite	54.76941	-2.77232	natural mineral
Gill Beck	-8.1	0.11	-51.2	0.26	Armathwaite	54.76941	-2.77232	spring
Waitrose Essentials	-7.7	0.39	-53.1	0.37	Armathwaite	54.808933	-2.769192	natural mineral
Glaceau Smartwater	-8.1	0.09	-52.9	0.03	Morpeth	55.16291	-1.66982	spring
Sainsburys Still	-7.5	0.06	-47.5	0.22	Campsie Fells	55.97415	-4.20242	drinking
Highland Spring	-8.4	0.04	-55.0	0.09	Blackford	56.26061	-3.77901	spring
Highland Spring (Sparkling)	-8.2	0.30	-54.0	1.02	Blackford	56.26061	-3.77901	spring
ALDI Still	-7.7	0.10	-51.5	0.43	Lichfield	52.71995	-1.84596	spring
ALDI Aqueo (Sparkling)	-7.7	0.49	-50.9	1.14	Lichfield	52.71995	-1.84596	spring
Boots Brecon Carreg	-6.0	0.06	-36.3	0.19	Brecon Beacons	51.84418	-3.94599	natural mineral
Greggs Still	-5.9	0.03	-36.4	0.02	Brecon Beacons	51.84418	-3.94599	natural mineral
Wilkos Mountain Falls	-5.5	0.08	-33.8	0.10	Brecon Beacons	51.84418	-3.94599	spring
JUST Water	-7.2	0.07	-46.7	0.24	Ballymena	54.85856	-6.27936	spring
Princes Gate	-5.6	0.01	-32.1	0.16	Narberth	51.77257	-4.67273	natural mineral
M&S Mountain Water	-9.6	0.18	-61.0	0.52	Glenlivet Estate/Speyside	57.444736	-3.128582	drinking
M&S Mountain Water	-7.8	0.42	-49.2	0.26	Glorat Estate	55.974	-4.1785	drinking
Waitrose Royal Deeside	-8.6	0.28	-57.0	0.27	Pannanich Wells	57.057799	-3.000099	natural mineral
Decante	-6.7		-38.0		Trofarth Farm	53.21754	-3.71218	natural mineral

1						
2	Ty Nant	-6.5	-43.0	Bethania	52.24719	-4.10756 natural mineral
3						
4						
5						
6						
7						
8						
9						
10						
11						
12						
13						
14						
15						
16						
17						
18						
19						
20						
21						
22						
23						
24						
25						
26						
27						
28						
29						
30						
31						
32						
33						
34						
35						
36						
37						
38						
39						
40						
41						
42						
43						
44						
45						
46						

For Peer Review Only