

# Northumbria Research Link

Citation: Hume, Emily, Muse, Hazel, Wallace, Kirstie, Wilkinson, Mick, Marshall Heslop, Karen, Nair, Arun, Clark, Stephen and Vogiatzis, Ioannis (2022) Feasibility and Acceptability of a Physical Activity Behavioural Modification Tele-Coaching Intervention in Lung Transplant Recipients. *Chronic Respiratory Disease*, 19. p. 147997312211165. ISSN 1479-9723

Published by: SAGE

URL: <https://doi.org/10.1177/14799731221116588>  
<<https://doi.org/10.1177/14799731221116588>>

This version was downloaded from Northumbria Research Link:  
<https://nrl.northumbria.ac.uk/id/eprint/49569/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

# Feasibility and acceptability of a physical activity behavioural modification tele-coaching intervention in lung transplant recipients

Chronic Respiratory Disease  
Volume 19: 1–13  
© The Author(s) 2022  
Article reuse guidelines:  
[sagepub.com/journals-permissions](https://sagepub.com/journals-permissions)  
DOI: 10.1177/14799731221116588  
[journals.sagepub.com/home/crd](https://journals.sagepub.com/home/crd)  


Emily Hume<sup>1</sup> , Hazel Muse<sup>2</sup>, Kirstie Wallace<sup>2</sup>, Mick Wilkinson<sup>1</sup>, Karen Heslop Marshall<sup>2</sup>, Arun Nair<sup>2</sup>, Stephen Clark<sup>1,2</sup> and Ioannis Vogiatzis<sup>1</sup>

## Abstract

**Background:** Despite improvements in pulmonary function following lung transplantation (LTx), physical activity levels remain significantly lower than the general population. To date, there is little research investigating interventions to improve daily physical activity in LTx recipients. This study assessed the feasibility and acceptability of a novel, 12-weeks physical activity tele-coaching (TC) intervention in LTx recipients.

**Methods:** Lung transplant recipients within 2 months of hospital discharge were recruited and randomised (1:1) to TC or usual care (UC). TC consists of a pedometer and smartphone app, allowing transmission of activity data to a platform that provides feedback, activity goals, education, and contact with the researcher as required. Recruitment and retention, occurrence of adverse events, intervention acceptability and usage were used to assess feasibility.

**Results:** Key criteria for progressing to a larger study were met. Of the 15 patients eligible, 14 were recruited and randomised to TC or UC and 12 completed (67% male; mean  $\pm$  SD age;  $58 \pm 7$  years; COPD  $n = 4$ , ILD  $n = 6$ , CF  $n = 1$ , PH  $n = 1$ ): TC ( $n = 7$ ) and UC ( $n = 5$ ). TC was well accepted by patients, with 86% indicating that they enjoyed taking part. Usage of the pedometer was excellent, with all patients wearing it for over 90% of days and rating the pedometer and telephone contact as the most vital aspects. There were no adverse events related to the intervention. After 12 weeks, only TC displayed improvements in accelerometry steps/day (by  $3475 \pm 3422$ ;  $p = .036$ ) and movement intensity (by  $153 \pm 166$  VMU;  $p = .019$ ), whereas both TC and UC groups exhibited clinically important changes in physical SF-36 scores (by  $11 \pm 14$  and  $7 \pm 9$  points, respectively).

**Conclusion:** TC appears to be a feasible, safe, and well-accepted intervention in LTx.

## Keywords

Lung transplantation, tele-rehabilitation, physical activity

## Introduction

Lung transplantation (LTx) is an established final treatment option for those with end-stage lung disease. Over recent decades, survival rates have improved, with the International Society for Heart and Lung Transplantation Registry reporting a 5-years survival rate of 59%.<sup>1</sup> In addition to increasing survival, an important goal of LTx is to enhance

<sup>1</sup>Department of Sport, Exercise and Rehabilitation, Faculty of Health & Life Sciences, Northumbria University, Newcastle upon Tyne, UK

<sup>2</sup>The Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle upon Tyne, UK

### Corresponding author:

Emily Hume Department of Sport, Exercise and Rehabilitation, Faculty of Health and Life sciences, Northumbria University, Northumberland Building, Newcastle upon Tyne NE1 8ST, UK.

Email: [emily.c.hume@northumbria.ac.uk](mailto:emily.c.hume@northumbria.ac.uk)



Creative Commons CC BY: This article is distributed under the terms of the Creative Commons Attribution 4.0 License (<https://creativecommons.org/licenses/by/4.0/>) which permits any use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

health-related quality of life (HRQoL) and physical function.<sup>2</sup> Despite improvements in lung function, significant skeletal muscle weakness and reduced exercise capacity persist after LTx, which may limit improvements in daily physical functioning and HRQoL.<sup>3</sup> This is due to a host of factors including deconditioning as a result of persistent sedentary time, as well as immunosuppressant medications and episodes of organ rejection which may hinder functional recovery.<sup>4</sup> Several studies have shown that objectively measured physical activity is significantly reduced in LTx recipients.<sup>5–7</sup> Collectively, these data are concerning as physical activity is a strong predictor of all-cause mortality, both in patients with chronic respiratory disease and healthy individuals.<sup>8,9</sup>

To date, there is little research investigating interventions to improve daily physical activity in LTx recipients.<sup>2</sup> One RCT implementing a 12-weeks supervised exercise training programme, demonstrated significantly greater improvements in daily physical parameters compared to usual care.<sup>10</sup> Although exercise training in the form of pulmonary rehabilitation is recommended for LTx recipients,<sup>11</sup> access, uptake and completion of these programmes is limited in the UK<sup>12</sup> and worldwide.<sup>13</sup> With only six lung transplant centres across the UK, patients often live far away from the transplant centre,<sup>14</sup> therefore rehabilitation beyond the immediate post-transplant hospital phase is typically only undertaken by a small minority of patients who have a prolonged hospital stay, and this will vary depending on the patient's geographical location.

Physical activity tele-coaching is a digital intervention that aims to promote physical activity in COPD by facilitating behaviour change techniques such as individually tailored feedback, self-monitoring and goal setting.<sup>15,16</sup> However, LTx recipients experience significant deconditioning and psychological distress throughout their transplant journey and already have a high treatment burden, involving intensive medication regimes, self-monitoring, diet management and regular hospital appointments<sup>17,18</sup> Thus, it is not known whether physical activity tele-coaching will be feasible and improve outcomes in these patients. Therefore, the primary objectives of this study were to evaluate: (1) the proportion of LTx recipients accepting participation in the trial; (2) retention of LTx recipients; (3) feasibility of randomisation; (4) participants' acceptability of the TC intervention and (5) compliance with the intervention and physical activity goals. The secondary objectives were to explore and compare the tele-coaching intervention to usual care to obtain preliminary data on short-term clinical impact and safety of tele-coaching, by measuring physical activity, anxiety/depression and HRQoL outcomes, as well as rates of adverse events.

## Methods

### Ethics approval

This study received ethical approval from the Northeast, Tyne and Wear South Research Ethics Committee (REC Reference 19/NE/0119; IRAS project ID 257479) and was prospectively registered on the [clinicaltrials.gov](https://clinicaltrials.gov) database (NCT03873597).

### Study design

This study was a single centre, parallel two-arm, randomised controlled feasibility study. The trial consisted of three visits, which were all conducted remotely and included: a screening assessment (T0), a baseline assessment (T1) and a post-intervention assessment (12 weeks) (T2).

### Participants


Patients who had undergone single or bilateral LTx and were discharged between February 2020 and October 2021 were recruited from Freeman Hospital, Newcastle upon Tyne NHS Foundation Trust, UK. Potentially eligible patients were identified by designated cardiothoracic transplant co-ordinators, who provided initial information about the trial. Patients received an invitation letter with a participant information sheet and were given time to consider participation in the trial before written informed consent was obtained upon confirmation of eligibility. Patients were consented within 2 months following hospital discharge, to coincide with the first outpatient appointment.

### Inclusion criteria included

- Undergone single or bilateral LTx with a primary diagnosis of Interstitial Lung Disease (ILD), COPD, Cystic Fibrosis (CF), Bronchiectasis or Pulmonary Vascular Disease.
- Within 2 months of discharge following LTx
- Aged >18 years
- Able to speak and read English.
- Able to provide informed consent.

### Exclusion criteria included

- Severe post-transplant critical illness neuromyopathy
- Bilateral diaphragmatic weakness
- Presence of any other significant disease or disorder which, in the opinion of the investigators, may either put the participant at risk because of participation in the study, or may influence the result of the study, or the participant's ability to participate in the study.



**Figure 1.** Overview of physical activity behavioural modification tele-coaching intervention.

### Randomisation and concealment

Participants were assigned to one of two conditions using a computer-generated random sequence, managed by a researcher not involved in the recruitment process. Randomisation (1:1) was stratified by 6MWT distance (6MWD:  $<300$  or  $\geq 300$  m),<sup>19,20</sup> which was performed routinely before hospital discharge, using a block size of two following T1. The tele-coaching group received usual care in addition to the intervention. The control group received usual care, which included a motivational interview session. Given the nature of the intervention, it was not possible to conceal the treatment that participants were assigned to.

### Physical activity tele-coaching intervention

The 12-week physical activity behavioural modification tele-coaching intervention consisted of a: (1) motivational interview with a coach exploring motivational factors, barriers, preferred and non-preferred activities and strategies to become more active; (2) a pedometer (iChoice Shark A20, Choice MMed America Co., Bristol, PA) providing direct feedback; (3) smartphone app (Linkcare v2.7.1) which uses data collected from the pedometer, transmitted to a smart phone via Bluetooth and simultaneously to the Linkcare web-based platform; (4) home exercise booklet containing general strengthening and stretching exercises in three levels of difficulty and (5) telephone support from the researcher. An overview of the intervention is depicted in Figure 1.

Patients were asked to wear the pedometer during waking hours and interact with the smartphone application every day by reviewing and completing the automated application tasks. Every evening (after 8pm), patients were required to upload their step data to the smartphone application (via Bluetooth) by pressing the button on the pedometer. Each week an activity goal was set by the app, based on the patient's physical activity levels (steps/day) in the previous week.<sup>16</sup> The goals were calculated using the mean and median of the 4 most active days.<sup>21</sup> If the mean value exceeded the weekly goal, the application displayed the option to increase their median goal by 500 steps/day or to keep it the same as the previous week. If the mean value was lower than the weekly goal and the median was more than 500 steps/day below the goal, the goal was reduced to the median of the 4 most active days +500 steps/day.<sup>16</sup> Otherwise, the goal remained the same. The app also provided patients with daily feedback, encouragement, and educational messages, which were displayed in text or picture format. Throughout the intervention, researchers could access patient data via their app linked web-based platform (Linkcare app v2.7.1, Caldicott approval: 7372) and monitor their physical activity progress and adherence to the intervention. Telephone contact from the researcher was triggered if patients: (1) did not send their step count data for three consecutive days, (2) did not reach their step target for 2 consecutive weeks, (3) reached the step target but were not willing to increase their goal for 2 consecutive weeks. Prior to commencing the intervention, all patients

**Table 1.** Characteristics of patients at baseline (hospital discharge).

Characteristic	Tele-coaching (n = 7)	Usual care (n = 5)
Age (years)	57 ± 9	58 ± 4
BMI (kg/m <sup>2</sup> )	22.7 ± 3.9	25.3 ± 2.9
Sex (male/Female)	4/3	4/1
FEV <sub>1</sub> (L)	2.22 ± 0.52	2.15 ± 0.67
FEV <sub>1</sub> (% predicted)	70 ± 11	73 ± 21
FVC (% predicted)	66 ± 11	73 ± 27
FVC (L)	2.56 ± 0.74	2.75 ± 1.12
FEV <sub>1</sub> /FVC %	88 ± 7	81 ± 10
6MWD (m)	325 ± 69	336 ± 43
Diagnosis		
COPD	3	1
CF	1	0
ILD	2	4
PAH	1	0
Hospital length of stay (days)	45 ± 23	38 ± 4

BMI = Body mass index, COPD = Chronic Obstructive Pulmonary Disease, CF = Cystic Fibrosis, ILD = Interstitial Lung Disease, PAH = Pulmonary Arterial Hypertension. Values are mean ± SD.

received an instruction guide on how to use the smartphone application.

### Usual care

Usual care for LTx recipients included physical mobilisation whilst in the intensive care unit and post-transplant ward. During this time, patients were provided with a set of individualised rehabilitation exercises to conduct at home following hospital discharge. Additionally, as part of the study, participants assigned to usual care underwent a motivational interview to encourage patients to be physically active. This included education on the benefits of being physically active, goal setting and self-monitoring of physical activities.

### Outcomes to assess feasibility

A priori progression criteria were used to consider whether it would be appropriate to progress to a full-scale study. Based on other similar feasibility studies<sup>22-24</sup> these included: (1) feasibility to recruit participants, (2) retention of participants, (3) feasibility of randomisation processes, (4) intervention acceptability, and (5) intervention usage (Online supplement, Table 1).

#### Criterion 1: Screening, eligibility and recruitment

The screening rate was defined as the number of patients that were approached by the research team and assessed for eligibility against the inclusion and exclusion criteria. This included those who decided not to take part. Eligibility was

determined by dividing the number of people screened by the number who met the inclusion criteria.

The research team recorded all patients that met the eligibility criteria and decided not to take part in the trial, along with the reason for their decision.

#### Criterion 2: Retention

The retention rate was defined as the number of participants who remained in the study and did not drop out.

#### Criterion 3: Randomisation feasibility

Randomisation feasibility was assessed by the number of participants that were willing to be randomised to either the intervention or usual care group.

#### Criterion 4: Patient acceptability

Acceptability of the intervention by patients was assessed through a project specific questionnaire at T2,<sup>15</sup> consisting of 16 multiple choice questions on their experiences with the intervention, including 10-point Likert scales to rate the usefulness of the intervention components. Patients were asked to complete this 15-min questionnaire at T2.

#### Criterion 5: Actual usage of the intervention and step goal compliance

Actual usage of the pedometer throughout the intervention was assessed objectively using the data on the web based LinkCare Platform, specifically the pedometer readings on a

day-to-day basis. Usage of the pedometer was determined by the presence of step count data (>70 steps for that day),<sup>15,16</sup> to verify actual usage of the pedometer each day. Compliance with the step goal was assessed using the step data and goals set on the platform. Self-reported usage of the pedometer and home exercise booklet was also assessed within the acceptability questionnaire.

### Contact time

All contact with patients was recorded in a case file, including details on the duration and reason for each contact.

### Adverse events

An adverse event was defined as any untoward occurrence that occurred during the conduct of the study. All adverse events were recorded in the adverse event log within the patients notes and were classified as serious or not, and attributable to the study or not, as per the 'Decision Tree for Adverse Event reporting' from the National Institute for Health Research, Clinical Research Network, Introduction to Good Clinical Practice Toolkit<sup>25</sup>

### Outcomes to assess clinical effectiveness

**Physical activity.** Physical activity was assessed objectively using an Actigraph accelerometer (Actigraph LLC Pensacola, Florida, USA) in the week following T1 and the week following T2. This accelerometer has been previously validated in patients with COPD.<sup>26</sup> Patients in both the tele-coaching and usual care groups were instructed to wear the accelerometer for seven consecutive days during waking hours. The accelerometer was positioned using an elasticated waistband on the participant's dominant side on the iliac crest at the anterior axillary line. Prior to wearing the accelerometer, participants were given written instructions with a visual demonstration on: (1) the correct positioning of the device; (2) the start and end date of the physical activity assessment; (3) the wearing period (I.e. wear the device during waking hours); (4) when the device should be removed (I.e. during water based activities such as showering or bathing). A valid physical activity measurement was defined as a minimum of four weekdays, with at least 8 hours of wear time. Weekends were excluded from the analysis, in line with taskforce recommendations for COPD patients.<sup>27</sup> The physical activity parameters assessed included daily steps, movement intensity, time spent in sedentary and at least light activity intensities.

The pedometer was used by the intervention group as part of the tele-coaching intervention, to provide direct feedback to patients on their daily steps.

### Additional assessments

Additional outcomes assessed at T1 and T2 included: (1) HRQoL through the SF-36 questionnaire and (2) Anxiety and Depression using the Hospital Anxiety and Depression Scale (HADS).<sup>10</sup>

### Analyses

All statistical analyses were performed using SPSS version 27 (IBM, UK). Prior to analysis, the assumption of normality for outcomes was assessed using the Shapiro Wilk Test. Descriptive statistics were reported to better understand the distribution and potential for change of the proposed outcomes.

Data from the project-tailored questionnaire were scored as categorical variables and reported as frequencies and percentages (number of patients indicating each answer), except for the usefulness ratings of the components, which were expressed as medians (IQR). Actual usage of the pedometer was expressed as the percentage of patients who wore the pedometer for at least 90% of the days, as well as the median (IQR) wear time (days per week). The 90% cut off point was derived from a study utilising a similar intervention in COPD patients,<sup>15</sup> to allow comparison between studies. Weekly compliance to the goal was presented as the percentage of goals met over the intervention period (12 weeks).

The feasibility study was not powered to test the effectiveness hypotheses associated with any planned main large-scale trial. Paired t-tests or Wilcoxon Signed Ranks Test were employed to assess the within group differences from T1 to T2, to identify whether the intervention or natural recovery had a significant effect on physical activity outcomes. Independent samples t-tests or Mann-Whitney U tests were used to detect differences in change scores ( $\Delta$ ) between groups. Statistical significance was set at  $p < .05$  for all analyses.


## Results

### Participants

In total, 14 LTx recipients provided consent for the study and were randomised to the tele-coaching intervention ( $n = 8$ ) or usual care ( $n = 6$ ). Twelve patients completed T2 (Figure 2, Table 1).

### Criterion 1: Screening, eligibility and recruitment

A total of 26 LTx recipients were discharged between February 2020 and October 2021. Of those 26, four were unable to be approached, due to the suspension of trial recruitment at the start of the COVID-19 pandemic. In total,



**Figure 2.** CONSORT Participant Flow Diagram.

22 patients were screened by accessing patient records or by direct contact in clinic. Of the 22 patients screened, 7 (32%) were not eligible to participate in the trial. The remaining 15 patients received information about the trial (Figure 2).

In total, 14 LTx recipients were recruited between February 2020 and October 2021. No patients were recruited from March to October 2020, as well as mid-January to May 2021 due to the suspension of LTx in response to the COVID-19 pandemic.<sup>28</sup> The consent rate for the study was high at 93%, with 14 out of 15 patients accepting participation.

### Criterion 2: Retention

The retention rate was 86% for patients that consented to take part in the study. The dropout rates were equal between the tele-coaching and usual care group and the reasons for drop out were: (1) extenuating personal circumstances and (2) chronic lung allograft dysfunction resulting in palliative care.

### Criterion 3: Randomisation

All 14 patients were willing to be randomised to either the intervention or usual care group following T1.

### Criterion 4: Acceptability of intervention

Overall, patient feedback on the intervention was positive, with 86% of patients indicating that they either “liked” (29%) or “liked the intervention a lot” (57%) (Table 2, Figure 3). Furthermore, 86% of patients reported that the intervention “helped them a lot” to improve their physical activity levels, with 86% of patients indicating that the smartphone app was either “very easy” or “easy” to use. Importantly, 86% of patients were willing to use at least one aspect of the intervention in the future.

### Criterion 5: Actual usage of the intervention and step goal compliance

Of those completing the intervention, 100% wore the pedometer for more than 90% of days over the 12-weeks intervention. Overall, patients wore the pedometer for a median of 7 (IQR: 7–7) days per week.

In terms of self-reported usage, 86% of patients indicated that they looked at the pedometer “several times a day,” the remaining 14% indicated “once daily.”

The number of weekly step goal targets met throughout the 12-weeks intervention was good, with a mean (SD) of  $82 \pm 14\%$  of step goals achieved (Figure 3).


**Table 2.** Overview of patient responses from acceptability questionnaire.

Question 1) How much did you enjoy taking part in the activity programme?	Liked it a lot 57%	Liked it 29%	Neutral 14%	I disliked it 0%	No opinion 0%
Question 2) Did the intervention help you to increase your physical activity levels?	Yes, helped a lot 86%	Yes, helped a little 0%	Not noticeable 14%	No, not at all 0%	No, it discouraged me 0%
Question 3) How did you experience the weekly goal increases during the intervention?	Much too low 0%	A little bit too low 0%	Reasonable 86%	A little bit too high 14%	Much too high 0%
Question 4) How was it for you to work with the smartphone intervention?	Very easy 29%	Easy 57%	Not easy, but managed 0%	Difficult 14%	Very difficult 0%
Question 5) In your opinion, what was the most important part of the intervention?	Step counter 57%	Smartphone app 0%	Telephone contact 43%	Exercise booklet 0%	Other 0%
Question 6) How often did you perform the following actions? a) Look at the step counter b) Do any home exercises	Several times per day 86% 57%	Once per day 14% 14%	Sometimes but not everyday 0% 0%	Once or twice per week 0% 0%	Never 0% 29%
Question 7) How would you rate the graphics used on the smartphone application?	Very helpful and supportive 0%	Helpful and supportive 57%	Neutral 29%	Poor, not supportive 14%	Very poor, not supportive at all 0%
Question 8) How would you rate the interaction between you and the app?	Very quick 0%	Quick 43%	Neutral 43%	Slow 0%	Very slow 14%
Question 9) Which part of the intervention would you be willing to use in the future?	Nothing 14%	Step counter 14%	Step counter, phone and feedback messages 0%	Step counter, phone and contact 29%	Whole intervention 43%

### Contact time

The total mean  $\pm$  SD contact time per patient was 52  $\pm$  23 min per patient. On average, patients had to be

contacted 9  $\pm$  4 times over the 12-weeks. If the patient was progressing well and no contact was triggered, general well-being checks were conducted every 2 weeks via brief phone calls. For instances where the patient did not send their step



**Figure 3.** (A) pedometer steps/day, (B) step goal compliance and (C) boxplots depicting the usefulness scores (1-10 likert scale) of the different intervention components rated by patients.

data for three consecutive days, did not reach their step target for 2 consecutive weeks, or chose not to increase their goal for 2 consecutive weeks, the mean number of contacts was increased as well as the time for consultation. This was to provide troubleshooting solutions and explore barriers of engagement with goal adjustment.

### Adverse events

Over the study period, there were no adverse events related to the intervention and the effort of patients to progressively increase their activity levels, or related to the study protocol or procedures.

### Hospital admissions and complications

Throughout the 12-week intervention period, six patients (Tele-Coaching:  $n = 4$  and Usual Care:  $n = 2$ ) were admitted to the hospital for more than 72 h. In the tele-coaching group, the reasons for admission were acute rejection resulting in reduction in pulmonary function ( $n = 3$ ), fever and suspected infection ( $n = 1$ ) and dyspnoea due to right main bronchus anastomotic stricture ( $n = 1$ ). In the usual care group, hospital admissions were for acute rejection resulting in reduction in pulmonary function ( $n = 1$ ) and acute kidney injury ( $n = 1$ ).

### Outcome measures

**Accelerometer-derived physical activity.** At 12 weeks there were clinically important<sup>29</sup> improvements in steps/day for both the tele-coaching (by  $3475 \pm 3422$  steps/day;  $p = .036$ ) and usual care (by  $1159 \pm 991$  steps/day;  $p = .059$ ) groups, however the increase in the tele-coaching group exceeded the usual care group by clinically important margins<sup>29</sup> (by 2316 steps/day) (Table 3).

Accelerometer movement intensity significantly improved within the tele-coaching group (by  $153 \pm 166$  VMU;

$p = .019$ ), but not the usual care group (by  $-3 \pm 61$  VMU;  $p = .908$ ), with a significant difference between groups (by 156 VMU;  $p = .037$ ). For time spent in at least light activity, there was a significant increase within the tele-coaching group (by  $37 \pm 24$  min/day;  $p = .006$ ) at 12 weeks, but not in the usual care group (by  $3 \pm 37$  min/day;  $p = .861$ ), with a significant difference between groups (by 34 min/day;  $p = .040$ ). Individual changes in steps/day and movement intensity for each disease entity in the tele-coaching and usual care groups are presented in Figure 4. For daily steps and movement intensity the mean improvement in the tele-coaching group was  $3896 \pm 5580$  steps/day and  $127 \pm 175$  VMU, respectively for ILD ( $n = 2$ ),  $2126 \pm 919$  steps/day and  $100 \pm 52$  VMU, respectively for COPD ( $n = 3$ ),  $8717 \pm 0$  steps/day and  $479 \pm 0$  VMU, respectively for Pulmonary Arterial Hypertension (PAH) ( $n = 1$ ) and  $1438 \pm 0$  steps/day and  $39 \pm 0$  VMU, respectively for CF ( $n = 1$ ). For usual care, the mean improvement in daily steps and movement intensity were  $1329 \pm 1057$  steps/day and  $17.8 \pm 44.7$  VMU, respectively for ILD ( $n = 4$ ) and  $479 \pm 0$  steps/day and  $-87.9 \pm 0$  VMU, respectively for COPD ( $n = 1$ ).

**HRQoL and psychological wellbeing.** At 12 weeks, there were clinically important ( $>2-3$  units<sup>30</sup>) increases in SF-36 physical component summary scores, in both the tele-coaching (+11 points) and usual care (+7 points) groups, however these changes were not statistically significant between groups.


### Discussion

This study showed that tele-coaching was feasible, safe, and well accepted by LTx recipients. Patient uptake and retention, acceptability and usage of the tele-coaching intervention was high, without occurrence of adverse events. When compared to usual care, tele-coaching elicited improvements in accelerometer derived physical activity

**Table 3.** Changes in PA parameters and HRQoL outcomes in the Tele-Coaching and Usual Care groups.

Outcome	Group	Baseline (T1)	12 weeks (T2)	Within group change	Between group p value
<b>Accelerometer Outcomes</b>					
Daily steps (steps/day)	TC	3558 ± 3188	7033 ± 5944	3475 ± 3422 <sup>##</sup>	0.089 <sup>#</sup>
	UC	4249 ± 3531	5408 ± 4444	1159 ± 991 <sup>#</sup>	
Movement intensity (VMU)	TC	237 ± 155	390 ± 311	153 ± 166 <sup>*</sup>	0.037
	UC	317 ± 153	314 ± 129	-3 ± 61	
Time spent in sedentary activity (min/day)	TC	570 ± 108	513 ± 296	-57 ± 128	0.114
	UC	441 ± 59	463 ± 85	22 ± 52	
Time spent in at least light activity (min/day)	TC	160 ± 58	197 ± 72	37 ± 24 <sup>*</sup>	0.040
	UC	191 ± 71	194 ± 58	3 ± 37	
<b>HADS</b>					
Anxiety	TC	5 ± 4	4 ± 4	1 ± 4	0.339
	UC	3 ± 4	4 ± 5	1 ± 2	
Depression	TC	3 ± 4	4 ± 4	0 ± 5	0.479
	UC	3 ± 3	3 ± 5	0 ± 2	
<b>SF-36</b>					
PCS Score	TC	28 ± 11	39 ± 14	11 ± 14 <sup>#</sup>	0.291
	UC	30 ± 7	37 ± 12	7 ± 9 <sup>#</sup>	
MCS score	TC	57 ± 7	53 ± 14	-4 ± 15	0.348
	UC	51 ± 14	51 ± 13	-1 ± 10	

Abbreviations: VMU = Vector Magnitude Units, MVPA = moderate to vigorous physical activity, HADS = Hospital Anxiety and Depression Scale, SF-36 = Short Form 36 Questionnaire, PCS = Physical Component Summary, MCS = Mental Component Summary, TC = Tele-Coaching, UC = Usual Care. Values are mean ± SD \* = statistically significant (P<0.05), # = clinically important change.



**Figure 4.** Individual changes in steps/day and movement intensity (VMU) in the tele-coaching (A&C) and usual care (B&D) groups from baseline to 12 weeks.

parameters that exceeded clinically important margins, highlighting the potential efficacy of this intervention to support patients post LTx.

Recruitment for the trial was significantly affected by the COVID-19 pandemic and the suspension of LTx during the early stages of the pandemic. Thus, the main reason for slow patient recruitment was due to the limited number of transplants performed. A centre-specific investigation reported this as a 77% reduction during the first peak of the pandemic.<sup>28</sup> Although the number of transplants was limited, uptake of the study was high with 93% of eligible participants accepting participation. This well exceeds criteria previously used to proceed to a full-scale trial (>30% of eligible patients recruited).<sup>22</sup> Additionally, there were low rates of attrition in both the tele-coaching and usual care groups (14% overall) over the 12-weeks. According to previous literature, attrition of <20% is unlikely to threaten the validity of a trial.<sup>31</sup> Additionally, this is significantly lower than the dropout rate previously reported in a meta-analysis of app-based interventions in chronic disease.<sup>32</sup>

Overall, the tele-coaching intervention was well accepted by patients, who rated their enjoyment similarly to a study using the same intervention in COPD patients.<sup>15</sup> Most patients (86%) reported that the intervention ‘helped them a lot’ to improve their physical activity, which is higher than that previously reported in COPD patients (59%).<sup>15</sup> The simplicity of the smartphone application may have contributed to the good acceptability of the intervention, as most patients reported finding it easy to use. In COPD patients, 47.8% rated the goal increases as either ‘high’ or ‘much too high’ compared to only 14% in the current study in LTx recipients, which is supported by high step goal compliance ( $82 \pm 14\%$ ).<sup>15</sup> This may suggest that LTx recipients are more ambitious with their physical activity targets, because of improved lung function and diminished symptoms of breathlessness.<sup>33</sup>

Alike to the findings in COPD patients,<sup>15</sup> LTx recipients considered the pedometer and telephone contact with the researcher as the most important components of the intervention. The regular contact with the researcher to resolve and advise on any safety concerns in the current study, may have enhanced patient’s self-efficacy to undertake more physical activity,<sup>34,35</sup> and highlights the significance of a collaborative approach between the patient and healthcare professional (HCP) in facilitating patient behaviour change and self-management.<sup>36</sup>

Although HCP contact was important, the average contact time required for each patient was only 52 min over the 12-weeks intervention. In the current study, coaching eight patients simultaneously over 12 weeks, would equate to around 35 min of HCP time per week, which is significantly less resource intensive than pulmonary rehabilitation. The low contact time could have been facilitated by several factors, such as the semi-automated nature of the

intervention, the instruction booklet provided to help with working the app, as well as the simplicity of the app, as 86% of patients indicated that they found the app either “very easy” or “easy” to use.

The high-level of perceived importance of the pedometer by patients was also reflected by the excellent actual and self-reported usage of the pedometer. All patients wore the pedometer for over 90% of the 12-weeks programme, which was higher than that previously reported in the study in COPD patients (59%).<sup>15</sup>

In terms of accelerometer physical activity outcomes, there were clinically important improvements in steps/day in both groups. The improvement in the usual care group highlights the natural recovery occurring in the early stages of LTx recovery, similarly to Langer et al.<sup>10</sup> who demonstrated an improvement of 750 steps/day in a usual care group within an exercise training study. A recent systematic review<sup>37</sup> highlighted that the majority of rehabilitation studies conducted post-transplantation are limited by the lack of a control group, making it difficult to differentiate the true effect of the intervention. Literature on interventions to improve physical activity in LTx recipients is scarce.<sup>2</sup> Improvements in daily steps in the current study exceeded those shown following exercise training.<sup>10,38</sup> This is likely due to step counts being the central focus of the intervention and the incorporation of behavioural techniques such as self-monitoring, goal setting and feedback, which have been deemed important for enhancing healthy activity behaviours.<sup>39</sup>

Although peripheral muscle abnormalities have been shown to be the predominant limiting factor to exercise capacity in lung transplant recipients,<sup>3</sup> the underlying lung disease entity and pathophysiology may also influence an individual’s exercise capacity and physical activity behaviour. When examining individual changes following tele-coaching (Figure 4), the largest improvements in daily steps and movement intensity were seen in PAH, whereas the lowest was in CF. Although CF recipients are often younger compared to other disease entities such as COPD, CF is a multi-organ disease in which co-morbidities such as diabetes mellitus and bone disease are common both pre- and post-transplant.<sup>40</sup> It is important to note that no conclusions can be drawn from this data due to the limited sample size, however this poses an interesting question for future research. Our findings on HRQoL support those of Finlen Copeland et al.,<sup>41</sup> showing that SF-36 PCS scores demonstrate a natural course of recovery following LTx, likely due to improved pulmonary function, symptoms and ability to perform daily activities. The lack of change in HADs and SF-36 MCS scores in either group reflects previous findings.<sup>10,42</sup> This may be due to several factors such as the uncertainty of organ rejection, adverse effects of immunosuppressive medications and recurring pain following LTx. However, a full

scale RCT is required to infer whether tele-coaching can induce greater improvements in HRQoL outcomes than usual care alone.

### Implications of the study

This study highlights the potential of digital health technology to increase physical activity levels in LTx recipients in the early stages of recovery. The present study may inform a fully powered RCT to determine whether a digital physical activity intervention can elicit significantly greater improvements in physical activity and HRQoL outcomes compared to usual care, as well as determine the longer-term impact of this intervention.

### Study limitations

There are several limitations that must be considered in this study. Firstly, this was a small-scale study, therefore, generalisability of the results to clinical practice may be limited. However, the main aim of this study was to explore the feasibility and acceptability of tele-coaching in LTx recipients, thus it was not powered to detect differences in study outcomes between groups. Secondly, acceptability of the intervention was assessed through a project specific questionnaire which was used previously by Loeckx et al.<sup>15</sup> in COPD patients. This makes it challenging to make comparisons with other studies implementing digital health interventions, however it provides useful insights into patient acceptability and can be compared to the findings by Loeckx et al.<sup>15</sup> to explore differences between different patient groups using the same intervention. Finally, randomisation to the tele-coaching and usual care groups was stratified based on functional exercise capacity (6MWD) as this has been demonstrated as a strong predictor of physical activity change,<sup>43</sup> consequently it was not possible to balance groups for all variables (e.g. sex and disease entities) and there was a large diversity of primary disease diagnosis and therefore underlying pathophysiology of physical activity limitation.<sup>44</sup>

### Conclusion

In conclusion, physical activity tele-coaching appears to be a feasible, safe, and acceptable intervention to support patients post LTx. Additionally, there is a degree of natural recovery in some physical activity and HRQoL parameters, however tele-coaching appears to elicit greater improvements in physical activity measures.

### Acknowledgements

The authors would like to thank the lung transplant recipients who took part in this study and wish to acknowledge the following

persons for their complementary support: Linkcare Health Services SL.: Jim Roldan, Jose Sanchez and Josep Benavent.

### Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

### Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The study was funded by the Freeman Heart and Lung Transplant Association (grant number: 7417FH), the Transplant Association (grant number: 1161340) and Northumbria University collaborative doctoral research studentship scheme (reference number: RST00428).

### ORCID iD

Emily Hume  <https://orcid.org/0000-0003-0462-2395>

### Supplemental Material

Supplemental material for this article is available online.

### References

1. Bos S, Vos R, Van Raemdonck DE, et al. Survival in adult lung transplantation: where are we in 2020? *Curr Opin Organ Transpl* 2020; 25: 268–273. DOI: [10.1097/mot.0000000000000753](https://doi.org/10.1097/mot.0000000000000753).
2. Langer D. Addressing the changing rehabilitation needs of patients undergoing thoracic surgery. *Chronic Respir Dis* 2021; 18: 1479973121994783. DOI: [10.1177/1479973121994783](https://doi.org/10.1177/1479973121994783).
3. Mathur S, Reid WD, Levy RD, et al. Exercise limitation in recipients of lung transplants. *Phys Ther* 2004; 84: 1178–1187. DOI: [10.1093/ptj/84.12.1178](https://doi.org/10.1093/ptj/84.12.1178).
4. Langer D. Rehabilitation in patients before and after lung transplantation. *Respiration; International Review Thoracic Diseases* 2015; 89: 353–362. DOI: [10.1159/000430451](https://doi.org/10.1159/000430451).
5. Wickerson L, Mathur S, Singer LG, et al. Physical activity levels early after lung transplantation. *Phys Ther* 2015; 95: 517–525. DOI: [10.2522/ptj.20140173](https://doi.org/10.2522/ptj.20140173).
6. Ulvestad M, Durheim MT, Kongerud JS, et al. Cardiopulmonary fitness and physical activity following lung transplantation: a national cohort study. *Respiration; International Review Thoracic Diseases* 2020; 99: 316–324. DOI: [10.1159/000506883](https://doi.org/10.1159/000506883).
7. Langer D, Gosselink R, Pitta F, et al. Physical activity in daily life 1 year after lung transplantation. *J Heart Lung Transplantation: The Official Publication Int Soc Heart Transplant* 2009; 28: 572–578. DOI: [10.1016/j.healun.2009.03.007](https://doi.org/10.1016/j.healun.2009.03.007).
8. Garcia-Aymerich J, Lange P, Benet M, et al. Regular physical activity reduces hospital admission and mortality in chronic obstructive pulmonary disease: a population based cohort

- study. *Thorax* 2006; 61: 772–778. DOI: [10.1136/thx.2006.060145](https://doi.org/10.1136/thx.2006.060145).
9. Lee IM and Skerrett PJ. Physical activity and all-cause mortality: what is the dose-response relation? *Med Sci Sports Exerc* 2001; 33: S459–S471.
  10. Langer D, Burtin C, Schepers L, et al. Exercise training after lung transplantation improves participation in daily activity: a randomized controlled trial. *Am Journal Transplantation* 2012; 12: 1584–1592. DOI: [10.1111/j.1600-6143.2012.04000.x](https://doi.org/10.1111/j.1600-6143.2012.04000.x).
  11. Spruit MA. Pulmonary rehabilitation. *Eur Respir Rev* 2014; 23: 55–63. DOI: [10.1183/09059180.00008013](https://doi.org/10.1183/09059180.00008013).
  12. Steiner M, Holzhauer-Barrie J, Lowe D, et al. *Pulmonary rehabilitation: steps to breathe better/National chronic obstructive pulmonary disease (COPD) audit programme: clinical audit of pulmonary rehabilitation services in england and wales national clinical audit report*. 2016. London: RCP.
  13. Rochester CL, Vogiatzis I, Holland AE, et al. An official American thoracic society/European respiratory society policy statement: enhancing implementation, use, and delivery of pulmonary rehabilitation. *Am J Respir Crit Care Med* 2015; 192: 1373–1386. DOI: [10.1164/rccm.201510-1966ST](https://doi.org/10.1164/rccm.201510-1966ST).
  14. NHS Blood and Transplant. *Annual report on cardiothoracic organ transplantation 2020/2021*, 2021.
  15. Loeckx M, Rabinovich RA, Demeyer H, et al. Smartphone-based physical activity telecoaching in chronic obstructive pulmonary disease: mixed-methods study on patient experiences and lessons for implementation. *JMIR Mhealth Uhealth* 2018; 6: e200.
  16. Demeyer H, Louvaris Z, Frei A, et al. Physical activity is increased by a 12-week semiautomated telecoaching programme in patients with COPD: a multicentre randomised controlled trial. *Thorax* 2017; 72: 415–423. DOI: [10.1136/thoraxjnl-2016-209026](https://doi.org/10.1136/thoraxjnl-2016-209026).
  17. Xu J, Adeboyejo O, Wagley E, et al. Daily burdens of recipients and family caregivers after lung transplant. *Prog Transpl* 2012; 22: 41–48. DOI: [10.7182/pit2012815](https://doi.org/10.7182/pit2012815).
  18. Wessels-Bakker MJ, van de Graaf EA, Kwakkel-van Erp JM, et al. The relation between psychological distress and medication adherence in lung transplant candidates and recipients: a cross-sectional study. *J Clin Nurs* 2021; 31: 716–725. n/a DOI: [10.1111/jocn.15931](https://doi.org/10.1111/jocn.15931).
  19. van Adrichem EJ, Reinsma GD, van den Berg S, et al. Predicting 6-minute walking distance in recipients of lung transplantation: longitudinal study of 108 patients. *Phys Ther* 2015; 95: 720–729. DOI: [10.2522/ptj.20140001](https://doi.org/10.2522/ptj.20140001).
  20. Maury G, Langer D, Verleden G, et al. Skeletal muscle force and functional exercise tolerance before and after lung transplantation: a cohort study. *Am Journal Transplantation* 2008; 8: 1275–1281. DOI: [10.1111/j.1600-6143.2008.02209.x](https://doi.org/10.1111/j.1600-6143.2008.02209.x).
  21. Demeyer H, Burtin C, Van Remoortel H, et al. Standardizing the analysis of physical activity in patients with COPD following a pulmonary rehabilitation program. *Chest* 2014; 146: 318–327. DOI: [10.1378/chest.13-1968](https://doi.org/10.1378/chest.13-1968).
  22. Hawkins J, Charles JM, Edwards M, et al. Acceptability and feasibility of implementing accelerometry-based activity monitors and a linked web portal in an exercise referral scheme: feasibility randomized controlled trial. *J Med Internet Res* 2019; 21: e12374.
  23. Haines M. Feasibility of procedures for a randomised pilot study of reduced exertion, high-intensity interval training (REHIT) with non-diabetic hyperglycaemia patients. *Pilot Feasibility Stud* 2020; 6: 28. DOI: [10.1186/s40814-020-00571-8](https://doi.org/10.1186/s40814-020-00571-8).
  24. Ward N, Stiller K, Rowe H, et al. Airway clearance by exercising in mild cystic fibrosis (ACE-CF): a feasibility study. *Respir Med* 2018; 142: 23–28. DOI: [10.1016/j.rmed.2018.07.008](https://doi.org/10.1016/j.rmed.2018.07.008).
  25. National Institute for Health Research. *Introduction to good clinical practice elearning (secondary care)*, 2018, <https://learn.nihr.ac.uk/course/view.php?id=282>
  26. Rabinovich RA, Louvaris Z, Raste Y, et al. Validity of physical activity monitors during daily life in patients with COPD. *Eur Respir J* 2013; 42: 1205–1215. DOI: [10.1183/09031936.00134312](https://doi.org/10.1183/09031936.00134312).
  27. Demeyer H, Mohan D, Burtin C, et al. Objectively measured physical activity in patients with COPD: recommendations from an international task force on physical activity. *Chronic Obstr Pulm Dis* 2021; 8: 528–550. DOI: [10.15326/jcopdf.2021.0213](https://doi.org/10.15326/jcopdf.2021.0213).
  28. Hardman G, Sutcliffe R, Hogg R, et al. The impact of the SARS-CoV-2 pandemic and COVID-19 on lung transplantation in the UK: lessons learned from the first wave. *Clin Transplant* 2021; 35: e14210, DOI: [10.1111/ctr.14210](https://doi.org/10.1111/ctr.14210).
  29. Demeyer H, Burtin C, Hornikx M, et al. The minimal important difference in physical activity in patients with COPD. *PLoS One* 2016; 11: e0154587. DOI: [10.1371/journal.pone.0154587](https://doi.org/10.1371/journal.pone.0154587).
  30. Singer LG and Chowdhury N. Estimation of minimal important differences in health-related quality of life measures for lung transplant recipients. *J Heart Lung Transplant* 2013; 32: S214–S215, DOI: [10.1016/j.healun.2013.01.539](https://doi.org/10.1016/j.healun.2013.01.539).
  31. Schulz KF and Grimes DA. Sample size slippages in randomised trials: exclusions and the lost and wayward. *Lancet* 2002; 359: 781–785. DOI: [10.1016/s0140-6736\(02\)07882-0](https://doi.org/10.1016/s0140-6736(02)07882-0).
  32. Meyerowitz-Katz G, Ravi S, Arnolda L, et al. Rates of attrition and dropout in app-based interventions for chronic disease: systematic review and meta-analysis. *J Med Internet Res* 2020; 22: e20283. DOI: [10.2196/20283](https://doi.org/10.2196/20283).
  33. Pêgo-Fernandes PM, Abrão FC, Fernandes FLA, et al. Spirometric assessment of lung transplant patients: one year follow-up. *Clinics (Sao Paulo)* 2009; 64: 519–525. DOI: [10.1590/s1807-59322009000600006](https://doi.org/10.1590/s1807-59322009000600006).
  34. Hartman JE, Boezen HM, de Greef MH, et al. Consequences of physical inactivity in chronic obstructive pulmonary

- disease. *Expert Rev Respir Med* 2010; 4: 735–745. DOI: [10.1586/ers.10.76](https://doi.org/10.1586/ers.10.76).
35. McAuley E, Szabo A, Gothe N, et al. Self-efficacy: implications for physical activity, function, and functional limitations in older adults. *Am J Lifestyle Med* 2011; 5: 361–369. DOI: [10.1177/1559827610392704](https://doi.org/10.1177/1559827610392704).
36. Benzo R. Collaborative self-management in chronic obstructive pulmonary disease: learning ways to promote patient motivation and behavioral change. *Chronic Respir Dis* 2012; 9: 257–258. DOI: [10.1177/1479972312458683](https://doi.org/10.1177/1479972312458683).
37. Hume E, Ward L, Wilkinson M, et al. Exercise training for lung transplant candidates and recipients: a systematic review. *Eur Respir Rev* 2020; 29: 200053. DOI: [10.1183/16000617.0053-2020](https://doi.org/10.1183/16000617.0053-2020).
38. Choi J, Hergenroeder AL, Burke L, et al. Delivering an in-home exercise program via telerehabilitation: a pilot study of lung transplant go (LTGO). *Int Journal Telerehabilitation* 2016; 8: 15–26. DOI: [10.5195/ijt.2016.6201](https://doi.org/10.5195/ijt.2016.6201).
39. Sullivan AN and Lachman ME. Behavior change with fitness technology in sedentary adults: a review of the evidence for increasing physical activity. *Front Public Health* 2017; 4: 289–289. DOI: [10.3389/fpubh.2016.00289](https://doi.org/10.3389/fpubh.2016.00289).
40. Meachery G, De Soyza A, Nicholson A, et al. Outcomes of lung transplantation for cystic fibrosis in a large UK cohort. *Thorax* 2008; 63: 725–731. DOI: [10.1136/thx.2007.092056](https://doi.org/10.1136/thx.2007.092056).
41. Finlen Copeland CA, Vock DM, Pieper K, et al. Impact of lung transplantation on recipient quality of life: a serial, prospective, multicenter analysis through the first post-transplant year. *Chest* 2013; 143: 744–750. DOI: [10.1378/chest.12-0971](https://doi.org/10.1378/chest.12-0971).
42. Singer JP, Chen J, Blanc PD, et al. A thematic analysis of quality of life in lung transplant: the existing evidence and implications for future directions. *Am Journal Transplantation* 2013; 13: 839–850. DOI: [10.1111/ajt.12174](https://doi.org/10.1111/ajt.12174).
43. Osadnik CR, Loeckx M, Louvaris Z, et al. The likelihood of improving physical activity after pulmonary rehabilitation is increased in patients with COPD who have better exercise tolerance. *Int Journal Chronic Obstructive Pulmonary Disease* 2018; 13: 3515–3527. DOI: [10.2147/COPD.S174827](https://doi.org/10.2147/COPD.S174827).
44. Vogiatzis I, Zakynthinos G, Andrianopoulos V, et al. Mechanisms of physical activity limitation in chronic lung diseases. *Pulm Med* 2012; 2012: 634761–634811. DOI: [10.1155/2012/634761](https://doi.org/10.1155/2012/634761).