# Northumbria Research Link

Citation: Travis, Eleanor, Scott-Bell, Andrea and Thornton, Claire (2024) The current state of concussion knowledge and attitudes in British American Football. Physician and Sportsmedicine, 52 (1). pp. 36-45. ISSN 0091-3847

Published by: Taylor & Francis

URL: https://doi.org/10.1080/00913847.2022.2156766 <https://doi.org/10.1080/00913847.2022.2156766>

This version was downloaded from Northumbria Research Link: https://nrl.northumbria.ac.uk/id/eprint/50943/

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <a href="http://nrl.northumbria.ac.uk/policies.html">http://nrl.northumbria.ac.uk/policies.html</a>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)







The Physician and Sportsmedicine

ISSN: (Print) (Online) Journal homepage: https://www.tandfonline.com/loi/ipsm20

### The current state of concussion knowledge and attitudes in British American Football

Eleanor Travis, Dr Andrea Scott-Bell & Dr Claire Thornton

To cite this article: Eleanor Travis, Dr Andrea Scott-Bell & Dr Claire Thornton (2024) The current state of concussion knowledge and attitudes in British American Football, The Physician and Sportsmedicine, 52:1, 36-45, DOI: 10.1080/00913847.2022.2156766

To link to this article: https://doi.org/10.1080/00913847.2022.2156766

© 2022 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



6

Published online: 15 Dec 2022.

_	
С	
	1.
L	<u>v</u>
_	

Submit your article to this journal 🗹

Article views: 1381



View related articles

View Crossmark data 🗹



Citing articles: 1 View citing articles 🗹

#### ORIGINAL RESEARCH

OPEN ACCESS OPEN ACCESS

Taylor & Francis

Taylor & Francis Group

## The current state of concussion knowledge and attitudes in British American Football

Eleanor Travis<sup>a</sup>, Dr Andrea Scott-Bell <sup>b</sup> and Dr Claire Thornton <sup>b</sup>

<sup>a</sup>Musculoskeletal Health Research Group, School of Health, Leeds Beckett University, Leeds, UK; <sup>b</sup>Department of Sport, Exercise and Rehabilitation, Northumbria University, Newcastle upon Tyne, UK

#### ABSTRACT

**Objectives:** To examine concussion knowledge and concussion attitudes of players, coaches, and support staff in British American Football (BAF).

**Methods:** Data from players, coaches and support staff (n = 236) were collected from across all leagues in BAF. An online survey tool was used which included the Rosenbaum Concussion Knowledge and Attitudes Survey (RoCKAS), and questions examining concussion education and perceived risk of participating in football.

**Results:** The mean score on the RoCKAS concussion knowledge was  $21.0 \pm 2.1$  of a possible score of 25 reflecting good knowledge. Of a possible score of 65, the mean concussion attitude score was  $55.6 \pm 6.1$  showing safe attitude. Whilst an overall safe attitude was seen, almost half of participants (45.3%) noted they would continue to play with a concussion. No relationship was found between CAI and prior concussion history. Fifty seven percent of participants agreed the benefits of playing football outweighed the risks. Forty eight percent reported that they had received no concussion-related education in the past 12 months.

**Conclusion:** BAF participants have good concussion knowledge and safe attitudes. However, risky behavior is demonstrated through unsafe likelihood to report and attitude to long-term health risks. Access to the British American Football Association (BAFA) concussion policy and education was poor raising questions over what sources of information stakeholders are drawing their knowledge from. These findings can help form the foundation of educational interventions (e.g. coaching workshops) to challenge current misconceptions and improve likelihood to report concussion in BAF.

**ARTICLE HISTORY** 

Received 30 August 2022 Accepted 4 December 2022

#### **KEYWORDS**

Concussion reporting; british american football; health education; injury risk; concussion attitude; concussion knowledge

#### Introduction

American football (AF) is a growing sport in the United Kingdom with 473 National League teams currently playing under the national governing body, the British American Football Association (BAFA) in both the flag and tackle game [1]. In the 2020–21 season there were 9075 players across BAFA flag and tackle leagues, with possible age ranges from 7 years to adult (no upper age limit). Due to the tackling and blocking element of this sport, a significant number (59.3%) of injuries including concussion-injuries are caused by player contact [2]. In 2020, 32% of athletes in the British university game of AF were at risk of sustaining at least one concussion over the course a season [3]. However, it should be noted that these findings were in university AF players only and at present there is no data on non-self reported diagnosed concussions.

Appropriate and timely recognition of concussion is important to the management of the injury, to provide both immediate and secondary stage care, in order to avoid longterm tertiary effects. Unfortunately, under-reporting of concussion is common within BAF, where almost 45% of players suspected they had a concussion but did not get this formally diagnosed [4]. Furthermore, when asked about concussion injury reporting behavior 23.5% of players reported hiding symptoms from coaches or medical staff [4]. Findings from this study reported players would downplay, ignore and deny injury [4]. This greatly increases the risk of subsequent acute or chronic illness (e.g. secondary impact syndrome [5], chronic traumatic encephalopathy or 'traumatic encephalopathy syndromes' (TESs)) [6,7]. Worryingly, a history of concussions puts the athlete at elevated risk of further concussions which are likely more severe and have a longer recovery period [8–11].

As a developing sport BAF has finite resources with many teams having limited access to suitably trained medical personnel and updated concussion protocols [4]. Thus, there is a greater need for players and game day staff (e.g. coaches, players and referees) to recognize signs and symptoms of concussion and for players to self-report suspected concussions. Coaches have gaps in knowledge regarding concussion, particularly around diagnostic methods and recovery of youth athletes [12]. This therefore needs to be addressed to ensure player safety.

The aim of this study was to investigate the knowledge and attitudes of concussion in players, coaches and staff involved in the game of BAF.

Investigation of the knowledge and attitudes of concussion in BAF is a key step to understanding the reasons behind the

CONTACT Eleanor Travis 🖾 e.travis@leedsbeckett.ac.uk 🖃 School of Health, Leeds Beckett University, City Campus, Leeds LS6 3QW, UK

© 2022 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The terms on which this article has been published allow the posting of the Accepted Manuscript in a repository by the author(s) or with their consent.

under-reporting of injury and to help guide the creation of future educational strategies of the national governing body for all parties involved within the game.

#### **Materials and methods**

#### **Participants**

Participants (n = 236) were drawn from across all leagues in British American Football in the UK during the autumn of 2021. They were contacted via central BAFA e-mail communication (to registered coaches, players, and support staff, those who had 'opted-in' to e-mail communication) and social media (Facebook and Instagram). The demographic sample can be found in Table 1.

Inclusion criteria: Participants had to be over the age of 18 and actively involved in British American Football as a coach, player, support personnel or combination of roles in the 18 months prior to the release of the survey. Exclusion criteria: Nonparticipating minors were not actively involved in the study.

#### Procedures

Ethical approval was granted by a British University Ethics Committee (ETHICS2020-61). The survey was available via a link to the website Online Surveys. Informed consent was required prior to proceeding with the questionnaires. The survey was open for 2 months and follow-up reminders were sent via e-mail.

#### Materials

The survey included sections on demographics (11 questions), previous concussion history (6 questions), policy awareness (2 questions) and football-related risk questions (3 questions).

In addition, the Rosenbaum Concussion Knowledge and Attitudes Survey (RoCKAS) was used which is considered a valid and reliable tool for assessing knowledge and attitude of concussion in collegiate athletes [13,14]. The tool consists of 53 items divided into 5 sections with two scores: Concussion

#### Table 1. Participant demographics.

Knowledge Index (CKI) and Concussion Attitude Index (CAI). The first subscale (CKI) comprises 14 true/false questions, three applied scenario true/false questions and nine symptom questions, with eight distractors (excluded from total score). Scoring was performed following the instructions from Rosenbaum and Arnett [14]. Possible scores for CKI ranged from 0 to 25 with a higher score representing greater concussion knowledge.

Responses to the CAI questions were measured on a 5-point Likert scale (1 = strongly disagree to 5 = strongly agree). Participants received a score between one and five points depending on the safety level of the response (1 = very unsafe response, 5 = very safe response). The range of the CAI scores ranged from 13 to 65, with 13 indicating very unsafe attitudes and 65 indicating very safe attitudes.

Perceived risk of football-related neurodegenerative disease was assessed using questions from a previously published study [15]. Using 7-point Likert scale participants indicated their perception and risk of developing dementia, Alzheimer's disease (AD), or chronic traumatic encephalopathy (CTE) later in life because of participating in football. Finally, participants were asked to indicate how strongly they agreed with three statements on the risks and benefits of playing football, in the short and long term as well as health benefits specifically. This was rated on a 7-point Likert scale ranging from 'Strongly disagree (1)' to 'Strongly agree (7).'

Participants had the opportunity to select if they received concussion education in the last 12 months and how this educational information was presented to them.

#### Data analysis

Descriptive statistics (means and standard deviations) were calculated using Excel (Version 1808) and SPSS Version 26 (IBM, Chicago, Illinois, USA) was used to calculate Pearson's correlation coefficient. An independent T-test was performed to calculate whether there was a significant difference between CAI scores in offensive and defensive players. The alpha level was set at p < 0.05 for all statistical tests.

				_	Gend Frequend	er cy (n)	
			Age				Playing years
Role		Frequency (n)	Mean $\pm$ SD	Male	Female	Prefer not to say	Mean $\pm$ SD
All participants		100.0 (236)	30.5 ± 11.2	75.4 (178)	23.7 (56)	0.8 (2)	5.2 ± 5.0
Player	Offence	40.7 (96)	29.4 ± 10.0	29.7 (70)	10.6 (25)	0.4 (1)	4.2 ± 3.8
	Defence	40.3 (95)	25.9 ± 6.3	29.2 (69)	10.6 (25)	0.4 (1)	5.5 ± 5.1
	Special Teams	0.8 (2)	24.0 ± 8.5	0.8 (2)	0.0 (0)	0.0 (0)	4.9 ± 4.8
	Total	64.0 (151)	25.6 ± 6.5	45.8 (108)	17.8 (42)	0.4 (1)	4.2 ± 3.8
Coach		13.6 (32)	43.1 ± 12.8	13.1 (31)	0.4 (1)	0.0 (0)	-
Support Personnel		2.5 (6)	45.7 ± 9.0	0.8 (2)	1.7 (4)	0.0 (0)	-
Combined role of player, coach or support personnel* Code		19.9 (47)	35.7 ± 11.6	15.7 (37)	3.8 (9)	0.4 (1)	-
	Tackle	81.8 (193)	29.3 ± 10.8	63.1 (149)	18.2 (43)	0.4(1)	-
	Flag	8.9 (21)	33.8 ± 10.2	5.1 (12)	3.8 (9)	0.0 (0)	-
	Tackle and Flag	9.3 (22)	37.6 ± 12	7.2 (17)	1.7 (4)	0.4 (1)	-

\* Support personnel was defined as a team manager, medical staff, conditioning staff or other (excluding coach or player).

Percentages based upon total participant number. Total participants n = 236

All playing positions apart from 'Punter' were covered.

#### **Results**

#### Football risks

Almost 79% (n = 186) of participants reported that developing dementia, Alzheimer's disease (AD) or chronic traumatic encephalopathy (CTE) would be very negative. Twenty nine percent of participants (n = 68) reported that they perceived it unlikely they would develop dementia, Alzheimer's disease (AD), or chronic traumatic encephalopathy (CTE) later in life as a result of playing football. Eighty-three percent agreed that the benefits of plaving football outweigh/ed the risks in the short term, with sixty two percent stating that the benefits of playing football outweigh/ed the risks in the long term. Fifty seven percent agreed that the benefits of playing football outweigh/ed the risks regarding health benefits specifically and sixteen percent reported they disagreed that the benefits of playing football outweigh/ed the risks regarding health benefits specifically. Twenty three percent (n = 62)neither agreed nor disagreed with this statement.

A mean score of 1.5 was reported for how positive or negative developing dementia, Alzheimer's disease (AD), or chronic traumatic encephalopathy (CTE) would be on a scale of 1 (very negative) to 7 (very positive). Of the typical neurodegenerative diseases, a mean score of 3.3 was reported for response to the question 'I will develop dementia, Alzheimer's disease (AD), or chronic traumatic encephalopathy (CTE) later in life because I played football' on a scale of 1 (definitely will not) to 7 (definitely will). On a scale of 1 (strongly disagree) to 7 (strongly agree) a mean score of 3.7 was noted for response to 'The benefits of playing football outweigh/ed the risks in the short term,' a mean score of 4.8 for response to 'The benefits of playing football outweigh/ed the risks in the long term' and finally a mean score of 4.7 for response to 'The benefits of playing football outweigh/ed the risks regarding health benefits specifically."

There was a negative correlation between football-related risks outweighing risks regarding health benefits and concussion reporting likelihood, r = -0.1, n = 236, p = 0.002 indicating that the more the athlete believes the benefits outweigh the risks the less likely they are to report a concussion, i.e. athletes are happier to continue to play football with injury.

#### **RoCKAS** survey

The total mean CKI score was  $21.0 \pm 2.1$  (range 12-25). Within the CKI, the most common knowledge question correctly identified was: Symptoms of a concussion can last for several weeks (98.3%, 232/236). The most common misconception identified was: An athlete who gets knocked out after a concussion is experiencing a coma (14.4%, 34/236). See Table 2.

The CAI component mean score was  $55.6 \pm 6.1$  (range 23–65). The most commonly accepted safe attitude was: coaches/managers need to be extremely cautious when determining whether an athlete should return to play (95.7%, 226/236). The most commonly accepted riskiest behavior was the athlete would continue playing sport whilst suffering with a headache as a result of concussion (45.3%, 107/236). See Table 3.

Figure 1 shows the mean CAI scores for positional groups. This includes players who have indicated other responsibilities in football.

The most common correct recognition of concussion symptoms were headache, difficulty concentrating and dizziness. The most concerning and least correctly identified symptoms was 'feeling slowed down.' See Table 4 for the percentage of respondents who correctly identified the concussion symptoms.

Correlations were performed to explore relationships between variables. See Table 5 for results.

Table 2. Percentage of correct responses to knowledge RoCKAS questions.

Question	%	n
Symptoms of a concussion can last for several weeks. (True)	98.3	232
Concussions can sometimes lead to emotional disruptions. (True)	97.5	230
There is a possible risk of death if a second concussion occurs before the first one has healed. (True)	90.7	214
People who have one concussion are more likely to have another concussion. (True)	69.5	164
In order to be diagnosed with a concussion, you have to be knocked out. (False)	99.6	235
A concussion can only occur if there is a direct hit to the head. (False)	91.1	215
Being knocked unconscious always causes permanent damage to the brain. (False)	84.3	199
Sometimes a second concussion can help a person remember things that were forgotten after the first concussion. (False)	95.3	225
After a concussion occurs, brain imaging (e.g. CAT scan, MRI, X-Ray, etc) typically shows visible physical damage (e.g. bruise, blood clot) to the brain. (False)	38.6	91
If you receive one concussion and you have never had a concussion before, you will become less intelligent. (False)	98.3	232
After 10 days, symptoms of a concussion are usually completely gone. (True)	48.3	114
After a concussion, people can forget who they are and not recognize others but be perfect in every other way. (False)	26.3	62
An athlete who gets knocked out after getting a concussion is experiencing a coma. (True)	14.4	34
There is rarely a risk to long-term health and wellbeing from multiple concussions. (False)	89.8	212

Table 3. Percentage of safer responses to risk attitude RoCKAS questions.

Question	%	n
I feel that coaches/managers need to be extremely cautious when determining whether an athlete should return to play.	95.76	226
I would continue playing a sport while also having a headache that resulted from a minor concussion.	45.34	107
I feel concussions are less important than other injuries.	91.10	215
I feel that an athlete has a responsibility to return to a game even if it means playing while experiencing symptoms of a concussion.	90.25	213
I feel that an athlete who is knocked unconscious should be taken to the emergency room.	87.29	206





Figure 1. Mean CAI scores according to position.

\*No significant difference between CAI score and offensive and defensive playing position only p < 0.097 level

Table 4	Percentage of	correct responses to	<b>RoCKAS 16-item</b>	symptom	recognition of	checklist.
				<i>, ,</i>		

	Player	Coach	Support Personnel	Player, Coach, Support Personnel	Player, Coach	Coach, Support Personnel	Player, Support Personnel	Total
Headache	99.3	100	100	100	100	100	83.3	99.1
Difficulty speaking	100	100	100	100	100	100	100	76.1
Sensitivity to light	96.0	96.9	83.3	75	100	40	100	94.9
Difficulty remembering	91.3	96.9	83.3	100	93.8	80	83.3	92.0
Panic attacks	100	100	100	100	100	100	100	26.7
Drowsiness	84.1	96.9	83.3	100	84.3	60	83.3	85.6
Feeling like in a 'fog'	85.4	100	83.3	100	100	80	100	89.8
Feeling slowed down	77.5	71.9	66.7	75	71.9	60	83.3	75.4
Reduced breathing rate	100	100	100	100	100	100	100	30.1
Difficulty concentrating	95.4	100	100	100	100	100	83.3	96.6
Dizziness	96.7	100	100	100	100	80	100	97.5
Hives	100	100	100	100	100	100	100	0.4
Arthritis	100	100	100	100	100	100	100	0.4
Weight Gain	98.0	90.6	50	25	90.6	40	50	1.3
Excessive Studying	98.0	90.6	50	25	90.6	40	50	1.7
Hair loss	100	100	100	100	100	100	100	0

\*The percentage (%) of respondents who correctly included each symptom and correct concussion symptoms are in bold.

There was significant positive correlation between CAI and the likelihood to report concussion and age. Negative correlations were noted between CAI and risk attitude to heath specifically and prior history of concussion. In addition, there were significant positive correlations between CKI and likelihood to report concussion. See Table 5 for the relationship of CAI and CKI with variables. Negative correlations were found between CAI and perceived benefits of playing AF versus the long- and short-term risk to health. No clear correlation was noted between CKI and perceived benefits of playing AF versus the long- and short-term risk to health.

#### Concussion, concussion education, and policy

The mean number of diagnosed concussions by a medical practitioner of all participants was 0.9 ( $\pm$  1.4). The mean number of suspected concussions of all participants was 2.3 ( $\pm$  3.7).

Table 5. Relationships of participants' response to concussion attitude index and concussion knowledge index to age, prior concussion diagnosis, reporting likelihood and attitude to risk.

		Age	Prior Concussion Diagnosis	Likelihood to report concussion to coach or medic	Attitude to health risks specifically vs the benefits of plaving AF	Attitudes toward the benefits of playing AF vs risks in the long term	Attitudes toward the benefits of playing AF vs risks in the short term
CAL	r	228	- 231	440	– 163		- 218
CAI	Sig. (2-tailed)	.000**	.000**	.000**	.012*	.003**	.001**
CKI	r	.124	059	.172	.002	.047	.038
	Sig. (2-tailed)	.058	.365	.008**	.978	.468	.563

\*Correlation is significant at 0.05 level \*\*Correlation is significant at 0.01 level

Total participants n = 236

Table 6. Participant's recognition of methods of receiving concussion education.

Q. If you received education about concussion, how was this information presented?	
Consent form required for participation	28 (24.8)
Education from my coach	34 (30.1)
Education from my parents	5 (4.4)
Education from a medical provider (i.e. sports therapist, physiotherapist, doctor etc)	39 (34.5)
Web based resource (e.g. educational blog)	66 (58.4)
Social Media	20 (17.7)
Magazines or other print material	8 (7.1)
Concussion fact sheet	35 (31.0)
Other	26 (23.0)

\*Brackets denote percentage of responses in each category.

Almost 47.9% (n = 113) of participants reported receiving education about sport-related concussion in the past 12 months. However, 48.3% (n = 114) reported that they had received no education in the past 12 months. A small percentage (3.8%, n = 9) reported that they did not know or were unsure if they had received education in the last 12 months. See Table 6 for methods of receiving concussion education information. Of those who selected the 'other' option, participants noted academic papers, coaching qualifications, formal training as a medical practitioner and online concussion awareness courses as forms of education.

When participants were asked about their awareness of the BAFA concussion policy, 63.1% (n = 149) reported they were aware. However, when asked whether they had previously accessed and read the BAFA Concussion policy, 63.6% (n = 150) reported that they had not accessed or read the policy.

#### Discussion

This is the first study to provide an insight into the current state of concussion knowledge and attitudes within BAF. The results suggest that concussion knowledge among BAF participants was generally good. Worryingly, regarding attitudes, the more an individual believes the benefits of playing outweigh the risks, the more likely they are to under-report symptoms and play through injury.

Over half of all participants had previously sustained a selfdiagnosed concussion, however less than half had been medically diagnosed. This supports previous findings in BAF [4] and could explain the increased confidence in knowledge of concussion. However, prior findings suggest knowledge may not be sufficient for injury self-diagnosis and give athletes a 'false sense of security' [16]. It is positive that participants felt able to self-diagnose concussion, but this does not explain why there was no formal diagnosis by a medical practitioner.

The primary finding of this study was that attitudes toward and knowledge of concussion may influence player behavior. Athletes with a safer attitude to concussion are overall more cautious in their approach to concussion and thus are more likely to report the injury to a medical practitioner or coach. However, it is concerning that almost half of participants stated that they would continue to play with a concussion. In addition, increased knowledge of concussion results in players being more likely to report concussive symptoms. However, intention to report is not always reflective of true reporting behavior [17] but this may improve with the introduction of concussion education. Furthermore, future research could explore the impact of educational delivery methods in BAF.

Many BAF players in the UK follow the NFL and, despite a current lack of research in BAF concussion, players might be aware of discussions from the US. In recent years, the NFL has addressed the 'concussion crisis' by actioning a response strategy of media and marketing [18], encouraging player educational programmes across many sports, under the Lystedt Law. Through this increased media coverage, we might expect that UK players might be more aware of the implications of playing with head injury and associated long-term risks. Moreover, as BAF is currently an amateur sport, players might be more cautious of the long-term effects of injury on their wider life (i.e. careers, family or studies), meaning they are safer in their approach to injury and reporting behavior. Indeed, prior research suggests athletes' risk-taking attitude is linked to career decisions [19].

Our results confirm that an increased knowledge of concussion does not affect the number of concussions a player has, however this prior knowledge might help players to selfdiagnose a concussion. With an increased ability to selfdiagnose, players may be able to prevent secondary or tertiary harm [20]. Furthermore, knowledge of concussion was not related to the number of concussions, suggesting that athletes with no prior history of concussion have a similar understanding as those who have had one, agreeing with research conducted with Japanese collegiate athletes [21]. Our findings are positive, suggesting that concussion education is reaching the whole community. However, there is a paradox given that 37% of participants reported to have not accessed and read the BAFA concussion policy. At present there is no mandate in BAF for those involved in the game to be educated on concussion and some participants have had no concussion training at all [4]. It therefore cannot be presumed that all participants are accessing this information from the NGB and are getting information elsewhere. Concussion educational material was reported to come from a range of sources (e.g. web-based), however the specific details about educational platforms warrants further investigation. Table 7 outlines a series of educational recommendations for the NGB.

In general, improved knowledge positively influenced likelihood to report concussion to a coach or medic. Greater knowledge of concussion was demonstrated through increased recognition of signs and symptoms. In line with other studies [21], most participants correctly identified the highly recognizable symptoms such as headache, difficultly

#### Table 7. Recommendations.

#### Recommendations for future directions

- 1. The development of BAF-specific concussion educational programmes to include the dangers of under-reporting injury and risk taking, This should be delivered to coaches, athletes, parents/guardians, game-day staff, referees and medical staff.
- 2. Regular concussion educational training to be mandated within the concussion policy. This educational training should be provided by a healthcare practitioner or academic at minimum of once per year/season.
- 3. The development of BAF-specific concussion posters and leaflets which can be shared with all NGB associated teams upon review of the concussion policy each year.
- 4. The NGB should regularly share concussion education on social media platforms.
- 5. Future studies should focus on one pool of participants e.g. players only, to allow for a targeted approach to informed education.

concentrating and dizziness while fewer identified symptoms such as feeling like in a 'fog.' However, just as there are common symptoms of concussion (e.g. headache), there are broader acute signs of concussion which might require medical examination. It is of concern that a number of symptom distractors were incorrectly selected such as 'feeling slowed down'. This is a concern because this is a common symptom of fatigue and so a player could be misdiagnosed.

Furthermore, our findings suggest that as age increases, risk attitudes become more cautious, agreeing with Kerr [22] and Hutchinson et al. [23]. This may not be unique to athletes as humans naturally become more cautious about health with age in general life [24]. We found no significant relationship between age and concussion knowledge however, despite previous findings which suggest that concussion knowledge increases with age [22]. No matter the age of an individual, knowledge may be perceived in different ways despite the amount or type of knowledge, and this requires further investigation.

It is concerning that the more athletes believe the benefits of playing outweigh the risk to health, the less likely they are to report injury. Although this study did not explore what the individuals perceived benefits might be, prior research suggests that individuals report the benefits of sport participation to be: improvements in daily life, improved physical and mental wellbeing and social outlets [25]. Thus, if BAF players perceive these benefits of being part of team as more important that personal wellbeing, they may choose to play through injury.

Amateur BAF participants may believe the long-term health implications seen at the professional level will not impact them. Indeed, amateur rugby players have been noted to downplay and ignore concussion suggesting they do not fully understand concussion but have trivialized it as part of the sport [26]. In addition, athletes may only remove themselves from play if they perceived the injury posed a long-term threat to their own health [27]. Despite these findings it should be noted that Liston et al. [27] reported data from a small pool of university rugby athletes and so may not be representative of all athletes. Comparable findings have also been seen at 'sub-elite' level with clinicians reporting that athletes do not fully understand their bodies and think they are invincible [28]. As such, the education of amateur athletes may be insufficient, and they do not believe that serious health implications can occur at that level of the game. Indeed, this study found that over half of participants perceived it unlikely they will develop a neurodegenerative disease as a result of playing football. At present, there is limited

research to confirm the likelihood of developing neurodegenerative disease in amateur contact athletes, however links have been noted in the professional game [29].

It has been suggested that athletes reporting decisions are dependent on the athlete's level of competitiveness and game circumstances, for example the availability of substitutions or the importance of the game [30]. To overcome the underreporting of injury in sports such as soccer, governing bodies such as the English Premier League have brought in the concussion substitute law which allows each team to have two permanent concussion substitutes whilst players are assessed by medics. However, unlike other sports with smaller squad sizes, substitutions are not an issue in BAF where teams have no maximum roster size nor are there restrictions on substitutions. Yet, at an amateur level, not all teams will have roster depth and so some players may be required to play in multiple positions, thus placing the athletes under pressure to stay on the field. Further investigation might be taken to investigate BAF players reasons for concussion reporting barriers.

Upon suspecting symptoms of concussion, athletes should be removed or remove themselves from physical activity. However, despite good knowledge or prior concussion history, underreporting is still present. Indeed, Liston et al. [26] found contact athletes preferred receiving a concussion over musculoskeletal injury which in their eyes would mean less time out of the game. Secondly, the participants somewhat rationalized the benefits of concussion, allowing them to play in a 'primal state' and so not capable of conscious cognition thus not needing to adhere to the social norms of reporting behavior [31]. However, studies have reported that contact athletes with prior injury history demonstrate greatest worry and concern of re-injury [32]. There is currently a lack of research looking at the role fear of re-injury has on symptom reporting and future research could seek to investigate this. This might be further explained by the manifestation of anxiety in the injured athlete. Fear of losing their position on the team, letting the team, family, friends or community down, believing they could manage the injury alone or losing their athletic ability which has been previously reported to pressure the athlete to continue to play [29]. Athletes will continue to play under the pressure of coaches and parents, changing reporting behavior which might intensify the athlete's reason to underreport [33].

Underreporting is present in BAF. Players may hold misconceptions about concussions which lead to underreporting of injury exposing the athlete to considerable health implications. As a developing sport, BAF has an opportunity to challenge these misconceptions through player education.

#### Limitations

This is the first known study of concussion knowledge and attitudes in BAF however there are some limitations. The authors recognize the limitation in analyzing the mix of groups however this is an exploratory study looking to capture current concussion knowledge. Therefore, future research may look to investigate individual groups or differences between these groups. This study was cross-sectional in design; therefore, the data should be regarded as a snapshot of behavior in a single moment of time and thus could be unrepresentative of group behavior as a whole. Participants in this study were contacted and invited to participate through varied avenues resulting in participants choosing to participate which might have resulted in selection bias. It was difficult to contact all registered BAFA members due to opt-out marketing communication options. Additionally, some areas of the survey required athletes to rely upon their ability to remember past events which may have led to recall bias and potential inaccuracies. Furthermore, it was assumed that respondents were honest in their responses and did not answer based on what they thought to be socially acceptable. However, these findings should not be discredited and provide primary data on knowledge and attitudes of concussion in BAF.

#### **Future directions**

With the support of key stakeholders, future research should aim to further investigate perceived risk and reporting behavior in BAF to inform future educational strategies and reduce risky concussion-related behavior. Our recommendations for the NGB are outlined in Table 7. We recommend that key stakeholders in the game emphasize the importance of a safe approach to concussion reporting and encourage regular concussion education. By creating a culture of reporting concussive symptoms, we may see safer reporting trends [16]. Furthermore, research might seek to understand the true sources of education in BAF participants to inform future educational strategies of the NGB and examine the effectiveness (both in cost and player knowledge) when introducing educational programmes in BAF.

#### Conclusion

This is the first study to examine attitudes and knowledge of concussion in BAF and could be used to inform BAF concussion education, implemented within BAF coaching workshops. The findings are generally positive, with knowledge of concussion and symptom recognition generally good. However, these findings show there is evidence of risky behavior seen in BAF through unsafe unlikeliness to report and poor attitudes to long-term health. Many respondents had not received concussion education in the last 12 months nor accessed the NGB concussion policy. Future interventions should look to challenge these misconceptions, improving attitude toward concussion, reporting behavior and educational strategy. Future research is needed to examine the relationship between concussion-related knowledge and attitudes and reporting behaviors in BAF athletes.

#### Acknowledgments

The authors would like to express thanks to the British American Football community for participating in this study and to the British American Football Association for their support in sharing the survey to its members.

#### **Disclosure statement**

No potential conflict of interest was reported by the author(s).

#### Funding

The author(s) reported there is no funding associated with the work featured in this article.

#### Notes on contributors

**Eleanor Travis** is a lecturer and Graduate Sports Therapist with an interest in American football and works with teams across the United Kingdom including the Great Britain tackle and flag teams. Additionally, Eleanor has been part of two BUCS National Championship winning teams whilst working in a coaching and sports therapy practitioner capacity.

Andrea Scott-Bell is a senior lecturer and programme leader at Northumbria University. Her PhD from Loughborough University examined the professionalization, formalization and bureaucratization of sport and exercise medicine in British Olympic sports. Andreas current research interests are broadly focused on athlete healthcare in sport.

**Claire Thornton** is a senior lecturer at Northumbria University and an experienced sports coach. Claire's teaching interests center on sport and exercise psychology and research methods. Claire's PhD explored the mechanisms for differences in pain tolerance, perception and coping between contact and non-contact athlete groups.

#### **Declaration of interest**

ET and ASB are members of the Sport Science and Medicine Committee for the British American Football Association. The views expressed in this study are not influenced by the committee.

This research was conducted in the absence of financial interest.

#### **Author's contributions**

ET and CT designed the study. ET collected and analysed the data with contributions from CT. ET, CT and ASB drafted the manuscript. All authors have read and approved the final version of the manuscript and agree with the order of presentation of the authors.

#### ORCID

Dr Andrea Scott-Bell D http://orcid.org/0000-0002-1022-9023 Dr Claire Thornton D http://orcid.org/0000-0002-5032-1836

#### References

- 1. BAFA. (2021). British *American* Football Annual Report 2021. Available: https://www.britishamericanfootball.org/wp-content /uploads/2021/11/BAFA-Annual-Report-2021-1.pdf. Last accessed 2022 Apr 19th.
- Willigenburg NW, Borchers JR, Quincy R, et al. Comparison of injuries in American collegiate football and club rugby: a prospective cohort study. Am J Sports Med. 2016;44(3):753–760.

- 3. Bayram JM, Hamilton DF, Saunders DH. Epidemiology of American football injuries at universities in the **United Kingdom**. Orthop J Sports Med. 2020;8(10):2325967120960206.
- Travis E, Thornton C, Scott-Bell A. Concussion reporting and safeguarding policy development in British *American* Football: an essential agenda. Front Sport Active Liv. 2021;3:118.
- Tator C, Starkes J, Dolansky G, et al. Fatal second impact syndrome in rowan stringer, a 17-year-old rugby player. Can J Neurol Sci / J Can Des Sci Neurol. 2019;46(3):351–354.
- Omalu B, Hammers J. Recommendation to create new neuropathologic guidelines for the post-mortem diagnosis of chronic traumatic encephalopathy. Neurosurgery. 2021;89(1):E97–E98.
- Harris SA, Chivers PT, McIntyre FL, et al. Exploring the association between recent concussion, subconcussive impacts and depressive symptoms in male Australian Football players. BMJ Open Sport Exerc Med. 2020;6(1):e000655.
- Collins MW, Lovell MR, Iverson GL, et al. Cumulative effects of concussion in high school athletes. Neurosurgery. 2002;51 (5):1175–1181.
- 9. Zemper ED. Two-year prospective study of relative risk of a second cerebral concussion. Am J Phys Med Rehabil. 2003;82 (9):653–659.
- Guskiewicz KM, McCrea M, Marshall SW, et al. Cumulative effects associated with recurrent concussion in collegiate football players: the NCAA concussion study. Jama. 2003;290(19):2549–2555.
- van Ierssel J, Osmond M, Hamid J, et al. What is the risk of recurrent concussion in children and adolescents aged 5–18 years? A systematic review and meta-analysis. Br J Sports Med. 2021;55 (12):663–669.
- 12. White PE, Newton JD, Makdissi M, et al. Knowledge about sports-related concussion: is the message getting through to coaches and trainers? Br J Sports Med. 2014;48(2):119–124.
- 13. Chapman EB, Nasypany A, May J, et al. Investigation of the rosenbaum concussion knowledge and attitudes survey in collegiate athletes. Clin J Sport Med. 2018;28(2):117–124.
- Rosenbaum AM, Arnett PA. The development of a survey to examine knowledge about and attitudes toward concussion in high-school students. J Clin Exp Neuropsychol. 2010;32(1):44–55.
- Baugh CM, Gedlaman MA, Daneshvar DH, et al. Factors influencing college football players' beliefs about incurring football-related dementia. Orthop J Sports Med. 2021;9(4):23259671211001129.
- Weber M, Edwards MG. Sport concussion knowledge in the UK general public. Arch Clin Neuropsychol. 2012;27(3):355–361.
- Register-Mihalik JK, Linnan LA, Marshall SW, et al. Using theory to understand high school aged athletes' intentions to report sportrelated concussion: implications for concussion education initiatives. Brain Inj. 2013;27(7–8):878–886.
- 18. LaGree D, Wilbur D, Cameron GT. A strategic approach to sports crisis management: assessing the NFL concussion crisis from

marketing and public relations perspectives. Int J Sports Mark Spons. 2019;20(3):407-429

- 19. Gakhar DV. Risk attitude and financial awareness of inter-collegiate athletes. 2020.
- 20. Register-Mihalik J, Baugh C, Kroshus EY, et al. A multifactorial approach to sport-related concussion prevention and education: application of the socioecological framework. J Athl Train. 2017;52 (3):195–205.
- 21. Suzuki K, Imamoto T, Nagai S, et al. Knowledge of, and attitudes toward, concussion in Japanese male collegiate athletes. Front Sport Active Liv. 2022;4:42.
- 22. Kerr ZY, Nedimyer AK, Kay MC, et al. Factors associated with concussion-symptom knowledge and attitudes toward concussion care seeking in a national survey of parents of middle-school children in the US. J Sport Health Sci. 2021;10(2):113–121.
- Hutchinson S, Ellison P, Levy A, et al. Knowledge and attitudes towards concussion in UK-based male ice hockey players: a need for attitude change? Int J Sports Sci Coaching. 2019;14(2):153–161.
- Grøtvedt L, Stavem K. Association between age, gender and reasons for smoking cessation. Scand J Public Health. 2005;33 (1):72–76.
- Lovell GP, El Ansari W, Parker JK. Perceived exercise benefits and barriers of non-exercising female university students in the United Kingdom. Int J Environ Res Public Health. 2010;7(3):784–798.
- Liston K, McDowell M, Malcolm D, et al. On being 'head strong': the pain zone and concussion in non-elite rugby union. Int Rev Sociol Sport. 2018;53(6):668–684.
- 27. Liston K, Reacher D, Smith A, et al. Managing pain and injury in non-elite rugby union and rugby league: a case study of players at a British university. Sport in Society. 2006;9(3):388–402
- Barrette A, Harman K. Athletes play through pain—what does that mean for rehabilitation specialists? J Sport Rehabil. 2019;29 (5):640–649.
- Roberts AL, Pascual-Leone A, Speizer FE, et al. Exposure to American football and neuropsychiatric health in former national football league players: findings from the football players health study. Am J Sports Med. 2019;47(12):2871–2880.
- Williams JM, Langdon JL, McMillan JL, et al. English professional football players concussion knowledge and attitude. J Sport Health Sci. 2016;5(2):197–204.
- Liston K, Malcolm D. Sports-related brain injury: concussion and chronic traumatic encephalopathy. In: Young, K the suffering body in sport. Bingley: Emerald Publishing Limited; 2019. p. 98.
- 32. Short SE, Reuter J, Brandt J, et al. The relationships among three components of perceived risk of injury, previous injuries and gender in contact sport athletes. Athletic Insight. 2004;6(3):78–85.
- Bloom GA, Trbovich AM, Caron JG, et al. Psychological aspects of sport-related concussion: an evidence-based position paper. J Appl Sport Psychol. 2020;34:1–23.

## Knowledge and attitudes of concussion of players in British American Football

\*Required response

Note: Below survey excludes logic flow of questions.

- (1) Please state your age e.g. '21'\*
- (2) Please state your gender\*
  - a. Male
  - b. Female
  - c. Prefer not to say
  - d. Other
    - (i) If you selected Other, please specify:
- (3) What form of British American Football are you involved in either as a coach, player or support personnel?\*
  - a. Tackle
  - b. Flag
  - c. Both
- (4) Please state how you are involved in American Football. If you have several roles, please select all that apply.\* (Select all that apply)
  - a. Player
  - b. Coach
  - c. Support Personnel
- (5) What is the highest level of the sport you coach?
  - a. Premiership
  - b. Division 1
  - c. Division 2
  - d. Division 3
  - e. Associate
  - f. Recreational
  - g. National
- (6) How many years' experience do you have coaching American football (tackle, flag football or combined flag/tackle)?
- (7) Is American Football the primary sport you coach?
  - a. Yes
  - b. No
    - (i) As you answered 'no' to the previous question, what is the primary sport you coach?
- (8) What is the highest level of the sport you have played for? Please select one option for tackle and one for flag football if this applies.
  - a. Premiership Tackle
  - b. Division 1 Tackle
  - c. Division 2 Tackle
  - d. Associate Tackle
  - e. National Tackle
  - f. Premiership Flag Football
  - g. Division 1 Flag Football
  - h. Associate Flag Football
  - i. Recreational Flag Football
  - j. National Flag Football
- (9) Primary playing Position:
  - a. Quarter Back
  - b. Tight End
  - c. Offensive tackle
  - d. Offensive guard
  - e. Center
  - f. Running back
  - g. Full back
  - h. Wide receiver
  - i. Defensive end
  - j. Defensive tackle
  - k. Outside line-backer
  - I. Insider line-backer
  - m. Cornerback
  - n. Safety
  - o. Kicker
  - p. Punter
  - q. Blitzer
- (10) Please state your total number of years playing tackle, flag or combined tackle and flag football

- (11) How many diagnosed concussions have you sustained during your American Football playing career? (Concussion is defined as an injury resulting from a blow to the head or body that caused an alteration in mental status and one or more of the following symptoms: headache, nausea, vomiting, dizziness/balance problems, fatigue, trouble sleeping, drowsiness, sensitivity to light or noise, blurred vision, difficulty remembering and difficulty concentrating.) \*
- (12) How many suspected concussions have you sustained during your American Football playing career? (Concussion defined as an injury resulting from a blow to the head or body that caused an alteration in mental status and one or more of the following symptoms: headache, nausea, vomiting, dizziness/balance problems, fatigue, trouble sleeping, drowsiness, sensitivity to light or noise, blurred vision, difficulty remembering and difficulty concentrating.)\*
- (13) Please state the date of most recent diagnosed or suspected concussion. If you have not had a concussion please put 'N/A'. \*
- (14) On a scale of 1–10, how likely are you to report to your medic or coach that you think you might have a concussion? (1 – not likely, 10 – very likely) \*
- (15) Have you received education about sport-related concussion in the past 12 months?
  - a. Yes
  - b. No
  - c. Do not know/unsure
- (16) If you have received education about concussion, how was the information presented?
  - a. Consent form required for participation
  - b. Education from my coach
  - c. Education from my parents
  - d. Education from a medical provider (i.e. sports therapist, physiotherapist, doctor etc.)
  - e. Web-based resource (e.g. educational blog)
  - f. Social Media
  - g. Magazines or other print material
  - h. Concussion fact sheet

pletely gone.

- i. Other
  - (i) If you selected Other, please specify:
- (17) Did you know that there is a BAFA Concussion Policy? \*
  - a. Yes
  - b. No
- (18) Have you previously accessed and read the BAFA Concussion Policy?
  - a. Yes
  - b. No
- (19) Part 1 DIRECTIONS: Please read each of the following scenarios and answer TRUE or FALSE for each question that follows the following scenarios. True/False response required \*
  - a. 1. There is a possible risk of death if a second concussion occurs before the first one has healed.
  - b. 2. Running everyday does little to improve cardiovascular health.
  - c. 3. People who have one concussion are more likely to have another concussion.
  - d. 4. Cleats help athletes feet grip the playing surface.
  - e. 5. In order to be diagnosed with a concussion, you have to be knocked out.
  - f. 6. A concussion can only occur if there is a direct hit to the head.
  - g. 7. Being knocked unconscious always causes permanent damage to the brain.
  - h. 8. Symptoms of a concussion can last for several weeks.
  - i. 9. Sometimes a second concussion can help a person remember things that were forgotten after the first concussion.
  - j. 10. Weightlifting helps to tone and/or build muscle.
  - After a concussion occurs, brain imaging (e.g. CAT scan, MRI, X-Ray, etc) typically shows visible physical damage (e.g. bruise, blood clot) to the brain.
  - I. 12. If you receive one concussion and you have never had a concussion before, you will become less intelligent.
    m. 13. After 10 days, symptoms of a concussion are usually com-

- n. 14. After a concussion, people can forget who they are and not recognize others but be perfect in every other way.
- o. 15. High-school athletes and college athletes tend to be the same age.
- p. 16. Concussions can sometimes lead to emotional disruptions.
- q. 17. An athlete who gets knocked out after getting a concussion is experiencing a coma.
- r. 18. There is rarely a risk to long-term health and wellbeing from multiple concussions.
- (20) Part 2a DIRECTIONS: Please read each of the following scenarios and circle TRUE or FALSE for each question that follows the scenario. Scenario 1: While playing in a game, Player Q and Player X collide with each other and each suffers a concussion. Player Q has never had a concussion in the past. Player X has had 4 concussions in the past. True/False response required \*
  - a. 1. It is likely that Player Q's concussion will affect his long-term health and wellbeing.
  - b. 2. It is likely that Player X's concussion will affect his long-term health and wellbeing.
- (21) Part 2b DIRECTIONS: Please read each of the following scenarios and circle TRUE or FALSE for each question that follows the scenario. Scenario 2: Player F suffered a concussion in a game. She continued to play in the same game despite the fact that she continued to feel the effects of the concussion. True/False response required \*
  - a. 3. Even though Player F is still experiencing the effects of the concussion, her performance will be the same as it would be had she not suffered a concussion.
- (22) Part 3 \* Please do not select more than 1 answer(s) per row. Please select exactly 8 answer(s). Please do not select more than 5 answer(s) in any single column. (1-strongly disagree, 2 – disagree, 3 – neutral, 4 – agree, 5 strongly agree)
  - a. 1. I would continue playing a sport while also having a headache that resulted from a minor concussion.
  - b. 2. I feel that coaches/managers need to be extremely cautious when determining whether an athlete should return to play.
  - c. 3. I feel that mouth guards protect teeth from being damaged or knocked out.
  - d. 4. I feel that professional athletes are more skilled at their sport than high-school athletes.
  - e. 5. I feel concussions are less important than other injuries.
  - f. 6. I feel that an athlete has a responsibility to return to a game even if it means playing while experiencing symptoms of a concussion.
  - g. 7. I feel that an athlete who is knocked unconscious should be taken to the emergency room.
  - h. 8. I feel that most highschool athletes will play professional sports in the future.
- (23) Part 4 DIRECTIONS: For each question read the scenarios and select the number that best describes your view. (For the questions that ask you what most athletes feel, base your answers on how you think MOST athletes would feel).

Scenario 1: Player R suffers a concussion during a game. Coach A decides to keep Player R out of the game. Player R's team loses the game. \* Please do not select more than 1 answer(s) per row. Please select exactly 2 answer(s). Please do not select more than 2 answer(s) in any single column. (1-strongly disagree, 2 – disagree, 3 -neutral, 4 – agree, 5 strongly agree)

- a. 1. I feel that Coach A made the right decision to keep Player R out of the game.
- b. 2. Most athletes would feel that Coach A made the right decision to keep Player R out of the game.

Scenario 2: Athlete M suffered a concussion during the first game of the season. Athlete O suffered a concussion of the same severity during the semifinal playoff game. Both athletes had persisting symptoms. \* Please do not select more than 1 answer(s) per row. Please select exactly 4 answer(s). Please do not select more than 4 answer(s) in any single

column. (1-strongly disagree, 2 – disagree, 3 – neutral, 4 – agree, 5 strongly agree)

- a. 3. I feel that Athlete M should have returned to play during the first game of the season.
- b. 4. Most athletes would feel that Athlete M should have returned to play during the first game of the season.
- c. 5. I feel that Athlete O should have returned to play during the semifinal playoff game.
- d. 6. Most athletes feel that Athlete O should have returned to play during the semifinal playoff game.

Scenario 3: Athlete R suffered a concussion. \*Athlete R's team has an sports therapist/physiotherapist on the staff. Please do not select more than 1 answer(s) per row. Please select exactly 2 answer(s). Please do not select more than 2 answer(s) in any single column. (1-strongly disagree, 2 – disagree, 3 -neutral, 4 – agree, 5 strongly agree)

- (1) 7. I feel that the sports therapist/physiotherapist, rather than Athlete R, should make the decision about returning Athlete R to play.
- (2) 8. Most athletes would feel that the sports therapist/physiotherapist, rather than Athlete R, should make the decision about returning Athlete R to play.
- (3) Part 5 DIRECTIONS: Think about someone who has had a concussion. From the following signs and symptoms select those that you believe someone may be likely to experience AFTER a concussion. \*
  - a. Hives
  - b. Headache
  - c. Difficulty speaking
  - d. Arthritis
  - e. Sensitivity to light
  - f. Difficulty remembering
  - g. Panic attacks
  - h. Drowsiness
  - i. Feeling like in a 'fog'
  - j. Weight gain
  - k. Feeling slowed down I. Reduced breathing rate
  - I. Reduced breathing rate m. Excessive studying
  - n. Difficulty concentrating
  - o. Dizziness
  - p. Hair loss
- (4) To what extent do you agree with the following statement? (1 very negative to 7 definitely will) \*
  - a. 'I will develop dementia, Alzheimer's disease (AD), or chronic traumatic encephalopathy (CTE) later in life because I played football'.

(5) Please rate the following \* (1 – very negative to 7 – very positive)

- Please rate how positive or negative (in your opinion) developing dementia, Alzheimer's disease (AD), or chronic traumatic encephalopathy (CTE) would be.
- (6) Football-related risk-benefit Please indicate how strongly you agree with the following statements \* Please do not select more than 1 answer(s) per row. Please select exactly 3 answer(s). Please do not select more than 3 answer(s) in any single column. (1 – strongly agree, 2 – disagree, 3 – somewhat disagree, 4 – neither agree nor disagree, 5 – somewhat agree, 6 – agree, 7 – strongly agree)
  - The benefits of playing football outweigh/ed the risks in the short term'.
  - b. The benefits of playing football outweigh/ed the risks in the long term'.
  - c. The benefits of playing football outweigh/ed the risks regarding health benefits specifically'.