

Northumbria Research Link

Citation: Avery, Leah, Exley, Catherine, McPherson, Stuart, Trenell, Michael, Anstee, Quentin M. and Hallsworth, Kate (2017) Lifestyle Behavior Change in Patients With Nonalcoholic Fatty Liver Disease: A Qualitative Study of Clinical Practice. *Clinical Gastroenterology and Hepatology*, 15 (12). pp. 1968-1971. ISSN 1542-3565

Published by: Elsevier

URL: <https://doi.org/10.1016/j.cgh.2017.06.011>
<<https://doi.org/10.1016/j.cgh.2017.06.011>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/31486/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

Lifestyle Behavior Change in Patients With Nonalcoholic Fatty Liver Disease: A Qualitative Study of Clinical Practice

Leah Avery,^{*} Catherine Exley,[‡] Stuart McPherson,[§] Michael I. Trenell,^{*} Quentin M. Anstee,^{*,§} and Kate Hallsworth^{*,§}

^{*}Institute of Cellular Medicine, Newcastle University, Newcastle upon Tyne, United Kingdom; [‡]Faculty of Health and Life Sciences, Northumbria University, Newcastle upon Tyne, United Kingdom; [§]Liver Unit, Newcastle Upon Tyne Hospitals National Health Service Foundation Trust, Newcastle upon Tyne, United Kingdom

Nonalcoholic fatty liver disease (NAFLD) is the most common liver condition worldwide and is linked largely to obesity and inactivity. Lifestyle modification is the primary treatment for NAFLD targeting dietary change, physical activity, and exercise to facilitate weight loss and weight loss maintenance.¹⁻³ This has been shown to reduce steatosis and ameliorate steatohepatitis. European Clinical Practice Guidelines for the management of NAFLD³ highlight the importance of targeting lifestyle behavior change in all patients with NAFLD regardless of disease severity. These guidelines recommend combining dietary restriction and a progressive increase in aerobic exercise and resistance training with a focus on tailoring interventions to the individual patient. Practice guidelines published by the American Association for the Study of Liver Diseases⁴ recommend weight loss of at least 3% to 5% of body weight via hypocaloric diet or diet combined with increased physical activity but state that these lifestyle interventions should target patients with nonalcoholic steatohepatitis. Given the benefits of lifestyle behavior change, this study explored the perceptions surrounding clinical care as currently offered to patients with NAFLD. The aim of this study was to establish whether current provision of lifestyle behavior change support is sufficient, whether health care professionals believe they have the tools to target lifestyle behavior changes effectively, and how targeting diet and physical activity/exercise to facilitate weight loss and weight loss maintenance in practice can be improved from the perspective of health care professionals and patients.

Methods

Semistructured qualitative interviews were conducted with 21 health care professionals from 2 UK National Health Service Hospital Trusts and 11 UK National Health Service Clinical Commissioning Groups across a range of specialties (hepatology, gastroenterology diabetology, and primary care) and 12 patients diagnosed with NAFLD. Interviews were conducted using 2 interview topic guides developed with reference to the American Association for

the Study of Liver Diseases and the European Association for the Study of the Liver, European Association for the Study of Diabetes, European Association for the Study of Obesity NAFLD guidelines.^{4,5} They explored perceptions and experiences of current clinical practice including the diagnostic process, management of NAFLD, and recommendations for intervention and optimization of the current care pathway. All interviews were audio recorded, transcribed verbatim, and analyzed independently by 2 researchers using directed content analysis. Study design and reporting were in accordance with the “consolidated criteria for reporting qualitative research”⁶ checklist.⁶

Results

Physicians reported that the process for diagnosing and assessing NAFLD improved after implementation of local guidelines (Table 1, subtheme 1.1); however, they reported that the drivers of referral from primary to secondary care varied considerably. For example, a number of primary care physicians referred patients with abnormal liver test results or abnormal liver imaging suspecting NAFLD. They regularly provided a NAFLD score but requested advice for ongoing management and treatment, specifically about whether further investigations were required (Table 1, subtheme 1.1). Once diagnosed, a lack of knowledge and tools to deliver effective lifestyle behavior change meant that health care professionals reported monitoring rather than actively managing NAFLD (eg, annual reviews consisted of assessing disease progression) (Table 1, subthemes 2.1, 2.2, 3.1, and 3.2). Patients reported a lack of information provision after diagnosis of NAFLD, specifically relating to NAFLD severity. Furthermore, some patients were

Abbreviation used in this paper: NAFLD, nonalcoholic fatty liver disease.

© 2017 by the AGA Institute. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1542-3565

<http://dx.doi.org/10.1016/j.cgh.2017.06.011>

Table 1. Illustrative Quotations by Theme From Data Generated by Health Care Professionals and Patients With NAFLD From Primary and Secondary Care

	Illustrative quotations	
117		175
118		Q12 176
119		177
120		178
121		Q13 179
122	1.0 Theme: diagnosis	180
123	1.1 Subtheme: local guidelines	181
124	have improved the	182
125	diagnostic process	183
126	1.2 Subtheme: inadequate	Q14 184
127	information provision	185
128	about what the diagnosis	186
129	means	187
130		188
131		189
132		190
133		191
134		192
135	2.0 Theme: management	193
136	2.1 Subtheme: monitoring	194
137	vs active management	195
138	2.2 Subtheme: general	196
139	lifestyle advice rather	197
140	than tailored intervention	198
141	2.3 Subtheme: monitoring	199
142	vs management	200
143	2.4 Subtheme: information	201
144	provision about NAFLD is	202
145	lacking	203
146		204
147		205
148		206
149		207
150		208
151	3.0 Theme: recommendations	209
152	3.1 Subtheme: training to	210
153	improve knowledge of	211
154	NAFLD, diagnosis, and	212
155	the referral pathway	213
156	3.2 Subtheme: training to	214
157	improve delivery of	215
158	lifestyle interventions	216
159	3.1 Subtheme:	217
160	information needs	218
161	3.2 Subtheme: tailored	219
162	support from a	220
163	multidisciplinary team	221
164		222
165		223
166		224
167		225
168		226
169		227
170		228
171		229
172		230
173		231
174		232

Table 1. Continued

	Illustrative quotations	
233		291
234		292
235		293
236		294
237		295
238		296
239		297
240		298
241		299
242	4.0 Theme: service delivery	300
243	4.1 Subtheme:	301
244	multidisciplinary	302
245	team input is important to	303
246	tailor management of NAFLD	304
247	4.2 Subtheme: tools to support	305
248	management of NAFLD	306
249	4.3 Subtheme: approach to	307
250	management that is flexible	308
251	and offers choice	309
252		310
253		311
254		312
255		313
256		314
257		315
258		316
259		317
260		318
261		319
262		320
263		321
264		322
265		323
266		324
267		325
268		326
269		327
270		328
271		329
272		330
273	HCP-PC, health care professional from primary care; HCP-SC, health care professional from secondary care.	331
274		332
275		333
276		334
277		335
278		336
279		337
280		338
281		339
282		340
283		341
284		342
285		343
286		344
287		345
288		346
289		347
290		348

advised that NAFLD is nothing to be concerned about when compared with other health conditions (ie, health conditions such as diabetes were viewed as more important) (Table 1, subtheme 1.1). This was particularly the case in primary care. Patients reported a lack of support thereafter to manage their condition effectively. They reported being advised to lose weight but did not receive any support to do so (Table 1, subtheme 2.1) and reported not knowing that NAFLD could be improved via lifestyle modification (Table 1, subtheme 3.1). This could explain in part why the majority of patients rarely succeed with weight loss in this context. Both participant groups reported the need for a multidisciplinary team to support the management of NAFLD, including a range of lifestyle intervention options that are sensitive to the needs and preferences of patients and that help to

support long-term behavior change (Table 1, subthemes 3.2 and 4.1–4.3).

Discussion

Although evidence has shown that lifestyle interventions targeting diet and physical activity/exercise for weight loss are effective for managing NAFLD, and published clinical guidelines recommend such intervention, we have identified a substantial disconnect between guidance and how clinical care is delivered in practice. A lack of resources and training on how to target lifestyle behavior change to help manage NAFLD long term effectively was reported by health care professionals, as well as the need for a joint way of working across

disciplines to avoid miscommunication to patients. Patients further reinforced this finding, indicating that information and support at the time of diagnosis and thereafter is severely lacking.

References

1. Thoma C, Day CP, Trenell MI. Lifestyle interventions for the treatment of non-alcoholic fatty liver disease: a systematic review. *J Hepatol* 2012;56:255–266.
2. Promrat K, Kleiner DE, Niemeier H, et al. Randomized controlled trial testing the effects of weight loss on nonalcoholic steatohepatitis. *Hepatology* 2010;51:121–129.
3. Hallsworth K, Avery L, Trenell MI. Targeting lifestyle behavior change in adults with NAFLD during a 20-min consultation: summary of the dietary and exercise literature. *Curr Gastroenterol Rep* 2016;18:11.
4. European Association for the Study of the Liver, European Association for the Study of Diabetes, European Association for the Study of Obesity. EASL–EASD–EASO Clinical Practice Guidelines for the management of non-alcoholic fatty liver disease. *J Hepatol* 2016;64:1388–1402.
5. Chalasani N, Younossi ZM, Lavine JE, et al. The diagnosis and management of non-alcoholic fatty liver disease: Practice

Guideline by the American Association for the Study of Liver Diseases, American College of Gastroenterology, and the American Gastroenterological Association. *Hepatology* 2012; 55:2005–2023.

6. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care* 2007; 19:349–357.

Reprint requests

Address requests for reprints to: Kate Hallsworth, MD, MoveLab, Newcastle University, William Leech Building, 4th Floor, Newcastle upon Tyne, NE2 4HH United Kingdom. e-mail: kate.hallsworth@ncl.ac.uk; fax: (44) 191-208-5685.

Conflicts of interest

The authors disclose no conflicts.

Funding

This article reports on independent research conducted as part of Clinical Lectureship CAT CL-2013-04-010 (K.H.), which was supported by the National Institute for Health Research and Health Education England. The project also was supported in part by the facilities of the Newcastle National Institute for Health Research Biomedical Research Centre. The views expressed in this article are those of the authors and not necessarily those of the National Health Service, the National Institute for Health Research, or the Department of Health.

412
413
414
415
416
417
418
419
420
421
422
Q1Q2 423
424
425
426
Q3 427
428
429
430
431
432
433
434
Q4 435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474