

Northumbria Research Link

Citation: Gelash, Andrey, Agafontsev, Dmitry, Zakharov, Vladimir, El, Gennady, Randoux, Stéphane and Suret, Pierre (2019) Bound state soliton gas dynamics underlying the spontaneous modulational instability. *Physical Review Letters*, 123 (23). p. 234102. ISSN 0031-9007

Published by: American Physical Society

URL: <https://doi.org/10.1103/physrevlett.123.234102>
<<https://doi.org/10.1103/physrevlett.123.234102>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/41424/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

Bound state soliton gas dynamics underlying the spontaneous modulational instability

Andrey Gelash^{1,2}, Dmitry Agafontsev^{1,3}, Vladimir Zakharov^{1,4},
Gennady El⁵, Stéphane Randoux^{6,7}, and Pierre Suret^{6,7,*}

¹ Skolkovo Institute of Science and Technology, 121205, Moscow, Russia

² Novosibirsk State University, 630090, Novosibirsk, Russia

³ P. P. Shirshov Institute of Oceanology of RAS, 117218, Moscow, Russia

⁴ P. N. Lebedev Physical Institute, 53 Leninsky ave., 119991, Moscow, Russia

⁵ Northumbria University, Newcastle upon Tyne, United Kingdom

⁶ Laboratoire de Physique des Lasers, Atomes et Molécules,

UMR-CNRS 8523, Université de Lille, France and

⁷ Centre d'Etudes et de Recherches Lasers et Applications (CERLA), 59655 Villeneuve d'Ascq, France*

We investigate the fundamental phenomenon of the spontaneous, noise-induced modulational instability (MI) of a plane wave. The statistical properties of the noise-induced MI observed previously in numerical simulations and in experiments have not been explained theoretically. In this paper, using the inverse scattering transform (IST) formalism, we propose a theoretical model of the asymptotic stage of the noise-induced MI based on N -soliton solutions of the focusing one-dimensional nonlinear Schrödinger equation (1D-NLSE). Specifically, we use ensembles of N -soliton bound states having a special semi-classical distribution of the IST eigenvalues, together with random phases for norming constants. To verify our model, we employ a recently developed numerical approach to construct ensemble of N -soliton solutions with a large number of solitons, $N \sim 100$. Our investigation reveals a remarkable agreement between spectral (Fourier) and statistical properties of the long-term evolution of the MI and those of the constructed multi-soliton, random-phase bound states. Our results can be generalized to a broad class of strongly nonlinear integrable turbulence problems.

Integrable partial differential equations such as Sine-Gordon, Korteweg-de Vries (KdV) and the one-dimensional nonlinear Schrödinger equation (1D-NLSE), are considered as universal models of nonlinear physics [1]. They describe at leading order various nonlinear systems and can be integrated using the Inverse Scattering Transform (IST), often seen as a nonlinear analogue of the Fourier transform [2–6]. The traditional IST theory deals with decaying fields [6, 7], but constant non-zero boundary conditions at infinity can also be considered [8, 9]. Periodic and quasi-periodic fields can finally be examined in the framework of the *finite gap theory* [10–12]. Deterministic initial conditions have been widely studied within these three different boundary conditions.

In contrast, the propagation of *random waves* in integrable systems remains an open theoretical problem of significant applied interest due to complexity of many real world nonlinear phenomena modeled by integrable equations. For this reason, *Integrable Turbulence* (IT) has been introduced as a “new chapter of turbulence theory” by V. Zakharov [13] and is now an active theoretical and experimental field of research [14–23].

In particular, the development of the noise-induced *Modulational Instability* (MI), also known as the Benjamin-Feir instability [24] and arising in the focusing regime of the 1D-NLSE represents a prominent example of the IT phenomena [15]. This instability can be observed in many physical systems, such as deep water waves [12], BEC [25] or nonlinear optical waves [26]. In the traditional formulation, the development of MI is seen as the amplification of an initially small sinusoidal perturbation of a plane wave – the *condensate* [11, 27].

In this case, the nonlinear stage of MI is described by homoclinic solutions of the 1D-NLSE – the Akhmediev Breathers [28–31].

When the small initial perturbation of the condensate is a random process, the numerical simulations of the focusing 1D-NLSE show that the long-time evolution is characterised by a stationary single-point statistics, which is Gaussian despite the presence of highly nonlinear breather-like structures [15, 17, 32]. It has also been shown recently that this long-time (stationary) statistics is typified by a quasi-periodic structure of the spatial autocorrelation function $g^{(2)}$ of the wavefield intensity [33]. While all these remarkable features of IT have been recently demonstrated experimentally in optical fiber [33], they are still not understood theoretically.

IT can be approached from a completely different perspective which is close to classical statistical mechanics. In 1971, V. Zakharov introduced the concept of soliton gas (SG) as an infinite collection of interacting KdV solitons that are randomly distributed in space [34]. SGs have been studied theoretically in different contexts [35–38] and generated very recently in shallow water experiments [39]. Originally introduced for the case of small density [34], the notion of SG has been extended to gases of finite density for both KdV and the focusing 1D-NLSE [36]. Importantly, the macroscopic properties of dense SGs are determined by pairwise collisions of solitons accompanied by phase shifts that are accumulated at long time leading to significant corrections to the average soliton velocities. The key role in the SG theory is played by the spectral (IST) distribution function which has the meaning of the density of states $f(\lambda, x, t)$, so that

$f(\lambda_0, x_0, t) d\lambda dx$ is the number of solitons with the spectral parameter λ in the interval $[\lambda_0, \lambda_0 + d\lambda]$ found in the space interval $[x_0, x_0 + dx]$. For a spatially non-uniform SG, the evolution of f is described by a kinetic equation [34, 36].

Given the above two approaches, one can naturally pose a question about a possibility of describing IT phenomena, for instance, the development of MI, by considering the SG dynamics for some special spectral (IST) distribution. In fact, the idea to explain the nonlinear stage of MI using soliton interactions has been put forward as early as in 1972 by V. Zakharov and A. Shabat [2]. However, rather paradoxically, up to now, this possible link between soliton interactions and the MI of the plane wave has not been explored.

In this Letter, we provide a bridge between the two fundamental phenomena of nonlinear physics by showing that *SG dynamics* explains the salient features of the nonlinear stage of the *noise-induced MI*. More precisely, we demonstrate a remarkable agreement between the spectral (Fourier) and statistical properties of an unstable plane wave in the long-time evolution and those of a SG representing a random infinite-soliton bound state. The two key ingredients in our analysis are (i) a special choice of the spectral (IST) distribution in the SG and (ii) random phases of the so-called norming constants.

Without loss of generality, we consider the focusing 1D-NLSE in the standard dimensionless form

$$i \frac{\partial \psi}{\partial t} + \frac{1}{2} \frac{\partial^2 \psi}{\partial x^2} + |\psi|^2 \psi = 0, \quad (1)$$

and the plane wave solution – the condensate – of unit amplitude $\psi_c(t, x) = \exp it$. Here, ψ represents the complex field, and x and t are space and time. The classical formulation of the noise-induced MI problem is to consider the initial condition composed of the condensate with some additional noise:

$$\psi(t=0, x) = 1 + \eta(x), \quad (2)$$

where η is a small noise, $\langle |\eta|^2 \rangle \ll 1$, with zero average, $\langle \eta \rangle = 0$. The condensate is unstable with respect to long-wave perturbations with the growth rate $\Gamma(k) = k\sqrt{1 - k^2/4}$. The spectral width of the the noise is assumed to be much larger than the larger unstable wave number $k = 2$. Previously, this problem was widely investigated, both numerically by using periodic boundary conditions in a box of large size [15, 33, 40, 41], and experimentally [33, 42].

Nonlinear wave fields ruled by Eq. (1) can be characterized by the so-called scattering data (the *IST spectrum*). In the general case, the IST spectrum of spatially localized wavefields $\psi(x)$ – with zero boundary conditions – consists of the *continuous and discrete* components. A

special class of solutions, the *N-soliton solutions* (*N-SS*), exhibits only discrete spectrum consisting of N complex-valued eigenvalues λ_n , $n = 1, \dots, N$, and complex coefficients C_n , called norming constants, defined for each λ_n . The key result of the IST theory is that, while the wavefield ψ exhibits a complex dynamics, the IST spectrum evolves trivially in time [7]:

$$\forall n : \lambda_n = \text{const}, \quad C_n(t) = C_n(0) e^{-2i\lambda_n^2 t}. \quad (3)$$

The asymptotic evolution of the *N-SS* for $t \rightarrow \infty$ generally leads to a superposition of N moving solitons (1-soliton solutions) separated in space. Each soliton corresponds to a point λ_n of the discrete spectrum so that $\text{Im}\lambda_n$ is proportional to the soliton's amplitude and $\text{Re}\lambda_n$ – to its velocity. The value C_n determines the soliton phase and position.

There is a special class of *N-SSs* with zero velocities, $\text{Re}\lambda_n = 0$, the so-called bound states. In the following, we consider an ensemble of *N-SSs* with the eigenvalues located on the imaginary axis and random phases in Eq. (3), i.e. $C_n(0) = |C_n(0)| e^{i\theta_n}$, where θ_n are uniformly distributed in $[0, 2\pi)$. For $N \gg 1$, we assume the following (Weyl's) normalized distribution of the IST eigenvalues $\lambda_n = i\beta_n$:

$$\varphi(\beta) = \beta / (\sqrt{1 - \beta^2}) \quad (4)$$

We shall call the limit at $N \rightarrow \infty$ of the described random *N-SS* ensemble a *bound state SG*.

The assumption of random norming constants phases in the long-term evolution of a stochastic field is natural because the phase rotations $-2i\lambda_n^2 t$ for large t introduce an effective randomization. Random phases of norming constants are proposed here to describe IT in the framework of IST. Similarly, in wave turbulence theory, the so-called “random phase approximation” corresponds to random phases of the Fourier components of weakly dispersive waves [43, 44].

Our motivation behind the eigenvalues distribution (4) is the Bohr-Sommerfeld quantization rule for the discrete spectrum of the semiclassical Zakharov-Shabat scattering problem with the potential $\psi(x)$ in the form of a real-valued rectangular box of unit amplitude and width $L_0 \gg 1$, see e.g. [2, 7] and also [45]:

$$\lambda_n = i\beta_n = i \sqrt{1 - \left[\frac{\pi(n - \frac{1}{2})}{L_0} \right]^2}, \quad n = 1, 2, \dots, N, \quad (5)$$

where $N = \text{int}[L_0/\pi]$. Importantly, for a “semiclassical” box, the contribution of the “non-soliton” part of the field, *i.e.* of the continuous IST spectrum, decays exponentially with L_0 and so can be neglected [7]. The continuous limit of Eq. (5) with $N \rightarrow \infty$, $\beta_n \rightarrow \beta$ yields the Weyl's distribution (4) for $\varphi(\beta) = \frac{1}{N} dn/d\beta$. In this case, the density of states

is simply $f(\beta) = \frac{1}{L_0} \frac{dn}{d\beta} = \frac{1}{\pi} \varphi(\beta)$. In fact, this distribution has been recently shown to follow from a certain criticality condition for the bound state SG [59].

The derivation of the general N -SS of the 1D-NLSE is a classical result of the IST theory [2]. However, the numerical computation of N -SS with large $N \sim 100$ has been realized for the first time only in 2018 [46]. Here, we use the same approach based on the specific implementation of the so-called dressing method [47] combined with 100-digits arithmetics (see [46] and Supplemental Material).

In the following, we shall compare dynamical and statistical properties of the nonlinear stage of MI with those of a specific bound state SG. The SG is built as a random ensemble of 10^3 realisations of N -SS with $N = 128$, the spectral distribution (4) and soliton phase parameters θ_n uniformly distributed in the interval $[0, 2\pi)$. In the results reported in the main Letter, the discretisation of the Weyl distribution (4) is taken at the Bohr-Sommerfeld quantization points (5). Similar results are obtained when soliton eigenvalues are randomly distributed using the probability function (4) (see Supplemental Material).

The density of the gas, *i.e.* the number of solitons per unit length, plays a crucial role in the dynamics. Our numerical investigations have shown that the higher the density is, the better is the agreement between SG and the stationary state of MI. In the dressing method, the density is empirically controlled by “space position parameters” x_{0n} (see [46]) and reaches its maximum when all $x_{0n} = 0$. Note that, when all $x_{0n} = 0$, all $|C_n| = 1$ and the N -SS is symmetric; in order to avoid this artificial symmetry, we use a random uniform distribution of space position parameters x_{0n} in a narrow interval $[-2, 2]$, so that the soliton density remains practically unchanged.

Finally, it is important to note that the norming constants used in the dressing methods somewhat differ from those appearing in the standard IST formulation (see e.g.[48]). However, the uniform random phase distribution employed in our dressing construction of N -SS also translates to uniform distribution for the traditional IST phases.

An example of the bound state N -SS with $N = 128$ constructed according to the above described rules is shown in Fig. 1. We first compare qualitatively the temporal evolutions of this bound state N -SS and of an unstable plane waves (Fig. 2). The evolution of the two states is simulated with the pseudo-spectral Runge-Kutta fourth-order method as described in [15] in a periodic box of size $L \simeq 570$; for the MI, the noise amplitude is $\sqrt{\langle |\eta|^2 \rangle} = 10^{-5}$ (see Supplemental Material). Remarkably, the features characterizing the spatio-temporal dynamics of the N -SS and of the noise-induced MI, which is shown in Fig. 2, are qualitatively very similar; in particular, one can recognize the emergence of the well-known




Figure 1. **Example of one realization of 128-SS with random norming constant phases.** (a) Intensity profile $|\psi(x, t = 0)|^2$. (b) Soliton eigenvalues computed from Eq. (5).




Figure 2. **Numerical simulations of 1D-NLSE : Space-Time diagrams of $|\psi(x, t)|^2$.** (a) Noise-induced Modulational Instability of a plane wave. (b) Dynamics of the random phase bound N -SS (the initial condition $|\psi(x, 0)|^2$ is shown in the Fig. 1.a)

structures resembling the Akhmediev breathers.

We now compare quantitatively the statistical properties of SG and of the asymptotic stage of MI. In particular, the long-term evolution of the noise-induced MI can be characterised by stationary values of the potential H_{nl} and kinetic H_l energy, and also by stationary shapes of the (Fourier) spectrum, the probability density function (PDF) of the intensity $I = |\psi|^2$, and the autocorrelation function of the intensity $g^{(2)}$ (see [15, 49]). H_{nl} and H_l are the two parts of the total energy (Hamiltonian) E , which is one of the infinite constants of motion of the 1D-NLSE [7]:

$$E = H_l + H_{nl}, \quad H_l = \frac{1}{2} \frac{1}{L} \int_{-L/2}^{L/2} |\psi_x|^2 dx, \\ H_{nl} = -\frac{1}{2} \frac{1}{L} \int_{-L/2}^{L/2} |\psi|^4 dx. \quad (6)$$

In the case of the MI, the kinetic and potential ener-




Figure 3. Time evolution of the ensemble-averaged kinetic $\langle H_I(t) \rangle$ and potential $\langle H_{nl}(t) \rangle$ energies for the noise-induced MI (black and magenta solid lines) and random phase SG (blue and red dashed lines).

gies reach after some oscillatory transient the stationary values of $\langle H_I \rangle = 0.5$ and $\langle H_{nl} \rangle = -1$, see [15] and Fig. 3. Meanwhile, the considered bound state SG is characterized by the same stationary values of $\langle H_I \rangle$ and $\langle H_{nl} \rangle$ (dashed lines in Fig. 3) from the start of the evolution. Note that $\langle \dots \rangle$ means here ensemble-averaging over 1000 random phase realizations.

In the following, for practical reasons, we perform ensemble averaging together with temporal averaging, both for the noise-induced MI and for the N -SS. In the case of the condensate, the temporal averaging is performed when the system is sufficiently close to the asymptotic stationary state by its statistical properties ($t \in [160, 200]$). In the case of the N -SS, time averaging is started from the very beginning of the system evolution. It is extremely important to note that the results obtained using solely the ensemble averaging over random phase realizations are the same as those obtained using both ensemble and time averaging (see Supplemental Material).

As demonstrated in Fig. 4a, the wave-action spectrum,

$$S_k \propto \langle |\psi_k|^2 \rangle, \quad \psi_k = \frac{1}{L} \int_{-L/2}^{L/2} \psi e^{-ikx} dx, \quad (7)$$

of the asymptotic state of the MI and of the considered SG coincide with excellent accuracy. Note that, for the SG case, S_k is renormalized proportionally to the spatial extension of the wavefield (see Supplemental Material).

Moreover, both the SG and the noise-induced MI exhibit nearly identical PDF $\mathcal{P}(I)$ of the field intensity $I = |\psi|^2$ (Fig. 4.b). The PDF of the N -SS thus reproduces quantitatively the exponential distribution discovered earlier as the asymptotic statistics of the unstable condensate [15, 33].

As has been shown very recently in [33], the long-term evolution of the MI is typified by an oscillatory structure




Figure 4. Comparison of ensemble averaged statistical characteristics of the asymptotic state of the MI development and of random phase 128-SSs. (a) Wave action spectrum S_k . (b) The PDF $\mathcal{P}(I)$. (c) Autocorrelation function of intensity (second order degree of coherence) $g^{(2)}(x)$

of autocorrelation of the intensity $g^{(2)}(x)$:

$$g^{(2)}(x) = \frac{\langle I(y, t) I(y - x, t) \rangle}{\langle I(y, t) \rangle^2} \quad (8)$$

which represents the second-order degree of coherence. The SG reproduces accurately this remarkable oscillatory shape (see Figs. 4.c).

Note that H_I , H_{nl} , the PDF and the $g^{(2)}$ functions for the SG case were computed in the central part of the 128-SSs. More precisely, we used the region $x \in [-150; 150]$ where the ensemble-average wavefield intensity of the N -SS is practically uniform and very close to unity, that allows us to mitigate the edge effects – see details in the Supplemental materials.

In this Letter we have demonstrated that the spectral (Fourier) and statistical properties of the asymptotic state of the MI precisely coincide with those of some

specific SG. This SG can be constructed with exact N -soliton solutions of the 1D-NLSE having large values of N and one specific distribution of IST eigenvalues coinciding with the semi-classical distribution (4) for the discrete spectrum of the box potential [2]. As it could be expected, the long-term statistical state of MI corresponds to a full stochasticization of IST (soliton) phases.

Our numerical investigations reveal that spectral (Fourier) and other statistical properties (PDF and $g^{(2)}$) of SG made of stochastic N -soliton bound states are very sensitive to the exact distribution of the IST eigenvalues. An extensive study of all possible types of distributions is beyond the scope of this paper. However, it is important to note that, for all the distributions that we have tried [except for Eq. (4)], the statistical properties of the SG do not coincide with the ones of the MI and/or are strongly non homogeneous in space. For example, if the eigenvalues are uniformly distributed in the interval $[0 : i]$, the average intensity $\langle |\psi(x, t)|^2 \rangle$ of the SG strongly depends on the position x (see Supplementary). Our investigations thus strongly suggest that the distribution (4) plays a very peculiar role in SG theory because (i) it models MI and (ii) it enables to build a dense homogeneous SG, having constant statistical properties over a significant part of the space, by using finite number of solitons.

We believe that our work opens a new promising direction in the theory of integrable turbulence by establishing a link between the MI and SGs dynamics. In particular, as is known, the nonlinear stage of the MI induced by small *harmonic* perturbations is characterized by the generation of Akhmediev breathers [28–31]. The picture is more complicated if the initial perturbation is a random noise, leading to IT with various breather structures appearing only locally [15, 41, 42, 50]. Breathers also appear locally in some multi-soliton interactions [51, 52]. Our approach, based on the well-known multi-soliton solutions, can shed light on the possible connection between soliton and breather gases [32, 36, 59].

Note that the mechanism underlying the MI induced by spatially homogeneous noise studied here is *a priori* different from the MI induced by localized perturbations which are studied within the framework of IST with nonzero boundary conditions [49, 53–58]. In this work, we have demonstrated that the IST formalism for wavefields with *decaying* boundary conditions can be successfully applied to describe accurately the asymptotic stationary state of the MI computed numerically by using *periodic* boundary conditions [15]. An important mathematical problem for the future studies is to explain the link between spatially periodic and localised descriptions of the MI in terms of the IST theory.

Finally note that, our model can be generalised to a broad class of IT problems when the (random) wavefield is strongly nonlinear, so that the impact of the non-soliton content can be neglected in the asymptotic state

($t \rightarrow \infty$). For such a case, our work suggests a general strategy to study the asymptotic state: build N -soliton solutions with the distribution $\varphi(\lambda)$ of IST eigenvalues characterizing the field and *random phases* of the norming constants.

ACKNOWLEDGMENTS

The authors thank A. Tikan and F. Copie for fruitful discussions. Simulations were performed at the Novosibirsk Supercomputer Center (NSU). This work has been partially supported by the Agence Nationale de la Recherche through the LABEX CEMPI project (ANR-11-LABX-0007) and by the Ministry of Higher Education and Research, Nord-Pas de Calais Regional Council and European Regional Development Fund (ERDF) through the Contrat de Projets Etat-Région (CPER Photonics for Society P4S). The work on the construction of multi-soliton ensembles reported in the first half of the paper was supported by the Russian Science Foundation (Grant No. 19-72-30028 to AG, DA and VZ). The work of GE was partially supported by EPSRC grant EP/R00515X/2.

* Corresponding author : Pierre.Suret@univ-lille.fr

- [1] J. Yang, *Nonlinear waves in integrable and nonintegrable systems*, Vol. 16 (SIAM, 2010).
- [2] V Zakharov and A Shabat, “Exact theory of two-dimensional self-focusing and one-dimensional self-modulation of waves in nonlinear media,” *Soviet physics JETP* **34**, 62 (1972).
- [3] V.E. Zakharov and A.B. Shabat, “A scheme for integrating the nonlinear equations of mathematical physics by the method of the inverse scattering problem. I,” *Functional analysis and its applications* **8**, 226–235 (1974).
- [4] M.J. Ablowitz, D.J. Kaup, A.C. Newell, and H. Segur, “The inverse scattering transform-fourier analysis for nonlinear problems,” *Studies in Applied Mathematics* **53**, 249–315 (1974).
- [5] VE Zakharov and AB Shabat, “Integration of nonlinear equations of mathematical physics by the method of inverse scattering. ii,” *Functional Analysis and Its Applications* **13**, 166–174 (1979).
- [6] M. J. Ablowitz and H. Segur, *Solitons and the inverse scattering transform*, Vol. 4 (Siam, 1981).
- [7] S Novikov, SV Manakov, LP Pitaevskii, and VE Zakharov, *Theory of solitons: the inverse scattering method* (Springer Science & Business Media, 1984).
- [8] Tutomu Kawata and Hiroshi Inoue, “Inverse scattering method for the nonlinear evolution equations under nonvanishing conditions,” *Journal of the Physical Society of Japan* **44**, 1722–1729 (1978).
- [9] Ludwig Faddeev and Leon Takhtajan, *Hamiltonian methods in the theory of solitons* (Springer Science & Business Media, 2007).
- [10] ED Belokolos, AI Bobenko, VZ Enol’skii, AR Its, and

- VB Matveev, “Algebro-geometric approach to nonlinear integrable equations,” (1994).
- [11] E. R. Tracy and H. H. Chen, “Nonlinear self-modulation: An exactly solvable model,” *Phys. Rev. A* **37**, 815–839 (1988).
- [12] A. Osborne, *Nonlinear ocean waves* (Academic Press, 2010).
- [13] VE Zakharov, “Turbulence in integrable systems,” *Stud. Appl. Math.* **122**, 219–234 (2009).
- [14] Pierre Walczak, Stéphane Randoux, and Pierre Suret, “Optical rogue waves in integrable turbulence,” *Phys. Rev. Lett.* **114**, 143903 (2015).
- [15] DS Agafontsev and Vladimir E Zakharov, “Integrable turbulence and formation of rogue waves,” *Nonlinearity* **28**, 2791 (2015).
- [16] M. Onorato, D. Proment, GA El, S. Randoux, and P. Suret, “On the origin of heavy-tail statistics in equations of the nonlinear schrodinger type,” *Physics Letters A* **380**, 173–3177 (2016).
- [17] J. M. Soto-Crespo, N. Devine, and N. Akhmediev, “Integrable turbulence and rogue waves: Breathers or solitons?” *Phys. Rev. Lett.* **116**, 103901 (2016).
- [18] S. Randoux, P. Walczak, M. Onorato, and P. Suret, “Nonlinear random optical waves: integrable turbulence, rogue waves and intermittency,” *Physica D: Nonlinear Phenomena* **333**, 323–335 (2016).
- [19] G. Roberti, GA El, S. Randoux, and P. Suret, “Early stage of integrable turbulence in 1D NLS equation: the semi-classical approach to statistics,” accepted for publication in *Phys. Rev. E*, arXiv:1901.02501 (2019).
- [20] P. Suret, R. El Koussaifi, A. Tikan, C. Evain, S. Randoux, C. Szwarz, and Serge Bielawski, “Single-shot observation of optical rogue waves in integrable turbulence using time microscopy,” *Nat. Commun.* **7** (2016).
- [21] A. Tikan, S. Bielawski, C. Szwarz, Stéphane Randoux, and P. Suret, “Single-shot measurement of phase and amplitude by using a heterodyne time-lens system and ultrafast digital time-holography,” *Nature Photonics* **12**, 228 (2018).
- [22] R. El Koussaifi, A. Tikan, A. Toffoli, S. Randoux, P. Suret, and M. Onorato, “Spontaneous emergence of rogue waves in partially coherent waves: a quantitative experimental comparison between hydrodynamics and optics,” *Physical Review E* **97**, 012208 (2018).
- [23] A. Cazaubiel, G. Michel, S. Lepot, B. Semin, S. Aumaitre, M. Berhanu, F. Bonnefoy, and E. Falcon, “Coexistence of solitons and extreme events in deep water surface waves,” *Physical Review Fluids* **3**, 114802 (2018).
- [24] T Brooke Benjamin and JE Feir, “The disintegration of wave trains on deep water part 1. theory,” *Journal of Fluid Mechanics* **27**, 417–430 (1967).
- [25] Kevin E Strecker, Guthrie B Partridge, Andrew G Truscott, and Randall G Hulet, “Formation and propagation of matter-wave soliton trains,” *Nature* **417**, 150 (2002).
- [26] G. P. Agrawal, *Nonlinear Fiber optics*, 3rd ed., Optics and Photonics (Academic Press, 2001).
- [27] VE Zakharov and LA Ostrovsky, “Modulation instability: the beginning,” *Physica D: Nonlinear Phenomena* **238**, 540–548 (2009).
- [28] NN Akhmediev, VM Eleonskii, and NE Kulagin, “Generation of periodic trains of picosecond pulses in an optical fiber: exact solutions,” *Sov. Phys. JETP* **62**, 894–899 (1985).
- [29] NN Akhmediev and VI Korneevev, “Modulation instability and periodic solutions of the nonlinear schrödinger equation,” *Theoretical and Mathematical Physics* **69**, 1089–1093 (1986).
- [30] N. Akhmediev, A. Ankiewicz, and M. Taki, “Waves that appear from nowhere and disappear without a trace,” *Phys. Lett. A* **373**, 675 – 678 (2009).
- [31] PG Grinevich and PM Santini, “The finite gap method and the analytic description of the exact rogue wave recurrence in the periodic nls cauchy problem. 1,” *Nonlinearity* **31**, 5258 (2018).
- [32] N Akhmediev, JM Soto-Crespo, and N Devine, “Breather turbulence versus soliton turbulence: Rogue waves, probability density functions, and spectral features,” *Physical Review E* **94**, 022212 (2016).
- [33] Adrien E. Kraych, Dmitry Agafontsev, Stéphane Randoux, and Pierre Suret, “Statistical properties of the nonlinear stage of modulation instability in fiber optics,” *Phys. Rev. Lett.* **123**, 093902 (2019).
- [34] VE Zakharov, *Sov. Phys. JETP* **33**, 538–540 (1971).
- [35] VS Gerdjikov, DJ Kaup, IM Uzunov, and EG Evstatiev, “Asymptotic behavior of n-soliton trains of the nonlinear schrödinger equation,” *Physical review letters* **77**, 3943 (1996).
- [36] G. A. El and A. M. Kamchatnov, *Phys. Rev. Lett.* **95**, 204101 (2005).
- [37] F. Carbone, D. Dutykh, and GA. El, *EPL (Europhysics Letters)* **113**, 30003 (2016).
- [38] Efim Pelinovsky and Ekaterina Shurgalina (Springer, 2017) pp. 295–306.
- [39] Ivan Redor, Eric Barthélemy, Hervé Michallet, Miguel Onorato, and Nicolas Mordant, “Experimental evidence of a hydrodynamic soliton gas,” *Phys. Rev. Lett.* **122**, 214502 (2019).
- [40] John M. Dudley, Frederic Dias, Miro Erkintalo, and Goery Genty, “Instabilities, breathers and rogue waves in optics,” *Nat. Photon.* **8**, 755 (2014).
- [41] Shanti Toenger, Thomas Godin, Cyril Billet, Frederic Dias, Miro Erkintalo, Goery Genty, and John M. Dudley, “Emergent rogue wave structures and statistics in spontaneous modulation instability,” *Sci. Rep.* **5** (2015).
- [42] Mikko Närhi, Benjamin Wetzal, Cyril Billet, Shanti Toenger, Thibaut Sylvestre, Jean-Marc Merolla, Roberto Morandotti, Frederic Dias, Goery Genty, and John M Dudley, “Real-time measurements of spontaneous breathers and rogue wave events in optical fibre modulation instability,” *Nat. Commun.* **7** (2016).
- [43] V.E. Zakharov, V.S. L’vov, and G. Falkovich, *Kolmogorov Spectra of Turbulence I*, Berlin (Springer, 1992).
- [44] Sergey. Nazarenko, *Wave Turbulence. 10.1007/978-3-642-15942-8*, Lecture Notes in Physics (Springer Berlin Heidelberg, Berlin, Heidelberg, 2011).
- [45] ZV Lewis, “Semiclassical solutions of the zaharov-shabat scattering problem for phase modulated potentials,” *Physics Letters A* **112**, 99–103 (1985).
- [46] A. A. Gelash and D. S. Agafontsev, “Strongly interacting soliton gas and formation of rogue waves,” *Phys. Rev. E* **98**, 042210 (2018).
- [47] V. E. Zakharov and A. V. Mikhailov, “Relativistically invariant two-dimensional models of field theory which are integrable by means of the inverse scattering problem method,” *Soviet Physics JETP* **47** (1978).
- [48] Yu-Hao Sun, “Soliton synchronization in the focusing nonlinear schrödinger equation,” *Physical Review E* **93**,

- 052222 (2016).
- [49] Adrien E. Kraych, Pierre Suret, Gennady El, and Stéphane Randoux, “Nonlinear evolution of the locally induced modulational instability in fiber optics,” *Phys. Rev. Lett.* **122**, 054101 (2019).
- [50] Stefan Wabnitz and Benjamin Wetzol, “Instability and noise-induced thermalization of fermi–pasta–ulam recurrence in the nonlinear schrödinger equation,” *Physics Letters A* **378**, 2750–2756 (2014).
- [51] A. Tikan, Cyril Billet, Gennady El, Alexander Tovbis, Marco Bertola, Thibaut Sylvestre, Francois Gustave, Stephane Randoux, Goëry Genty, P. Suret, and John M. Dudley, “Universality of the peregrine soliton in the focusing dynamics of the cubic nonlinear schrödinger equation,” *Phys. Rev. Lett.* **119**, 033901 (2017).
- [52] Gennady A El, Eduardo G Khamis, and Alexander Tovbis, “Dam break problem for the focusing nonlinear schrödinger equation and the generation of rogue waves,” *Nonlinearity* **29**, 2798 (2016).
- [53] V. E. Zakharov and A. A. Gelash, “Nonlinear stage of modulation instability,” *Phys. Rev. Lett.* **111**, 054101 (2013).
- [54] B. Kibler, A. Chabchoub, A. Gelash, N. Akhmediev, and V. E. Zakharov, “Superregular breathers in optics and hydrodynamics: Omnipresent modulation instability beyond simple periodicity,” *Phys. Rev. X* **5**, 041026 (2015).
- [55] G. Biondini, S. Li, and D. Mantzavinos, “Oscillation structure of localized perturbations in modulationally unstable media,” *Phys. Rev. E* **94**, 060201(R) (2016).
- [56] Gino Biondini and Dionyssios Mantzavinos, “Universal nature of the nonlinear stage of modulational instability,” *Physical review letters* **116**, 043902 (2016).
- [57] AA Gelash, “Formation of rogue waves from a locally perturbed condensate,” *Physical Review E* **97**, 022208 (2018).
- [58] Matteo Conforti, Sitai Li, Gino Biondini, and Stefano Trillo, “Auto-modulation versus breathers in the nonlinear stage of modulational instability,” *Optics letters* **43**, 5291–5294 (2018).
- [59] G.A. El and A. Tovbis, Spectral theory of soliton and breather gases in the focusing nonlinear Schrödinger equation, arXiv:1910.05732 [nlin.PS]