

Northumbria Research Link

Citation: Das, Kinkar, Çevik, Ahmet, Cangul, Ismail and Shang, Yilun (2021) On Sombor Index. *Symmetry*, 13 (1). p. 140. ISSN 2073-8994

Published by: MDPI

URL: <https://doi.org/10.3390/sym13010140> <<https://doi.org/10.3390/sym13010140>>

This version was downloaded from Northumbria Research Link:
<http://nrl.northumbria.ac.uk/id/eprint/45243/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

Article

On Sombor Index

Kinkar Chandra Das ^{1,*}, Ahmet Sinan Çevik ², Ismail Naci Cangul ³ and Yilun Shang ^{4,*}

¹ Department of Mathematics, Sungkyunkwan University, Suwon 16419, Korea

² Department of Mathematics, Faculty of Science, Selçuk University, Campus, 42075 Konya, Turkey; sinan.cevik@selcuk.edu.tr

³ Department of Mathematics, Faculty of Art and Science, Bursa Uludag University, Gorukle, 16059 Bursa, Turkey; cangul@uludag.edu.tr

⁴ Department of Computer and Information Sciences, Northumbria University, Newcastle NE1 8ST, UK

* Correspondence: kinkardas2003@gmail.com (K.C.D.); yilun.shang@northumbria.ac.uk (Y.S.)

Abstract: The concept of Sombor index (SO) was recently introduced by Gutman in the chemical graph theory. It is a vertex-degree-based topological index and is denoted by Sombor index SO : $SO = SO(G) = \sum_{v_i, v_j \in E(G)} \sqrt{d_G(v_i)^2 + d_G(v_j)^2}$, where $d_G(v_i)$ is the degree of vertex v_i in G . Here, we present novel lower and upper bounds on the Sombor index of graphs by using some graph parameters. Moreover, we obtain several relations on Sombor index with the first and second Zagreb indices of graphs. Finally, we give some conclusions and propose future work.

Keywords: graph; sombor index; maximum degree; minimum degree; independence number

MSC: 05C50; 05C12; 15A18

1. Introduction

Consider a simple graph $G = (V, E)$ with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$ and edge set $E = E(G)$, where $|V(G)| = n$ is the number of vertices and $|E(G)| = m$ is the number of edges. For $i = 1, 2, \dots, n$ denote by $d_G(v_i)$ the degree of vertex v_i . For $v_i \in V(G)$, let $\mu_G(v_i)$ represent the *average degree* of the vertices adjacent to vertex v_i . Let δ be the *minimum vertex degree* and Δ be the *maximum vertex degree*. It is known that any vertex v of degree 1 is a *pendant vertex*. Pendant vertex is also called *leaf*. A *pendant edge* is the edge incident with a pendant vertex. We denote by $v_i v_j \in E(G)$ when vertices v_i and v_j are adjacent.

In graph theory, a number that is invariant under graph automorphisms is referred to as a graphical invariant. It is often regarded as a structural invariant relevant to a graph. The term topological index is often reserved for graphical invariant in molecular graph theory. In the mathematical and chemical literature, several dozens of vertex-degree-based graph invariants (usually referred to as “topological indices”) have been introduced and extensively studied. Their general formula is

$$TI = TI(G) = \sum_{v_i, v_j \in E(G)} F(d_G(v_i), d_G(v_j)), \quad (1)$$

where $F(x, y)$ is some function with the property $F(x, y) = F(y, x)$. If $F(d_G(v_i), d_G(v_j)) = d_G(v_i) + d_G(v_j)$ or $d_G(v_i) d_G(v_j)$, then TI is the first Zagreb index or the second Zagreb index of graph G , respectively, which are put forward in [1] by Gutman and Trinajstić. They studied the dependence of total π -electron energy related to molecular structure. Some further development can be found for example in [2]. Given a molecular graph G , we have the *first Zagreb index* $M_1(G)$ as

$$M_1(G) = \sum_{v_i, v_j \in E(G)} (d_G(v_i) + d_G(v_j)) = \sum_{v_i \in V(G)} d_G(v_i)^2$$



Citation: Das, K.C.; Çevik, A.S.; Cangul, I.N.; Shang, Y. On Sombor Index. *Symmetry* **2021**, *13*, 140. <https://doi.org/10.3390/sym13010140>

Received: 19 December 2020

Accepted: 14 January 2021

Published: 16 January 2021

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

and the *second Zagreb index* $M_2(G)$ as

$$M_2(G) = \sum_{v_i v_j \in E(G)} d_G(v_i) d_G(v_j).$$

Many fundamental mathematical properties such as lower and upper bounds involving other important graphical invariants can be bound in, e.g., [3–15]. More recent results are reported in [16–25]. Zagreb indices characterize the degree of branching in molecular carbon-atom skeleton and are regarded as powerful molecular structure-descriptors [26,27].

Gutman mentioned a list of topological indices (26 indices including two Zagreb indices) in [28]. In the same paper, Gutman presented a novel approach to the vertex-degree-based topological index of (molecular) graphs. For this we need the following definition:

Definition 1 ([28]). *The ordered pair (x, y) , where $x = d_G(v_i)$, $y = d_G(v_j)$, is the degree-coordinate (or d -coordinate) of the edge $v_i v_j \in E(G)$. In the (2-dimensional) coordinate system, it pertains to a point called the degree-point (or d -point) of the edge $v_i v_j \in E(G)$. The point with coordinates (y, x) is the dual-degree-point (or dd -point) of the edge $v_i v_j \in E(G)$. The distance between the d -point (x, y) and the origin of the coordinate system is the degree-radius (or d -radius) of the edge $v_i v_j \in E(G)$, denoted by $r(x, y)$. Based on elementary geometry (using Euclidean metrics), we have $r(x, y) = \sqrt{x^2 + y^2}$. From this, we immediately see that a d -point and the corresponding dd -point have equal degree-radii. One can easily see that for any molecular graphs ($d_G(v) \leq 4$), two degree-points have equal degree-radii if and only if they coincide, that is, if and only if both have the same degree-coordinates. Unfortunately, this property is not valid for general graphs.*

Since the function $F(x, y) = \sqrt{x^2 + y^2}$ has not been used before in the theory of vertex-degree-based topological indices, from the above considerations motivated by the author in [28], introduce a new such index defined as

$$SO = SO(G) = \sum_{v_i v_j \in E(G)} \sqrt{d_G(v_i)^2 + d_G(v_j)^2}$$

and called the Sombor index. In the same paper, several mathematical properties have been discussed.

Given $W \subseteq E(G)$, we denote by $G - W$ the subgraph of G which is obtained by removing any edge within W . If $W = \{v_i v_j\}$, we will write $G - v_i v_j$ instead of the subgraph $G - W$ for ease of expression. For a pair of nonadjacent vertices v_i and v_j in G , we write $G + v_i v_j$ for the graph obtained by adding the edge $v_i v_j$ to G . Let $G[S]$ be the induced subgraph of G by $S \subseteq V(G)$. S is said to be an independent set of G if $G[S]$ is formed by $|S|$ isolated vertices. The number of vertices in the largest independent set is called the *independence number* of a given graph, which is denoted conventionally by α . If a graph on n vertices contains a clique of $n - \alpha$ vertices and the rest α vertices is a stable set, where every vertex within the clique is linked to every vertex in the stable set, then the graph is called a complete split graph and is denoted by $CS(n, \alpha)$, $1 \leq \alpha \leq n - 1$. Another interesting graph class is called (Δ, δ) -semiregular bipartite graph, where G is a bipartite graph with a bipartition U and W . Here, each vertex v_i in U admits constant degree Δ while each vertex v_j in W admits constant degree δ . Clearly, if $\Delta = \delta$, G becomes regular. As usual, K_n is a complete graph and $K_{a,b}$ with $(a + b = n)$ is a complete bipartite graph over n vertices. We refer the reader to the book [29] for other standard graph theoretical notations.

The rest of the paper is organized as follows. In Section 2, we obtain some lower and upper bounds on $SO(G)$ in terms of graph parameters. In Section 3, we present some relations between $SO(G)$ and the Zagreb indices $M_1(G)$ and $M_2(G)$. In Section 4, we give some conclusions and future work.

2. Bounds on Sombor Index of Graphs

In this section, we give several lower and upper bounds on $SO(G)$ building on some useful graph parameters. From the definition of Sombor index, the following result can be summarized.

Lemma 1. For a graph G , we have

- (i) $SO(G) > SO(G - e)$, where $e = v_i v_j$ is any edge in G ,
- (ii) $SO(G + e) > SO(G)$, where $e = v_i v_j$ and vertices v_i & v_j are non-adjacent in G .

First we give the upper and lower bounds on $SO(G)$ building on n , Δ and δ .

Theorem 1. Suppose that G is a graph over n vertices. If G has maximum degree Δ and minimum degree δ ,

$$\frac{n \delta^2}{\sqrt{2}} \leq SO(G) \leq \frac{n \Delta^2}{\sqrt{2}}$$

with equality (left and right) if and only if G becomes a regular graph.

Proof. Recall that Δ is the maximum degree of G and δ is the minimum degree of G . By employing the Handshaking lemma, we obtain

$$n \delta \leq \sum_{v_i \in V(G)} d_G(v_i) = 2m \leq n \Delta$$

with equality holding (left and right) if and only if $d_G(v_i) = \Delta$ for any $v_i \in V(G)$. It follows from the definition of the Sombor index, we obtain

$$\begin{aligned} SO(G) &= \sum_{v_i v_j \in E(G)} \sqrt{d_G(v_i)^2 + d_G(v_j)^2} \\ &\leq \sqrt{2} \Delta m \leq \frac{n \Delta^2}{\sqrt{2}}. \end{aligned}$$

Moreover, the equality herein holds if and only if $d_G(v_i) = \Delta$ for any $v_i \in V(G)$, i.e., G is regular. Similarly, we get the lower bound on Sombor index and equality holds if and only if G becomes regular. \square

Since $\Delta \leq n - 1$, we get the following corollary.

Corollary 1 ([28]). Let G be a graph of order n . Then

$$SO(G) \leq \frac{n(n-1)^2}{\sqrt{2}}$$

with equality if and only if $G \cong K_n$.

For triangle-free graph G , we obtain an upper bound on $SO(G)$ based on n , m , Δ and δ .

Theorem 2. Let G be a triangle-free graph of order n with m edges and maximum degree Δ , minimum degree δ . Then

$$SO(G) \leq \begin{cases} m \sqrt{\delta^2 + (n - \delta)^2} & \text{if } \Delta + \delta \leq n, \\ m \sqrt{\Delta^2 + (n - \Delta)^2} & \text{if } \Delta + \delta \geq n. \end{cases}$$

Proof. Let $d_G(v_i)$ be the degree of the vertex v_i in G . Since G is triangle-free graph, we have $d_G(v_i) + d_G(v_j) \leq n$ for any edge $v_i v_j \in E(G)$. Let us consider a function

$h(x) = x^2 + (n - x)^2$ for $\delta \leq x \leq \Delta$. Then one can easily see that $h(x)$ is an increasing function on $n/2 \leq x \leq \Delta$ and a decreasing function on $\delta \leq x \leq n/2$. Hence

$$d_G(v_i)^2 + (n - d_G(v_i))^2 \leq \begin{cases} \sqrt{\delta^2 + (n - \delta)^2} & \text{if } \Delta + \delta \leq n, \\ \sqrt{\Delta^2 + (n - \Delta)^2} & \text{if } \Delta + \delta \geq n. \end{cases}$$

With the results obtained above, we derive

$$\begin{aligned} SO(G) &= \sum_{v_i v_j \in E(G)} \sqrt{d_G(v_i)^2 + d_G(v_j)^2} \\ &\leq \sum_{v_i v_j \in E(G)} \sqrt{d_G(v_i)^2 + (n - d_G(v_i))^2} \\ &\leq \begin{cases} m \sqrt{\delta^2 + (n - \delta)^2} & \text{if } \Delta + \delta \leq n, \\ m \sqrt{\Delta^2 + (n - \Delta)^2} & \text{if } \Delta + \delta \geq n. \end{cases} \end{aligned}$$

□

Gutman [28] proved that the path P_n gives the minimum value of Sombor index for any connected graph of order n . Therefore, the path P_n gives the minimum value of Sombor index for any connected bipartite graph of order n . We now give an upper bound on the Sombor index of bipartite graphs.

Theorem 3. *Let G be a bipartite graph over n vertices. Then*

$$SO(G) \leq \begin{cases} \frac{n^3}{4\sqrt{2}} & \text{if } n \text{ is even,} \\ \frac{(n^2 - 1)\sqrt{n^2 + 1}}{4\sqrt{2}} & \text{if } n \text{ is odd,} \end{cases}$$

with equality if and only if $G \cong K_{\lceil \frac{n}{2} \rceil, \lfloor \frac{n}{2} \rfloor}$.

Proof. Let G be a bipartite graph of order n ($n = p + q$, $p \geq q$) with two partite sets having p and q vertices, respectively. Since G is bipartite graph, by Lemma 1, we obtain $SO(G) \leq SO(K_{p,q})$ with equality if and only if $G \cong K_{p,q}$. Hence

$$SO(G) \leq S(K_{p,q}) = pq \sqrt{p^2 + q^2} = p(n - p) \sqrt{p^2 + (n - p)^2}.$$

Let us consider a function

$$f(x) = x(n - x) \sqrt{x^2 + (n - x)^2}, \quad \left\lceil \frac{n}{2} \right\rceil \leq x \leq n - 1.$$

Then

$$f'(x) = -\frac{(2x - n)[x^2 + (n - x)^2 - 1]}{\sqrt{x^2 + (n - x)^2}} \leq 0 \text{ for } \left\lceil \frac{n}{2} \right\rceil \leq x \leq n - 1.$$

Thus $f(x)$ is a decreasing function on $\lceil \frac{n}{2} \rceil \leq x \leq n - 1$, and hence

$$SO(G) \leq p(n - p) \sqrt{p^2 + (n - p)^2} \leq \left\lceil \frac{n}{2} \right\rceil \left\lfloor \frac{n}{2} \right\rfloor \sqrt{\left\lceil \frac{n}{2} \right\rceil^2 + \left\lfloor \frac{n}{2} \right\rfloor^2}.$$

The required inequality has been proved. Besides, the equality holds if and only if $G \cong K_{\lfloor \frac{n}{2} \rfloor, \lfloor \frac{n}{2} \rfloor}$. \square

Next, we present an upper bound on $SO(G)$ by using n and independence number α .

Theorem 4. *Let G be a connected graph of order n with independence number α . Then $SO(G) \leq \sqrt{2} \binom{n-\alpha}{2} (n-1) + \alpha (n-\alpha) \sqrt{(n-1)^2 + (n-\alpha)^2}$ with equality if and only if $G \cong CS(n, \alpha)$.*

Proof. We obtain

$$SO(CS(n, \alpha)) = \sqrt{2} \binom{n-\alpha}{2} (n-1) + \alpha (n-\alpha) \sqrt{(n-1)^2 + (n-\alpha)^2}.$$

Since G has order n and independence number α , by Lemma 1, we obtain

$$SO(G) \leq SO(CS(n, \alpha)) = \sqrt{2} \binom{n-\alpha}{2} (n-1) + \alpha (n-\alpha) \sqrt{(n-1)^2 + (n-\alpha)^2}$$

with equality if and only if $G \cong CS(n, \alpha)$. \square

We now offer an additional upper bound on $SO(G)$ in terms of m , δ and $M_1(G)$.

Theorem 5. *Let G be a graph of size m and minimum degree δ . Then*

$$SO(G) \leq M_1(G) - (2 - \sqrt{2}) \delta m,$$

where $M_1(G)$ is the first Zagreb index of graph G . Moreover, the equality holds if and only if G is a regular graph.

Proof. For any edge $v_i v_j \in E(G)$ ($d_G(v_i) \geq d_G(v_j)$), one can easily check that

$$\sqrt{d_G(v_i)^2 + d_G(v_j)^2} \leq d_G(v_i) + (\sqrt{2} - 1) d_G(v_j)$$

with equality if and only if $d_G(v_i) = d_G(v_j)$. Now,

$$\begin{aligned} SO(G) &= \sum_{v_i v_j \in E(G)} \sqrt{d_G(v_i)^2 + d_G(v_j)^2} \\ &\leq \sum_{\substack{v_i v_j \in E(G) \\ d_G(v_i) \geq d_G(v_j)}} (d_G(v_i) + (\sqrt{2} - 1) d_G(v_j)) \\ &= \sum_{v_i v_j \in E(G)} (d_G(v_i) + d_G(v_j)) - \sum_{\substack{v_i v_j \in E(G) \\ d_G(v_i) \geq d_G(v_j)}} (2 - \sqrt{2}) d_G(v_j) \\ &\leq M_1(G) - (2 - \sqrt{2}) \delta m. \end{aligned}$$

Moreover, the above two inequalities are equalities if and only if G is a regular graph. \square

We give an upper bound on $SO(G) + SO(\overline{G})$ by employing the number n only.

Theorem 6. *Let G be a graph over n vertices. We have*

$$SO(G) + SO(\overline{G}) \leq \frac{n(n-1)^2}{\sqrt{2}}$$

with equality if and only if $G \cong K_n$ or $G \cong \overline{K}_n$.

Proof. Since $\Delta \leq n - 1$, it can be easily checked that

$$\sqrt{d_G(v_i)^2 + d_G(v_j)^2} \leq \sqrt{2}(n - 1) \text{ for any edge } v_i v_j \in E(G).$$

Since $|E(G)| + |E(\overline{G})| = \frac{n(n-1)}{2}$, we obtain

$$\begin{aligned} SO(G) + SO(\overline{G}) &= \sum_{v_i v_j \in E(G)} \sqrt{d_G(v_i)^2 + d_G(v_j)^2} + \sum_{v_i v_j \in E(\overline{G})} \sqrt{d_{\overline{G}}(v_i)^2 + d_{\overline{G}}(v_j)^2} \\ &\leq \frac{n(n-1)}{2} \times \sqrt{2}(n-1) = \frac{n(n-1)^2}{\sqrt{2}}. \end{aligned}$$

Moreover, the equality holds if and only if $d_G(v_i)^2 + d_G(v_j)^2 = 2(n-1)^2$ for any edge $v_i v_j \in E(G)$ or $d_{\overline{G}}(v_i)^2 + d_{\overline{G}}(v_j)^2 = 2(n-1)^2$ for any edge $v_i v_j \in E(\overline{G})$, that is, $G \cong K_n$ or $G \cong \overline{K}_n$. \square

3. Relation between Sombor Index with Zagreb Indices of Graphs

Topological indices in mathematical chemistry are well studied in the literature. In particular we have seen several mathematical and chemical properties on topological indices of graphs, some of them are very similar, but some of them are totally different. So it is natural to ask how two topological indices are related or to find some relations between two topological indices of graphs. In last 10 years several papers have been published on this topic in the literature, see [23,25,30–37]. In this section we try to find some relations between Sombor index and the (first & second) Zagreb indices of graphs. For this we need the following result:

Lemma 2. [38] Let a_1, a_2, \dots, a_k and b_1, b_2, \dots, b_k be real numbers so that there are constants s and t satisfying for any $i, i = 1, 2, \dots, k$ we have $s a_i \leq b_i \leq t a_i$. Then

$$\sum_{i=1}^k b_i^2 + s t \sum_{i=1}^k a_i^2 \leq (s + t) \sum_{i=1}^k a_i b_i \quad (2)$$

with equality if and only if for at least one $i, 1 \leq i \leq k$ holds $s a_i = b_i = t a_i$.

Next, we investigate the relation between Sombor index $SO(G)$ and the first Zagreb index $M_1(G)$ of graph G .

Theorem 7. Let G be a graph containing n vertices and m edges. The maximum degree is denoted by Δ and its minimum degree is $\delta > 0$. Then

$$\left(\sqrt{2(\Delta^2 + \delta^2)} + \Delta + \delta\right) SO(G) \geq \sqrt{\Delta^2 + \delta^2} M_1(G) + \sqrt{2}(\Delta + \delta) \delta m, \quad (3)$$

where $M_1(G)$ is the first Zagreb index of graph G . Moreover, the equality holds if and only if G is a regular graph.

Proof. One can easily see that

$$\frac{\sqrt{d_G(v_i)^2 + d_G(v_j)^2}}{d_G(v_i) + d_G(v_j)} \geq \frac{1}{\sqrt{2}}, \text{ that is, } \left(d_G(v_i) - d_G(v_j)\right)^2 \geq 0$$

with equality if and only if $d_G(v_i) = d_G(v_j)$. Since $0 < \delta \leq d_G(v_i) \leq \Delta$, for any $v_i \in V(G)$, we have

$$\frac{\delta}{\Delta} \leq \frac{d_G(v_i)}{d_G(v_j)} \leq \frac{\Delta}{\delta} \quad (4)$$

with right (left) equality if and only if $d_G(v_i) = \Delta$ and $d_G(v_j) = \delta$ ($d_G(v_i) = \delta$ and $d_G(v_j) = \Delta$). Let $f(x) = \frac{\sqrt{1+x^2}}{1+x}$, $x \geq 1$. Then we have

$$f'(x) = \frac{x-1}{\sqrt{1+x^2}(1+x)^2}, \quad x \geq 1.$$

Thus $f(x)$ is an increasing function on $x \geq 1$. Using the above results, we obtain

$$\frac{1}{\sqrt{2}} \leq \frac{\sqrt{d_G(v_i)^2 + d_G(v_j)^2}}{d_G(v_i) + d_G(v_j)} = \frac{\sqrt{1 + \frac{d_G(v_i)^2}{d_G(v_j)^2}}}{1 + \frac{d_G(v_i)}{d_G(v_j)}} \leq \frac{\sqrt{\Delta^2 + \delta^2}}{\Delta + \delta}.$$

For any edge $v_i v_j \in E(G)$, one can easily check that

$$\frac{d_G(v_i)^2 + d_G(v_j)^2}{d_G(v_i) + d_G(v_j)} \geq \delta$$

with equality if and only if $d_G(v_i) = \delta = d_G(v_j)$, that is,

$$\sum_{v_i v_j \in E(G)} \frac{d_G(v_i)^2 + d_G(v_j)^2}{d_G(v_i) + d_G(v_j)} \geq m \delta \tag{5}$$

with equality if and only if G is a regular graph.

Setting $s = \frac{1}{\sqrt{2}}$, $t = \frac{\sqrt{\Delta^2 + \delta^2}}{\Delta + \delta}$, $a_i \rightarrow \sqrt{d_G(v_i) + d_G(v_j)}$ and $b_i \rightarrow \sqrt{\frac{d_G(v_i)^2 + d_G(v_j)^2}{d_G(v_i) + d_G(v_j)}}$ in Lemma 2, we obtain

$$\begin{aligned} \sum_{v_i v_j \in E(G)} \frac{d_G(v_i)^2 + d_G(v_j)^2}{d_G(v_i) + d_G(v_j)} + \frac{\sqrt{\Delta^2 + \delta^2}}{\sqrt{2}(\Delta + \delta)} \sum_{v_i v_j \in E(G)} (d_G(v_i) + d_G(v_j)) \\ \leq \left(\frac{\sqrt{\Delta^2 + \delta^2}}{\Delta + \delta} + \frac{1}{\sqrt{2}} \right) \sum_{v_i v_j \in E(G)} \sqrt{d_G(v_i)^2 + d_G(v_j)^2}, \end{aligned} \tag{6}$$

that is,

$$m \delta + \frac{\sqrt{\Delta^2 + \delta^2}}{\sqrt{2}(\Delta + \delta)} M_1(G) \leq \left(\frac{\sqrt{\Delta^2 + \delta^2}}{\Delta + \delta} + \frac{1}{\sqrt{2}} \right) SO(G),$$

that is,

$$\left(\sqrt{2(\Delta^2 + \delta^2)} + \Delta + \delta \right) SO(G) \geq \sqrt{\Delta^2 + \delta^2} M_1(G) + \sqrt{2}(\Delta + \delta) \delta m,$$

by (5). The first part of the proof is done.

Suppose that equality holds in (3). Then all the above inequalities must be equalities. By Lemma 2, from the equality in (6), we obtain

$$\frac{1}{\sqrt{2}} = \frac{\sqrt{\Delta^2 + \delta^2}}{\Delta + \delta}, \quad \text{that is, } (\Delta - \delta)^2 = 0, \quad \text{that is, } \Delta = \delta.$$

Moreover, from the equality in (5), we obtain that G is a regular graph.

Conversely, let G be an r -regular graph. Then we have

$$\left(\sqrt{2(\Delta^2 + \delta^2)} + \Delta + \delta\right) SO(G) = 2\sqrt{2} n\Delta^3 = \sqrt{\Delta^2 + \delta^2} M_1(G) + \sqrt{2} (\Delta + \delta) \delta m,$$

□

We now obtain another relation between Sombor index $SO(G)$ and the second Zagreb index $M_2(G)$ of graph G .

Theorem 8. Let G be a graph over n vertices. Suppose it has maximum degree Δ and minimum degree $\delta > 0$. We obtain

$$\sqrt{2}(\Delta + \delta) SO(G) \geq 2 M_2(G) + n \Delta \delta^2, \tag{7}$$

where $M_2(G)$ is the second Zagreb index of graph G . Moreover, the equality holds in (7) if and only if G is a regular graph.

Proof. Since $0 < \delta \leq d_G(v_i) \leq \Delta$ ($v_i \in V(G)$), for any edge $v_i v_j \in E(G)$, we obtain

$$\frac{\sqrt{2}}{\Delta} \leq \frac{\sqrt{d_G(v_i)^2 + d_G(v_j)^2}}{d_G(v_i) d_G(v_j)} = \sqrt{\frac{1}{d_G(v_i)^2} + \frac{1}{d_G(v_j)^2}} \leq \frac{\sqrt{2}}{\delta}.$$

Let $\mu_G(v_i)$ be the average degree of the adjacent vertices of vertex v_i in G . Then

$$\mu_G(v_i) = \frac{\sum_{v_j: v_i v_j \in E(G)} d_G(v_j)}{d_G(v_i)}, \text{ that is, } d_G(v_i) \mu_G(v_i) = \sum_{v_j: v_i v_j \in E(G)} d_G(v_j).$$

From the definition of the average degree of vertex v_i , we have $\delta \leq \mu_G(v_i) \leq \Delta$. Now,

$$\begin{aligned} \sum_{v_i v_j \in E(G)} \left(\frac{d_G(v_i)}{d_G(v_j)} + \frac{d_G(v_j)}{d_G(v_i)} \right) &= \sum_{v_i \in V(G)} \sum_{v_j: v_i v_j \in E(G)} \frac{d_G(v_j)}{d_G(v_i)} \\ &= \sum_{v_i \in V(G)} \mu_G(v_i) \\ &\geq n \delta \end{aligned} \tag{8}$$

with equality holding if and only if $d_G(v_i) = \delta$ for any $v_i \in V(G)$.

Setting $s = \frac{\sqrt{2}}{\Delta}$, $t = \frac{\sqrt{2}}{\delta}$, $a_i \rightarrow \sqrt{d_G(v_i) d_G(v_j)}$ and $b_i \rightarrow \sqrt{\frac{d_G(v_i)^2 + d_G(v_j)^2}{d_G(v_i) d_G(v_j)}}$ in Lemma 2, we obtain

$$\begin{aligned} \sum_{v_i v_j \in E(G)} \frac{d_G(v_i)^2 + d_G(v_j)^2}{d_G(v_i) d_G(v_j)} + \frac{2}{\Delta \delta} \sum_{v_i v_j \in E(G)} d_G(v_i) d_G(v_j) \\ \geq \frac{\sqrt{2}(\Delta + \delta)}{\Delta \delta} \sum_{v_i v_j \in E(G)} \sqrt{d_G(v_i)^2 + d_G(v_j)^2}, \end{aligned}$$

that is,

$$\sum_{v_i v_j \in E(G)} \left(\frac{d_G(v_i)}{d_G(v_j)} + \frac{d_G(v_j)}{d_G(v_i)} \right) + \frac{2}{\Delta \delta} M_2(G) \leq \frac{\sqrt{2}(\Delta + \delta)}{\Delta \delta} SO(G),$$

that is,

$$\sqrt{2}(\Delta + \delta) SO(G) \geq 2 M_2(G) + n \Delta \delta^2,$$

by (8). The first part of the proof is done.

Similarly, the proof of the Theorem 7, we conclude that the equality holds in (7) if and only if G is a regular graph. \square

The following inequality is due to Radon [39].

Lemma 3. (Radon’s inequality) *If $a_k, x_k > 0, p > 0, k \in \{1, 2, \dots, r\}$, then the following inequality holds:*

$$\sum_{k=1}^r \frac{x_k^{p+1}}{a_k^p} \geq \frac{\left(\sum_{k=1}^r x_k \right)^{p+1}}{\left(\sum_{k=1}^r a_k \right)^p}$$

with equality holding $\frac{x_1}{a_1} = \frac{x_2}{a_2} = \dots = \frac{x_r}{a_r}$.

We now present a relation between Sombor index $SO(G)$ and the second Zagreb index $M_2(G)$.

Theorem 9. *Let G be a graph over n vertices. Suppose G has maximum degree Δ and minimum degree $\delta > 0$. We have*

$$SO(G)^2 \leq \left(\frac{\Delta}{\delta} + \frac{\delta}{\Delta} \right) m M_2(G), \tag{9}$$

with equality if and only if G is a bipartite semiregular graph or G is a regular graph.

Proof. For any edge $v_i v_j \in E(G)$ ($d_G(v_i) \geq d_G(v_j)$) and $\delta > 0$, by (4), we obtain

$$\begin{aligned} \left(\frac{d_G(v_i)}{d_G(v_j)} + \frac{d_G(v_j)}{d_G(v_i)} \right)^2 &= \left(\frac{d_G(v_i)}{d_G(v_j)} - \frac{d_G(v_j)}{d_G(v_i)} \right)^2 + 4 \\ &\leq \left(\frac{\Delta}{\delta} - \frac{\delta}{\Delta} \right)^2 + 4 = \left(\frac{\Delta}{\delta} + \frac{\delta}{\Delta} \right)^2, \end{aligned}$$

that is,

$$\frac{d_G(v_i)}{d_G(v_j)} + \frac{d_G(v_j)}{d_G(v_i)} \leq \frac{\Delta}{\delta} + \frac{\delta}{\Delta} \tag{10}$$

with equality if and only if $d_G(v_i) = \Delta, d_G(v_j) = \delta$, or $d_G(v_i) = \delta, d_G(v_j) = \Delta$.

Setting $p = 1$, $x_k \rightarrow \sqrt{d_G(v_i)^2 + d_G(v_j)^2}$ and $a_k \rightarrow d_G(v_i) d_G(v_j)$ in Lemma 3 and using the above result, we obtain

$$\frac{SO(G)^2}{M_2(G)} = \frac{\left(\sum_{v_i, v_j \in E(G)} \sqrt{d_G(v_i)^2 + d_G(v_j)^2} \right)^2}{\sum_{v_i, v_j \in E(G)} d_G(v_i) d_G(v_j)} \leq \sum_{v_i, v_j \in E(G)} \frac{d_G(v_i)^2 + d_G(v_j)^2}{d_G(v_i) d_G(v_j)} \quad (11)$$

$$= \sum_{v_i, v_j \in E(G)} \left(\frac{d_G(v_i)}{d_G(v_j)} + \frac{d_G(v_j)}{d_G(v_i)} \right) \leq \left(\frac{\Delta}{\delta} + \frac{\delta}{\Delta} \right) m. \quad (12)$$

The first part of the proof is done.

Suppose that equality holds in (9). Then all the above inequalities must be equalities. By Lemma 3, from the equality in (11), for any edges $v_i v_j, v_k v_\ell \in E(G)$, we obtain

$$\frac{\sqrt{d_G(v_i)^2 + d_G(v_j)^2}}{d_G(v_i) d_G(v_j)} = \frac{\sqrt{d_G(v_k)^2 + d_G(v_\ell)^2}}{d_G(v_k) d_G(v_\ell)},$$

that is,

$$\frac{1}{d_G(v_i)^2} + \frac{1}{d_G(v_j)^2} = \frac{1}{d_G(v_k)^2} + \frac{1}{d_G(v_\ell)^2}.$$

From the equality in (12), we obtain $d_G(v_i) = \Delta, d_G(v_j) = \delta$ for any edge $v_i v_j \in E(G)$, by (10). Using the above results, we conclude that G is a (Δ, δ) -semiregular bipartite graph (when G is bipartite) or G is a regular graph (when G is non-bipartite).

Conversely, let G be an r -regular graph. Then

$$SO(G)^2 = 2m^2 r^2 = \left(\frac{\Delta}{\delta} + \frac{\delta}{\Delta} \right) m M_2(G).$$

Let G be a (Δ, δ) -semiregular bipartite graph. Then

$$SO(G)^2 = m^2(\Delta^2 + \delta^2) = \left(\frac{\Delta}{\delta} + \frac{\delta}{\Delta} \right) m M_2(G).$$

The proof is then complete. \square

4. Conclusions

Topological indices are graph invariants and are used for quantitative structure - activity relationship (QSAR) and quantitative structure - property relationship (QSPR) studies. Many topological indices have been defined in the literature and several of them have found applications as a means to model physical, chemical, pharmaceutical, and other properties of molecules. Gutman introduced the SO index as a new topological index in mathematical chemistry. In this paper, we presented some upper and lower bounds on the SO index and characterized extremal graphs. Moreover, we obtained some relations between the Sombor index and the (first & second) Zagreb indices of graphs. The minimal and the maximal Sombor index (SO), in the case of unicyclic graphs and bicyclic graphs, remains an open problem. Motivation to better understand the Sombor index has been mentioned in the

literature [28]. Finding the chemical applications of this Sombor index is an attractive task for the near future.

Author Contributions: Investigation, K.C.D., A.S.Ç., I.N.C. and Y.S.; writing—original draft preparation, K.C.D., A.S.Ç., I.N.C. and Y.S.; writing—review and editing, K.C.D., A.S.Ç., I.N.C. and Y.S.; funding acquisition, Y.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Northumbria University under No. 201920A1001.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Gutman, I.; Trinajstić, N. Graph theory and molecular orbitals. Total π -electron energy of alternant hydrocarbons. *Chem. Phys. Lett.* **1972**, *17*, 535–538.
2. Gutman, I.; Ruščić, B.; Trinajstić, N.; Wilcox, C.F. Graph theory and molecular orbitals. XII. Acyclic Polyenes. *J. Chem. Phys.* **1975**, *62*, 3399–3405.
3. Das, K.C. Maximizing the sum of the squares of the degrees of a graph. *Discrete Math.* **2004**, *285*, 57–66.
4. Das, K.C. Sharp bounds for the sum of the squares of the degrees of a graph. *Kragujevac J. Math.* **2003**, *25*, 31–49.
5. Das, K.C. On comparing Zagreb indices of graphs. *MATCH Commun. Math. Comput. Chem.* **2010**, *63*, 433–440.
6. Das, K.C.; Gutman, I. Some properties of the Second Zagreb Index. *MATCH Commun. Math. Comput. Chem.* **2004**, *52*, 103–112.
7. Das, K.C.; Gutman, I.; Horoldagva, B. Comparison between Zagreb indices and Zagreb coincides. *MATCH Commun. Math. Comput. Chem.* **2012**, *68*, 189–198.
8. Das, K.C.; Gutman, I.; Zhou, B. New upper bounds on Zagreb indices. *J. Math. Chem.* **2009**, *46*, 514–521.
9. Das, K.C.; Xu, K.; Gutman, I. On Zagreb and Harary indices. *MATCH Commun. Math. Comput. Chem.* **2013**, *70*, 301–314.
10. Gutman, I.; Das, K.C. The first Zagreb index 30 years after. *MATCH Commun. Math. Comput. Chem.* **2004**, *50*, 83–92.
11. Xu, K.; Das, K.C.; Balachandran, S. Maximizing the Zagreb indices of (n, m) -graphs. *MATCH Commun. Math. Comput. Chem.* **2014**, *72*, 641–654.
12. Xu, K. The Zagreb indices of graphs with a given clique number. *Appl. Math. Lett.* **2011**, *24*, 1026–1030.
13. Yan, Z.; Liu, H.; Liu, H. Sharp bounds for the second Zagreb index of unicyclic graphs. *J. Math. Chem.* **2007**, *42*, 565–574.
14. Zhou, B. Upper bounds for the Zagreb indices and the spectral radius of series-parallel graphs. *Int. J. Quantum Chem.* **2007**, *107*, 875–878.
15. Zhou, B.; Gutman, I. Further properties of Zagreb indices. *MATCH Commun. Math. Comput. Chem.* **2005**, *54*, 233–239.
16. Ali, A.; Das, K.C.; Akhter, S. On the Extremal Graphs for Second Zagreb Index with Fixed Number of Vertices and Cyclomatic Number, Miskolc Mathematical Notes. in press. Available online: https://www.researchgate.net/publication/334881908_On_the_Extremal_Graphs_for_Second_Zagreb_Index_with_Fixed_Number_of_Vertices_and_Cyclomatic_Number(accessed on 19 December 2020)
17. An, M.; Das, K.C. First Zagreb index, k -connectivity, beta-deficiency and k -hamiltonicity of graphs. *MATCH Commun. Math. Comput. Chem.* **2018**, *80*, 141–151.
18. Borovičanić, B.; Das, K.C.; Furtula, B.; Gutman, I. Zagreb indices: Bounds and extremal graphs. *MATCH Commun. Math. Comput. Chem.* **2017**, *78*, 17–100.
19. Buyantogtokh, L.; Horoldagva, B.; Das, K.C. On reduced second Zagreb index. *J. Combin. Optim.* **2020**, *39*, 776–791.
20. Das, K.C.; Ali, A. On a conjecture about the second Zagreb index. *Discrete Math. Lett.* **2019**, *2*, 38–43.
21. Horoldagva, B.; Das, K.C. On Zagreb indices of graphs. *MATCH Commun. Math. Comput. Chem.* **2021**, *85*, 295–301.
22. Horoldagva, B.; Das, K.C.; Selenge, T.-A. Complete characterization of graphs for direct comparing Zagreb indices. *Discrete Appl. Math.* **2016**, *215*, 146–154.
23. Shang, Y. On the number of spanning trees, the Laplacian eigenvalues, and the Laplacian Estrada index of subdivided-line graphs. *Open Math.* **2016**, *14*, 641–648.
24. Shang, Y. Lower bounds for Gaussian Estrada index of graphs. *Symmetry* **2018**, *10*, 325.
25. Xu, K.; Gao, F.; Das, K.C.; Trinajstić, N. A formula with its applications on the difference of Zagreb indices of graphs. *J. Math. Chem.* **2019**, *57*, 1618–1626.
26. Todeschini, R.; Consonni, V. *Handbook of Molecular Descriptors*; Wiley-VCH: Weinheim, Germany, 2000.
27. Balaban, A.T.; Motoc, I.; Bonchev, D.; Mekenyan, O. Topological indices for structure-activity correlations. *Top. Curr. Chem.* **1983**, *114*, 21–55.
28. Gutman, I. Geometric approach to degree-based topological indices: Sombor indices. *MATCH Commun. Math. Comput. Chem.* **2021**, *86*, 11–16.
29. Bondy, J.A.; Murty, U.S.R. *Graph Theory with Applications*; MacMillan: New York, NY, USA, 1976.
30. Shang, Y. Lower bounds for the Estrada index using mixing time and Laplacian spectrum. *Rocky Mt. J. Math.* **2013**, *43*, 2009–2016.

31. Shang, Y. Estrada and L -Estrada indices of edge-independent random graphs. *Symmetry* **2015**, *7*, 1455–1462.
32. Das, K.C. Comparison between Zagreb eccentricity indices and the eccentric connectivity index, the second geometric-arithmetic index and the Graovac-Ghorbani index. *Croat. Chem. Acta* **2016**, *89*, 505–510.
33. Das, K.C.; Dehmer, M. Comparison between the zeroth-order Randić index and the sum-connectivity index. *Appl. Math. Comput.* **2016**, *274*, 585–589.
34. Das, K.C.; Mohammed, M.A.; Gutman, I.; Atan, K.A. Comparison between Atom-Bond Connectivity Indices of Graphs. *MATCH Commun. Math. Comput. Chem.* **2016**, *76*, 159–170.
35. Das, K.C.; Su, G.; Xiong, L. Relation between Degree Distance and Gutman Index of Graphs. *MATCH Commun. Math. Comput. Chem.* **2016**, *76*, 221–232.
36. Xu, K.; Alizadeh, Y.; Das, K.C. On two eccentricity-based topological indices of graphs. *Discrete Appl. Math.* **2017**, *233*, 240–251.
37. Xu, K.; Das, K.C.; Liu, H. Some extremal results on the connective eccentricity index of graphs. *J. Math. Anal. Appl.* **2016**, *433*, 803–817.
38. Diaz, J.B.; Metcalf, F.T. Stronger forms of a class of inequalities of G. Pólya-G. Szegő and L. V. Kantorovich. *Bull. Am. Math. Soc.* **1963**, *69*, 415–418.
39. Radon, J. über Die Absolut Additiven Mengenfunktionen. *Wien. Sitzungsber* **1913**, *122*, 1295–1438.