

Northumbria Research Link

Citation: Shang, Yilun (2023) On connectivity and robustness of random graphs with inhomogeneity. *Journal of Applied Probability*, 60 (1). pp. 284-294. ISSN 0021-9002

Published by: Cambridge University Press

URL: <https://doi.org/10.1017/jpr.2022.32> <<https://doi.org/10.1017/jpr.2022.32>>

This version was downloaded from Northumbria Research Link:
<https://nrl.northumbria.ac.uk/id/eprint/50675/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

ON CONNECTIVITY AND ROBUSTNESS OF RANDOM GRAPHS WITH INHOMOGENEITY

YILUN SHANG,* *Northumbria University*

Abstract

The study of threshold functions has a long history in random graph theory. It is known that the thresholds for minimum degree k , k -connectivity, as well as k -robustness coincide for a binomial random graph. In this paper, we consider an inhomogeneous random graph model, which is obtained by including each possible edge independently with an individual probability. Based on an intuitive concept of neighborhood density, we show two sufficient conditions, respectively, guaranteeing k -connectivity and k -robustness, which are asymptotically equivalent. Our framework sheds some light on extending uniform threshold values in homogeneous random graphs to threshold landscapes in inhomogeneous random graphs.

Keywords: random graph; connectivity; robustness; threshold

2020 Mathematics Subject Classification: Primary 60C05

Secondary 05C80; 05C40

1. Introduction

Classical binomial random graph theory founded by Gilbert [1] and Erdős and Rényi [2, 3] in the late 1950s considers the random graph model $G(n, p_n)$ of all graphs over the vertex set $V = \{1, 2, \dots, n\}$, in which each edge $e_{ij} \in E$ appears independently with probability p_n . Here, $E = \{e_{ij} = e_{ji} : 1 \leq i \neq j \leq n\}$ consists of all edges in the complete graph K_n over V . The probability of a random graph in $G(n, p_n)$ holding a graph property is typically understood in the large graph limit as $n \rightarrow \infty$. One of the most important results on random graphs, shown by Erdős and Rényi [3], is the

* Postal address: Department of Computer and Information Sciences, Northumbria University, NE1 8ST, Newcastle upon Tyne, UK

* Email address: yilun.shang@northumbria.ac.uk

following sharp threshold for the k -connectivity. Recall that a graph G is said to be k -connected if it remains connected when any set of at most $k - 1$ vertices is deleted.

Theorem 1. [3] *Let $p_n = \frac{1}{n}(\ln n + (k - 1) \ln \ln n + \omega_n)$ for an integer $k \geq 1$. Then*

$$\lim_{n \rightarrow \infty} \mathbb{P}(G(n, p_n) \text{ is } k\text{-connected}) = \lim_{n \rightarrow \infty} \mathbb{P}(G(n, p_n) \text{ has a minimum degree no less than } k) = \begin{cases} 0, & \text{if } \omega_n \rightarrow -\infty, \\ 1, & \text{if } \omega_n \rightarrow \infty. \end{cases}$$

The above type of zero-one law is known to be universal for all monotone properties in random graphs [4]. The theorem indicates that although having minimum degree at least k is weaker than k -connectedness, both properties share the same asymptotic threshold function (In this case, $\frac{1}{n}(\ln n + (k - 1) \ln \ln n)$).

Connectivity is a fundamental property of graph theory and is essential to many distributed computation problems over large-scale complex networks. For example, synchronization of oscillators or mobile agents is not possible without a connected underlying communication network. To cope with undesirable disturbances such as node faults and malicious attacks in realistic networked systems, a novel graph concept known as k -robustness was recently introduced by LeBlanc et al. [5] together with a class of distributed resilient consensus protocols called Weighted-Mean Subsequence Reduced (W-MSR) algorithms. In these algorithms, a node (or vertex) calculates its value in each iteration based on the current neighbors' states and cuts the links with some neighbors that have extreme values. The notion of k -robustness (see Section 2 for the definition) of the communication network has been proposed as a sufficient condition to guarantee the global consensus of the network against malicious neighbors. This notion has been found central in a number of related network control protocols; see e.g. [6, 7, 8]. In this context, a sufficiently connected but inadequately robust network is not able to deliver the consensus result. As will be seen below, k -robustness is a stronger property than k -connectivity. Interestingly, Zhang et al. [9] showed that Theorem 1 holds verbatim for k -robustness.

Theorem 2. [9] *Let $p_n = \frac{1}{n}(\ln n + (k - 1) \ln \ln n + \omega_n)$ for an integer $k \geq 1$. Then*

$$\lim_{n \rightarrow \infty} \mathbb{P}(G(n, p_n) \text{ is } k\text{-robust}) = \begin{cases} 0, & \text{if } \omega_n \rightarrow -\infty, \\ 1, & \text{if } \omega_n \rightarrow \infty. \end{cases}$$

In other words, a random graph in $G(n, p_n)$ becomes k -robust as soon as it becomes k -connected (and as soon as its last vertex of degree $k - 1$ vanishes).

The aforementioned threshold $\frac{1}{n}(\ln n + (k - 1) \ln \ln n)$ is obviously precise and plain (note that Theorem 1 is a weaker statement than the result in [3]). However, would there be a possibility to zoom in and examine details of the threshold landscape (across all edges)? A natural way is to introduce unequal edge probabilities. To this end, we in this paper consider an inhomogeneous random graph with a list of edge probabilities $\mathbf{p}_n = \{p_n(e_{ij})\}_{1 \leq i < j \leq n}$ by including each edge $e_{ij} = e_{ji}$ of K_n independently with probability $p_n(e_{ij})$. We denote this inhomogeneous random graph model by $G(n, \mathbf{p}_n)$.

As an effort to diversify and extend previous results regarding binomial random graphs, it is not surprising that $G(n, \mathbf{p}_n)$ model has been studied by a few researchers across a relatively long time period under different names, e.g., generalized binomial random graphs [10, 12], anisotropic random graphs [11, 15], and edge-independent random graphs [13, 14]. We mention that Chapter 9 of the monograph [15] presents an updated survey of connectivity related results for this and other inhomogeneous random graphs. In the seminal work [16], the authors systematically analyze a very general class of inhomogeneous random graphs and networks that have been studied during the recent decade.

In this paper, we attempt to generalize the threshold result presented in Theorem 1 for both k -connectivity and k -robustness by considering the $G(n, \mathbf{p}_n)$ model and shed some light on the shape of the threshold landscape. We show two sufficient conditions for k -connectivity and k -robustness, which are asymptotically equivalent as $n \rightarrow \infty$. It is hoped that the approach developed in the present work could facilitate relevant research in this direction on other topics in random graphs. The main results of this paper and their implications are discussed in Section 2 and proofs are given in Section 3.

2. Statements of main results and discussions

By convention, we will use the standard Landau asymptotic notations such as $O(\cdot)$, $o(\cdot)$, $\Theta(\cdot)$, \sim etc. throughout the paper; see e.g. [17]. For a simple, undirected graph $G = (V_G, E_G)$ with the vertex set $V_G = \{1, 2, \dots, n\}$ and the edge set E_G , let

$N_G(i) = \{j \in V_G : e_{ij} \in E_G\}$ be the set of neighbors (i.e., open neighborhood) of vertex $i \in V_G$. The degree of vertex i is denoted by $d_G(i) = |N_G(i)|$. Let $\bar{N}_G(i) = \{i\} \cup N_G(i)$ be the closed neighborhood of i . We also define the neighborhood of a set $S \subseteq V_G$ as $N_G(S) = \{j \in V_G : e_{ij} \in E_G \text{ for some } i \in S\}$. As mentioned in the introduction, the following notion of graph robustness was put forward in [5] to analyze the convergence of W-MSR algorithms in resilient control in the presence of adversaries.

Definition 1. [5] (a) For an integer $k \geq 1$ and a graph $G = (V_G, E_G)$, a subset $S \subseteq V_G$ is called k -reachable if there is some vertex $i \in S$ satisfying $|N_G(i) \setminus S| \geq k$. (b) A graph G is called k -robust if for any two nonempty, disjoint sets in V_G , at least one of them is k -reachable.

From Definition 1, it is not difficult to see that k -robustness is a stronger property compared to k -connectedness in general. In fact, if a graph G is k -connected, for any two nonempty disjoint sets in V_G at least one of them has k neighbors (collectively) outside of the set itself. This does not guarantee k -reachability of either of these sets for $k \geq 2$. Generally, we have the following relationship between the three properties: $\{k\text{-robustness}\} \subseteq \{k\text{-connectedness}\} \subseteq \{\text{minimum degree } k\}$, and for the binomial random graph $G(n, p_n)$ they share the same threshold function $p_n = \frac{1}{n}(\ln n + (k - 1) \ln \ln n)$.

To accommodate the inhomogeneous random graph setting, we consider the neighborhood density for a vertex $i \in V$ with respect to a set $S \subseteq V$. For $i \in V \setminus S$, define $\rho_n(i, S) = |S|^{-1} \sum_{j \in S} p_n(e_{ij})$. This quantifies the expected fraction of neighbors of i inside S . Let $p_n = \frac{1}{n}(\ln n + (k - 1) \ln \ln n + \omega_n)$, where $k \geq 1$ is an integer and $\omega_n \rightarrow \infty$ as $n \rightarrow \infty$. For a constant $\beta \geq 1$, define

$$\rho_n^\beta(i, S) = \frac{1}{|S|} \sum_{j \in S} \min\{p_n(e_{ij}), \beta p_n\}. \quad (1)$$

A sufficient condition ensuring k -connectivity and minimum degree k is formalized in the following.

Theorem 3. *Suppose that there is some constant $\beta \geq 1$ and an integer $k \geq 1$ satisfies*

$$\min_{1 \leq i \leq n} \min_{\substack{S: i \notin S \\ |S| \geq \lfloor n/2 \rfloor}} \rho_n^\beta(i, S) \geq p_n \quad (2)$$

for all large n . Then

$$\begin{aligned} & \lim_{n \rightarrow \infty} \mathbb{P}(G(n, \mathbf{p}_n) \text{ is } k\text{-connected}) \\ &= \lim_{n \rightarrow \infty} \mathbb{P}(G(n, \mathbf{p}_n) \text{ has a minimum degree no less than } k) = 1. \end{aligned}$$

Theorem 3 is a generalization of the “one” part of the zero-one law in Theorem 1, which can be seen by taking $p_n(e_{ij}) \equiv p_n$ for all distinct $i, j \in V$ in Theorem 3. Roughly speaking, Theorem 3 says if the edge probabilities are large enough, then the random graph $G(n, \mathbf{p}_n)$ is sufficiently connected with high probability. The quantity $\rho_n^\beta(i, S)$ can be viewed as a capped neighborhood density with each potential edge contributing no more than βp_n . If (2) holds for a constant β , then it holds for any larger β . Moreover, if $\max_{i, j \in V} p_n(e_{ij}) \leq \beta p_n$, then $\rho_n^\beta(i, S) = \rho_n(i, S)$ and (2) reduces to

$$\min_{1 \leq i \leq n} \min_{\substack{S: i \notin S \\ |S| \geq \lfloor n/2 \rfloor}} \rho_n(i, S) \geq p_n \quad (3)$$

concerning the neighborhood density. Since k -connectivity and minimum degree no less than k are monotonic properties, without loss of generality we will only present the proofs in Section 3 under this maximum probability condition.

To appreciate the above presented result, we consider an example of non-trivial edge probabilities satisfying (2).

Example 1. To build an inhomogeneous random graph $G(n, \mathbf{p}_n)$, we consider the following assignment probabilities. For $1 \leq i < j \leq \lfloor n^\alpha \rfloor$ for some $\alpha \in (0, 1)$, let $p_n(e_{ij}) = 0$. For all other $1 \leq i < j \leq n$, let $p_n(e_{ij}) = \frac{1}{n}(\ln n + (k-1) \ln \ln n + \omega_n)$ with $k \geq 1$ and $\omega_n \rightarrow \infty$. Let $\beta = 1$.

Notice that for any $S \subseteq V$ with $|S| \geq \lfloor n/2 \rfloor$ and $i \in V \setminus S$,

$$\begin{aligned} \rho_n^\beta(i, S) &\geq \frac{1}{|S|} \left(n^\alpha \cdot 0 + (|S| - n^\alpha) \cdot \frac{\ln n + (k-1) \ln \ln n + \omega_n}{n} \right) \\ &\geq \frac{\ln n + (k-1) \ln \ln n + \omega'_n}{n}, \end{aligned}$$

where $\omega'_n \rightarrow \infty$ as $n \rightarrow \infty$. It is easy to see that (2) is satisfied and hence by Theorem 3 $G(n, \mathbf{p}_n)$ is k -connected a.a.s. (i.e., asymptotically almost surely [17]). Loosely speaking, this example indicates that with a sub-linear order portion of the graph being arbitrarily sparse and the rest of the graph k -connected, the entire graph can still be k -connected a.a.s.

The next example looks into a classical result regarding connectivity of $G(n, \mathbf{p}_n)$ by Alon, which is sharp up to a multiplicative factor c .

Example 2. It is known that [12] for every constant $b > 0$ there exists a constant $c > 0$ so that if, for any $\emptyset \neq S \subset V$,

$$\sum_{j \in S, i \in V \setminus S} p_n(e_{ij}) \geq c \ln n, \quad (4)$$

then

$$\mathbb{P}(G(n, \mathbf{p}_n) \text{ is connected}) \geq 1 - n^{-b}.$$

By dividing both sides of the condition (4) by $|S|$, we can rewrite it as

$$\sum_{i \in V \setminus S} \rho_n(i, S) \geq \frac{c \ln n}{|S|}. \quad (5)$$

Write $\rho_n(i) = (n-1)^{-1} \sum_{j \in V \setminus \{i\}} p_n(e_{ij})$ for the neighborhood density of a vertex $i \in V$ in the entire graph. Firstly, taking $S = V \setminus \{i\}$, the condition (3) becomes $\min_{i \in V} \rho_n(i) \geq \frac{1}{n}(\ln n + \omega_n)$ when $k = 1$, while (5) reduces to $\min_{i \in V} \rho_n(i) \geq \frac{c}{n} \ln n$. As c is unknown, it is not difficult to see that these two conditions are incomparable in general. Secondly, the form of condition (5) takes a sum over all $i \notin S$, which mitigates the risk of uneven contribution of edge probabilities. Our condition (2) considers a single vertex i with minimum neighborhood density. Therefore, we have to consider a cap on the contribution to neighborhood density. Finally, for a given random graph $G(n, \mathbf{p}_n)$, the condition (4) is not always easy to check compared to (2) due to the undetermined constant c .

Next, we present our result for robustness.

Theorem 4. *Suppose that there is some constant $\beta \geq 1$ and an integer $k \geq 1$ satisfies the condition (2) for all large n . Then*

$$\lim_{n \rightarrow \infty} \mathbb{P}(G(n, \mathbf{p}_n) \text{ is } k\text{-robust}) = 1.$$

Several remarks are in order. Firstly, Theorem 4 can be viewed as a generalization of the “one” part of the zero-one law Theorem 2 for robustness of binomial random graphs, which can be recovered by taking $p_n(e_{ij}) \equiv p_n$ for all distinct $i, j \in V$ in Theorem 4.

Secondly, the same condition is adopted here as in Theorem 3 and analogous comments underneath it are applicable here. That said, k -robustness may be essentially stronger than k -connectedness in the current setting as we are not able to match a “zero” part statement. As an example (which appeared in [9]) of a heterogeneously connected graph, we consider the union graph G of a complete bipartite graph $K_{n/2, n/2}$ and a perfect matching between the two partite sets. G has connectivity $n/2$ but is just 1-robust, suggesting a gap might exist in the thresholds for connectivity and robustness in $G(n, \mathbf{p}_n)$ model.

Finally, it would be useful to perform some computational experiments to vouch for the above conjecture. Unfortunately, as shown in [9, 18] determining robustness is significantly harder than checking connectivity, namely, NP-hard versus P in time complexity. The most current robustness checking algorithm can only effectively handle graphs of order about $n = 30$.

Example 1. (revisited) For the model $G(n, \mathbf{p}_n)$ set up in Example 1, we can argue in the same way and show that $G(n, \mathbf{p}_n)$ is not only k -connected but k -robust a.a.s. by Theorem 4.

3. Proofs of Theorem 3 and Theorem 4

In this section, we present the proofs of our main results. As commented above, we will assume $\max_{i,j \in V} p_n(e_{ij}) \leq \beta p_n$ for some constant $\beta \geq 1$ and that the condition (3) holds.

Proof of Theorem 3. Here, we will actually rely on the following inequality instead of (3):

$$\min_{1 \leq i \leq n} \min_{\substack{S: i \notin S \\ |S| \geq \lfloor \frac{n-k+1}{2} \rfloor}} \rho_n(i, S) \geq p_n. \quad (6)$$

The minimum in condition (6) has the same asymptotic behavior as that in condition (3). In fact, although $\lfloor (n-k+1)/2 \rfloor < \lfloor n/2 \rfloor$ for any $k \geq 1$, the difference between the two terms is $O(1)$. Hence, it can only make a difference of order $O(n^{-1})$ to the right-hand side of (3) or (6).

For an integer $d \geq 0$, let X_d be the random variable counting the number of vertices in $G(n, \mathbf{p}_n)$ having degree d . Write $N_n(i) := N_{G(n, \mathbf{p}_n)}(i)$ and $\bar{N}_n(i) := \bar{N}_{G(n, \mathbf{p}_n)}(i)$,

respectively, for the open and closed neighborhood of vertex i in G_{n, \mathbf{p}_n} . Accordingly, we use $N_n^*(i)$ and $\bar{N}_n^*(i)$ to represent a potential instance of an open and closed neighborhood of i , respectively. By the construction of our inhomogeneous random graph model, we have

$$\mathbb{E}X_d = \sum_{i=1}^n \sum_{N_n^*(i): |N_n^*(i)|=d} \prod_{j \in N_n^*(i)} p_n(e_{ij}) \prod_{j \in V \setminus \bar{N}_n^*(i)} (1 - p_n(e_{ij})).$$

Capitalizing on the neighborhood density condition (6) and the estimate for binomial coefficient $\binom{n}{d} = (1 + o(1))n^d/d!$ for fixed d as $n \rightarrow \infty$, we obtain

$$\begin{aligned} \mathbb{E}X_d &\leq \sum_{i=1}^n \sum_{N_n^*(i): |N_n^*(i)|=d} \beta^d p_n^d e^{-|V \setminus \bar{N}_n^*(i)| \cdot \rho_n(i, V \setminus \bar{N}_n^*(i))} \\ &\leq n \binom{n-1}{d} \beta^d p_n^d e^{-(n-1-d)p_n} \\ &\leq (1 + o(1))n \cdot \frac{n^d}{d!} \beta^d \frac{(\ln n)^d}{n^d} \cdot \frac{e^{-\omega_n}}{n(\ln n)^{k-1}} \\ &= (1 + o(1)) \frac{\beta^d (\ln n)^d e^{-\omega_n}}{d! (\ln n)^{k-1}}. \end{aligned} \quad (7)$$

It follows from (7) that

$$\begin{aligned} \sum_{d=0}^{k-2} \mathbb{E}X_d &\leq (1 + o(1)) \frac{e^{-\omega_n}}{(\ln n)^{k-1}} \sum_{d=0}^{k-2} \beta^d \frac{(\ln n)^d}{d!} \\ &\leq (1 + o(1)) \frac{e^{-\omega_n}}{\ln n} (k-1) \beta^{k-2} \end{aligned} \quad (8)$$

and

$$\mathbb{E}X_{k-1} \leq (1 + o(1)) \frac{\beta^{k-1}}{(k-1)!} e^{-\omega_n}. \quad (9)$$

Since k and β are fixed constants and $\omega_n \rightarrow \infty$ as $n \rightarrow \infty$, by (8) and (9) we obtain $\sum_{d=0}^{k-2} \mathbb{E}X_d = o(1)$ and $\mathbb{E}X_{k-1} = o(1)$. Let d_n^{\min} represent the minimum degree of $G(n, \mathbf{p}_n)$. Then $\mathbb{P}(d_n^{\min} \geq k) \rightarrow 1$ as $n \rightarrow \infty$ by Markov's inequality.

Notice that

$$\begin{aligned} &\mathbb{P}(G(n, \mathbf{p}_n) \text{ is } k\text{-connected}) \\ &\geq \mathbb{P}(G(n, \mathbf{p}_n) \text{ is } k\text{-connected} \mid d_n^{\min} \geq k) \cdot \mathbb{P}(d_n^{\min} \geq k). \end{aligned}$$

To show Theorem 3, it suffices to prove

$$\lim_{n \rightarrow \infty} \mathbb{P}(G(n, \mathbf{p}_n) \text{ is } k\text{-connected} \mid d_n^{\min} \geq k) = 1. \quad (10)$$

Given two subsets $S_1, S_2 \subseteq V$ satisfying $0 \leq |S_1| \leq k-1$ and $k - |S_1| + 1 \leq |S_2| \leq \lceil \frac{1}{2}(n - |S_1|) \rceil$, consider an event

$$\mathcal{E}(S_1, S_2) = \{S_2 \text{ is a connected component of } G(n, \mathbf{p}_n) \setminus S_1\}.$$

Notice that if $G(n, \mathbf{p}_n) \setminus S_1$ is not connected, it must contain a component having order at most $\frac{1}{2}(n - |S_1|)$. In the following, we will bound the probability P_{12} that there exist two vertex sets S_1 and S_2 such that the event $\mathcal{E}(S_1, S_2)$ occurs. For this purpose, without loss of generality, we assume S_1 is minimal for the given S_2 . This implies that each vertex in S_1 is adjacent to at least one vertex in S_2 [Otherwise S_1 can be made smaller by removing a vertex without neighbors in S_2]. Moreover $N_{G(n, \mathbf{p}_n)}(S_2) = S_1$ since S_2 is a component. Let $|S_1| = s_1$ and $|S_2| = s_2$. Therefore, we have

$$\begin{aligned} P_{12} &\leq \sum_{s_1=0}^{k-1} \sum_{s_2=k-s_1+1}^{(n-s_1)/2} \sum_{\substack{S_1: |S_1|=s_1 \\ S_1 \subseteq V}} \sum_{\substack{S_2: |S_2|=s_2 \\ S_2 \subseteq V}} \sum_{\substack{T \subseteq S_2 \\ T \text{ is a spanning tree of } S_2}} \left(\prod_{e_{ij} \in T} p_n(e_{ij}) \right) \\ &\quad \cdot \left(\sum_{E_n^*(S_1, S_2): |E_n^*(S_1, S_2)|=s_1} \prod_{\substack{i: i \in S_1 \\ j \in S_2}} p_n(e_{ij}) \right) \cdot \prod_{e_{ij} \in E_n^*(V \setminus (S_1 \cup S_2), S_2)} (1 - p_n(e_{ij})) \\ &\leq \sum_{s_1=0}^{k-1} \sum_{s_2=k-s_1+1}^{(n-s_1)/2} \sum_{\substack{S_1: |S_1|=s_1 \\ S_1 \subseteq V}} \sum_{\substack{S_2: |S_2|=s_2 \\ S_2 \subseteq V}} \sum_{\substack{T \subseteq S_2 \\ T \text{ is a spanning tree of } S_2}} (\beta p_n)^{s_2-1} \\ &\quad \cdot \left(\sum_{E_n^*(S_1, S_2): |E_n^*(S_1, S_2)|=s_1} (\beta p_n)^{s_1} \right) \cdot e^{-\sum_{e_{ij} \in E_n^*(V \setminus (S_1 \cup S_2), S_2)} p_n(e_{ij})}, \quad (11) \end{aligned}$$

where $E_n(S_1, S_2) = \{e_{ij} \in G(n, \mathbf{p}_n) : i \in S_1, j \in S_2\}$ is the random edge set between two sets S_1 and S_2 , and $E_n^*(S_1, S_2)$ represents a potential instance accordingly. Using the neighborhood density assumption (2) and the relation $x - \lceil x/2 \rceil = \lfloor x/2 \rfloor$ for any nonnegative integer x , we have

$$e^{-\sum_{e_{ij} \in E_n(V \setminus (S_1 \cup S_2), S_2)} p_n(e_{ij})} = e^{-\sum_{j \in S_2} \sum_{i \in V \setminus (S_1 \cup S_2)} p_n(e_{ij})} \leq e^{-s_2(n-s_1-s_2)p_n}.$$

Since S_1 is minimal for the given S_2 , S_2 must be connected. There are at most $s_2^{s_2-2}$ different labelled spanning trees over S_2 by Cayley's formula. Hence, the right-hand

side of (11) is upper bounded by

$$\begin{aligned} & \sum_{s_1=0}^{k-1} \sum_{s_2=k-s_1+1}^{(n-s_1)/2} \binom{n}{s_1} \binom{n}{s_2} s_2^{s_2-2} \binom{s_1 s_2}{s_1} (\beta p_n)^{s_1+s_2-1} \cdot e^{-s_2(n-s_1-s_2)p_n} \\ & =: A_1 + A_2, \end{aligned} \tag{12}$$

where A_1 represents the value for $s_1 = 0$ and A_2 the sum for $1 \leq s_1 \leq k-1$.

We first estimate A_1 . Recall $k \geq 1$ and $\binom{n}{s_2} \leq (ne/s_2)^{s_2}$. We have

$$\begin{aligned} A_1 &= \sum_{s_2=k+1}^{n/2} \binom{n}{s_2} s_2^{s_2-2} (\beta p_n)^{s_2-1} e^{-s_2(n-s_2)p_n} \\ &\leq (\beta p_n)^{-1} \sum_{s_2=k+1}^{\ln n} \left(ne\beta p_n e^{-(n-s_2)p_n} \right)^{s_2} \\ &\quad + (\beta p_n)^{-1} \sum_{s_2=1+\ln n}^{n/2} \left(ne\beta p_n e^{-(n-s_2)p_n} \right)^{s_2} \\ &:= A_{11} + A_{12}. \end{aligned}$$

The term $ne\beta p_n e^{-(n-s_2)p_n}$ in the above sums is less than 1. Using the definition of p_n and $k \geq 1$, we obtain

$$\begin{aligned} A_{11} &\leq \frac{n}{\beta \ln n} \cdot (\ln n) \cdot \left(ne\beta p_n e^{-(n-\ln n)p_n} \right)^{k+1} \\ &= \frac{n}{\beta} \left((1+o(1))e\beta(\ln n)e^{-(1+o(1))\ln n} \right)^{k+1} = o(1) \end{aligned}$$

and

$$\begin{aligned} A_{12} &\leq \frac{n}{\beta \ln n} \sum_{s_2=1+\ln n}^{n/2} \left(ne\beta p_n e^{-(n/2)p_n} \right)^{s_2} \\ &\leq \frac{n}{\beta \ln n} \sum_{s_2=1+\ln n}^{\infty} \left((1+o(1))e\beta(\ln n + (k-1)\ln \ln n + \omega_n) \right. \\ &\quad \left. \cdot e^{-\frac{\ln n + (k-1)\ln \ln n + \omega_n}{2}} \right)^{s_2} \\ &= (1+o(1)) \frac{n}{\beta \ln n} \left((1+o(1))e\beta \cdot \frac{\ln n + (k-1)\ln \ln n + \omega_n}{e^{\frac{\ln n + (k-1)\ln \ln n + \omega_n}{2}}} \right)^{1+\ln n} \\ &= o(1). \end{aligned}$$

Therefore, $A_1 = A_{11} + A_{12} = o(1)$ as $n \rightarrow \infty$. Similarly, A_2 for $k \geq 2$ can be estimated

as follows.

$$\begin{aligned}
A_2 &= \sum_{s_1=1}^{k-1} \sum_{s_2=k-s_1+1}^{(n-s_1)/2} \binom{n}{s_1} \binom{n}{s_2} s_2^{s_2-2} \binom{s_1 s_2}{s_1} (\beta p_n)^{s_1+s_2-1} \cdot e^{-s_2(n-s_1-s_2)p_n} \\
&\leq (\beta p_n)^{-1} \sum_{s_1=1}^{k-1} \sum_{s_2=k-s_1+1}^{(n-s_1)/2} (ne^2 s_2 \beta p_n e^{s_2 p_n})^{s_1} \cdot (ne \beta p_n e^{-(n-s_2)p_n})^{s_2} \\
&= (\beta p_n)^{-1} \sum_{s_1=1}^{k-1} \sum_{s_2=k-s_1+1}^{\ln n} (ne^2 s_2 \beta p_n e^{s_2 p_n})^{s_1} \cdot (ne \beta p_n e^{-(n-s_2)p_n})^{s_2} \\
&\quad + (\beta p_n)^{-1} \sum_{s_1=1}^{k-1} \sum_{s_2=1+\ln n}^{(n-s_1)/2} (ne^2 s_2 \beta p_n e^{s_2 p_n})^{s_1} \cdot (ne \beta p_n e^{-(n-s_2)p_n})^{s_2} \\
&:= A_{21} + A_{22},
\end{aligned}$$

where the estimate $\binom{n}{s} \leq (ne/s)^s$ is used in the first inequality. Using the definition of p_n we write $p_n = \frac{1}{n}(\ln n + (1+o(1))(k-1)\ln \ln n) = (1+o(1))\frac{\ln n}{n}$. Therefore, we derive

$$\begin{aligned}
A_{21} &= O(1) \cdot \frac{n}{\ln n} \sum_{s_1=1}^{k-1} \sum_{s_2=k-s_1+1}^{\ln n} \left((1+o(1))\beta e^2 s_2 (\ln n) \cdot n^{\frac{s_2}{n}} (\ln n)^{\frac{(1+o(1))s_2(k-1)}{n}} \right)^{s_1} \\
&\quad \cdot \left((1+o(1))\beta e (\ln n) \cdot n^{\frac{s_2}{n}-1} (\ln n)^{(1+o(1))(k-1)(\frac{s_2}{n}-1)} \right)^{s_2} \\
&= O(1) \cdot \frac{n}{\ln n} \sum_{s_1=1}^{k-1} (O(\ln n))^{2s_1} \cdot \sum_{s_2=2}^{\ln n} \left(O\left(\frac{\ln n}{n}\right) \right)^{s_2} \\
&\leq nk (\ln n)^{2k} \cdot \left(\frac{\ln n}{n}\right)^2 = o(1),
\end{aligned}$$

where in the second equality above we note the limit $\lim_{n \rightarrow \infty} (n \ln n)^{\frac{c \ln n}{n}} = 1$ for any positive constant c and $s_2 \leq \ln n$. Likewise,

$$\begin{aligned}
A_{22} &= O(1) \cdot \frac{n}{\ln n} \sum_{s_1=1}^{k-1} \sum_{s_2=1+\ln n}^{(n-s_1)/2} \left((1+o(1))\beta e^2 s_2 (\ln n) \cdot n^{\frac{s_2}{n}} (\ln n)^{\frac{(1+o(1))s_2(k-1)}{n}} \right)^{s_1} \\
&\quad \cdot \left((1+o(1))\beta e (\ln n) \cdot n^{\frac{s_2}{n}-1} (\ln n)^{(1+o(1))(k-1)(\frac{s_2}{n}-1)} \right)^{s_2} \\
&= O(1) \cdot \frac{n}{\ln n} \sum_{s_1=1}^{k-1} \left(O\left(n^{\frac{3}{2}+o(1)}\right) \right)^{s_1} \cdot \sum_{s_2=1+\ln n}^{(n-s_1)/2} \left(O\left(n^{-\frac{1}{2}+o(1)}\right) \right)^{s_2} \\
&\leq \frac{nk}{\ln n} \left(n^{\frac{3}{2}+o(1)} \right)^k \cdot \left(n^{-\frac{1}{2}+o(1)} \right)^{1+\ln n} = o(1),
\end{aligned}$$

where in the second equality above we note the relationship $(\ln n)^c = n^{o(1)}$ for any

positive constant c and $s_2 \leq n/2$. Therefore, $A_2 = A_{21} + A_{22} = o(1)$ as $n \rightarrow \infty$. It follows from (12) that $P_{12} \leq A_1 + A_2 = o(1)$.

Recall that $G(n, \mathbf{p}_n) \setminus S_1$ contains a component having order no more than $\frac{1}{2}(n - |S_1|)$ if it is not connected. Moreover, when $d_n^{\min} \geq k$, any vertex in S_1 has no less than $k - |S_1| + 1$ neighbors outside S_1 . Recalling the definition of the event $\mathcal{E}(S_1, S_2)$, we know that the probability $P_{12} = o(1)$ implies that if $d_n^{\min} \geq k$, then a.a.s. $G(n, \mathbf{p}_n)$ is k -connected. Hence, (10) holds true. The proof of Theorem 3 is completed. \square

Proof of Theorem 4. Arguing similarly as in the beginning of Theorem 3, we here will rely on the following inequality instead of (3):

$$\min_{1 \leq i \leq n} \min_{\substack{S: i \notin S \\ |S| \geq \lceil n/2 \rceil - k + 1}} \rho_n(i, S) \geq p_n. \quad (13)$$

In view of Definition 1, it suffices to show $\mathbb{P}(\text{Any nonempty set } S \subseteq V \text{ with } |S| \leq \lfloor n/2 \rfloor \text{ is } k\text{-reachable in } G(n, \mathbf{p}_n)) \rightarrow 1 \text{ as } n \rightarrow \infty$.

Let P_0 be the probability that some set of size no more than $\lfloor n/2 \rfloor$ is not k -reachable. Furthermore, let P_l represent the probability that some set of size $l \geq 1$ is not k -reachable in $G(n, \mathbf{p}_n)$. Fix a set $S \in V$ with $|S| = l$. For any vertex $i \in S$,

$$\begin{aligned} & \mathbb{P}(i \text{ has less than } k \text{ neighbors in } V \setminus S) \\ &= \sum_{r=0}^{k-1} \sum_{N_n(i) \setminus S: |N_n(i) \setminus S|=r} \prod_{j \in N_n(i) \setminus S} p_n(e_{ij}) \prod_{j \in V \setminus (S \cup N_n(i))} (1 - p_n(e_{ij})) \\ &\leq \sum_{r=0}^{k-1} \sum_{N_n(i) \setminus S: |N_n(i) \setminus S|=r} (\beta p_n)^r e^{-|V \setminus (S \cup N_n(i))| \cdot \rho_n(i, V \setminus (S \cup N_n(i)))} \\ &\leq \sum_{r=0}^{k-1} \binom{n-l}{r} (\beta p_n)^r e^{-(n-l-r)p_n}, \end{aligned}$$

where we used the neighborhood density condition (13). Since $\binom{n-l}{r} \leq n^r$, the above probability is bounded from above by $k(n\beta p_n e^{p_n})^{k-1} \cdot e^{-(n-l)p_n}$. By independence, the probability that S is not k -reachable is at most $\left(k(n\beta p_n e^{p_n})^{k-1} \cdot e^{-(n-l)p_n}\right)^l$. Therefore,

$$\begin{aligned} P_l &\leq \binom{n}{l} \left(k(n\beta p_n e^{p_n})^{k-1} \cdot e^{-(n-l)p_n}\right)^l \\ &\leq \left(\frac{ekn}{l} (n\beta p_n e^{p_n})^{k-1} \cdot e^{-(n-l)p_n}\right)^l. \end{aligned} \quad (14)$$

Using the assumption of p_n , the right-hand side of (14) equals

$$\begin{aligned} & \left(\frac{ek}{l} (\beta(\ln n + (k-1)\ln \ln n + \omega_n) e^{p_n})^{k-1} \frac{1}{(\ln n)^{k-1}} e^{lp_n - \omega_n} \right)^l \\ & \leq (C\varphi(l)e^{-\omega_n})^l, \end{aligned} \quad (15)$$

where $C > 0$ is a constant and the function $\varphi(l) := e^{lp_n}/l$. A quick study of φ , when seen as a function of a real variable in the interval $[1, n/2]$, shows that it is convex and hits its minimum at p_n^{-1} . Thus its maximum is either $\varphi(1) = e^{p_n}$ or $\varphi(n/2) = O(\frac{1}{\sqrt{n}})$. It follows from (14) and (15) that $P_l \leq (Ce^{1-\omega_n})^l$.

Finally, an application of the union bound yields

$$P_0 \leq \sum_{l=1}^{n/2} (Ce^{1-\omega_n})^l \leq \sum_{l=1}^{\infty} (Ce^{1-\omega_n})^l = \frac{Ce^{1-\omega_n}}{1 - Ce^{1-\omega_n}} = o(1)$$

as $n \rightarrow \infty$, i.e., $\omega_n \rightarrow \infty$. This completes the proof of Theorem 4. \square

Acknowledgements

The author is very grateful to the anonymous referees for their comprehensive comments that helped improve the presentation significantly.

Funding information

There are no funding bodies to thank relating to this creation of this article.

Competing interests

There were no competing interests to declare which arose during the preparation or publication process of this article.

References

- [1] GILBERT, E. N. (1959). Random graphs. *Ann. Math. Statist.* **30**, 1141–1144.
- [2] ERDŐS, P. AND RÉNYI, A. (1959). On random graphs, I. *Publ. Math. Debrecen* **6**, 290–297.
- [3] ERDŐS, P. AND RÉNYI, A. (1961). On the strength of connectedness of a random graph. *Acta Math. Acad. Sci. Hungar.* **12**, 261–267.

- [4] FRIEDGUT, E. AND KALAI, G. (1996). Every monotone graph property has a sharp threshold. *Proc. Amer. Math. Soc.* **124**, 2993–3002.
- [5] LEBLANC, H., ZHANG, H., KOUTSOUKOS, X. AND SUNDARAM, S. (2013). Resilient asymptotic consensus in robust networks. *IEEE J. Sel. Areas Commun.* **31**, 766–781.
- [6] SHANG, Y. (2018). Resilient consensus of switched multi-agent systems. *Syst. Contr. Lett.* **122**, 12–18.
- [7] SHANG, Y. (2022). Resilient cluster consensus of multiagent systems. *IEEE Trans. Syst. Man Cybern. Syst.* **52**, 346–356.
- [8] MUSTAFA, A., MODARES, H. AND MOGHADAM, R. (2020). Resilient synchronization of distributed multi-agent systems under attacks. *Automatica* **115**, 108869.
- [9] ZHANG, H., FATA, E. AND SUNDARAM, S. (2015). A notion of robustness in complex networks. *IEEE Trans. Contr. Netw. Syst.* **2**, 310–320.
- [10] KOVALENKO, I. N. (1971). The theory of random graphs. *Cybern. Syst. Anal.* **7**, 575–579.
- [11] COHEN, J. E. (1986). Connectivity of finite anisotropic random graphs and directed graphs. *Math. Proc. Camb. Phil. Soc.* **99**, 315–330.
- [12] ALON, N. (1995). A note on network reliability. In *Discrete Probability and Algorithms*, D. Aldous, P. Diaconis, J. Spencer, J. M. Steele (eds.), IMA Vol. Math. Appl., 72, Springer-Verlag, New York, 11–14.
- [13] CHUNG, F. AND RADCLIFFE, M. (2011). On the spectra of general random graphs. *Elect. J. Combin.* **18**, #P215.
- [14] SHANG, Y. (2016). Bounding extremal degrees of edge-independent random graphs using relative entropy. *Entropy* **18**, art. no. 53.
- [15] FRIEZE, A. AND KAROŃSKI, M. (2016). *Introduction to Random Graphs*, Cambridge University Press, Cambridge.
- [16] BOLLOBÁS, B., JANSON, S. AND RIORDAN, O. (2007). The phase transition in inhomogeneous random graphs. *Rand. Struct. Algor.* **31**, 3–122.
- [17] JANSON, S., ŁUCZAK, T. AND RUCIŃSKI, A. (2000). *Random Graphs*, John Wiley & Sons, New York.
- [18] USEVITCH, J. AND PANAGOUD, D. (2020). Determining r - and (r, s) - robustness of digraphs using mixed integer linear programming. *Automatica* **111**, 108586.