

# Northumbria Research Link

Citation: McGovern, R., Bogowicz, P., Meader, N., Kaner, E., Alderson, H., Craig, D., Geijer-Simpson, E., Jackson, K., Muir, C., Salonen, D., Smart, D. and Newham, James (2023) The association between maternal and paternal substance use and child substance use, internalising and externalising problems: a systematic review and meta-analysis. *Addiction*, 118 (5). pp. 804-818. ISSN 0965-2140

Published by: Wiley-Blackwell

URL: <https://doi.org/10.1111/add.16127> <<https://doi.org/10.1111/add.16127>>

This version was downloaded from Northumbria Research Link:  
<https://nrl.northumbria.ac.uk/id/eprint/51176/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)

McGovern Ruth (Orcid ID: 0000-0002-4119-4353)

**The association between maternal and paternal substance use and child substance use, internalising and externalising problems: a systematic review and meta-analysis**

McGovern R<sup>1</sup>, Bogowicz P<sup>1</sup>, Meader N<sup>1</sup>, Kaner E<sup>1</sup>, Alderson H<sup>1</sup>, Craig D<sup>1</sup>, Geijer-Simpson E<sup>1</sup>, Jackson K<sup>1</sup>, Muir C<sup>1</sup>, Salonen D<sup>1</sup>, Smart D<sup>1</sup>, Newham JJ<sup>2</sup>.

<sup>1</sup>Population Health Sciences Institute, Newcastle University, United Kingdom

<sup>2</sup>Faculty of Health and Life Sciences, Northumbria University, United Kingdom

**Running head:** Maternal and paternal substance use

**Funding:** Ruth McGovern has been funded to undertake this systematic review by the National Institute for Health Research (NIHR) personal fellowships (NIHR PDF-2014-07-045). Eileen Kaner is in receipt of support from NIHR Senior Investigator Award and is the Director of the NIHR Applied Research Collaboration, North East and North Cumbria. At the time of conducting the review Domna Salonen was supported by the NIHR HEE/NIHR ICA Programme Pre-doctoral Clinical Academic Fellowship (NIHR301146).

**Declaration of interests:** none

Word count: 4149

**Corresponding author:**

Dr Ruth McGovern, Population Health Sciences Institute, Newcastle University, Baddiley-Clark Building, Richardson Road, Newcastle upon Tyne, NE2 4AX, UK

Tel: +44 191 2087994 Fax: +44 191 2086043

Email: [r.mcgovern@ncl.ac.uk](mailto:r.mcgovern@ncl.ac.uk)

**ABSTRACT**

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/add.16127

This article is protected by copyright. All rights reserved.

**Background and aim:** There is substantial evidence showing an association between parental substance use and child substance use and/or mental health problems. Most research focuses upon maternal substance use, with the influence of fathers who use substance often being overlooked. We aimed to investigate the differential effects of maternal and paternal substance use upon children aged 0-18 years.

**Method:** We used systematic review methods to identify observational studies examining the association between either maternal or paternal substance use and child substance use and/or mental health problems. The odds ratio (OR) effect measure was used, for ease of computation. We used a random effects model with the inverse variance method to meta-analyse the findings from eligible studies.

**Findings:** We included 17 unique studies with a total of 47,374 child participants. Maternal and paternal substance use were both associated with increased odds of child any drug use (OR=2.09; 95% confidence interval (CI) [1.53, 2.86]; n=12,349 participants; three studies, and OR=2.86; 95% CI [1.25, 6.54]; n=5,692 participants; three studies, respectively), child alcohol problem use (OR=2.16; 95% CI [1.73, 2.71]; n=7,339 participants; four studies, and OR=1.70; 95% CI [1.36, 2.12]; n=14,219 participants; six studies), child externalising problems (OR=1.81; 95% CI [1.01, 3.22]; n=1,748 participants; three studies, and OR=1.60; 95% CI [1.18, 2.17]; n=2,508 participants; six studies), and child internalising problems (OR=1.60; 95% CI [1.25, 2.06]; n=1,748 participants; three studies, and OR=1.42; 95% CI [1.12, 1.81]; n=2,248 participants; five studies). Child any alcohol use was associated with maternal substance use only (OR=2.26; 95% CI [1.08, 4.70]; n=28,691 participants; five studies).

**Conclusion:** Both maternal and paternal substance use are associated with child substance use and mental health problems.

**Key words:** parental substance use; child substance use; child internalising problems; child externalising problems

## INTRODUCTION

It is estimated that between 5% and 30% of all children in Europe live with at least one parent who misuses alcohol and/or drugs (1). In England, the Children's Commissioner data reports that in 2019-2020 there were 478,000 children living with a problem alcohol or drug using parent; a rate of 40 in 1000 children (2). Parents substantially influence the development of their children through a range of factors including genetics, the physical environment they provide, child-rearing practices and relationship quality (3, 4). When the parent has a substance use disorder, a number of those mechanisms may come into play, creating adverse conditions which can have profound and lasting effect (5). This includes an increased likelihood of child substance use including alcohol consumption (6-9), alcohol

intoxication (7, 9-11), and alcohol use disorders (12, 13), as well as illicit drug use (9, 11, 14). Further, association has been found between parental substance use and child externalising (12, 15) and internalising problems such as behavioural problems, depression and anxiety (16). Children and adolescents who use substances and/or experience mental health problems are more likely to experience poor outcomes in childhood than their peers, including increased disease burden (17, 18). Substance use in adolescence is also associated with criminal involvement, unprotected or regretted sexual activity, self-harm and suicide (19) and the later development of substance dependency (20). Adolescents who experience mental health problems have been found to be at increased risk of not completing secondary school, later unemployment, and unplanned pregnancy/parenthood (21). Moreover problems with substance use and/or mental health in childhood often persist in adulthood (18, 22), resulting in reduced life opportunities (21).

Whilst numerous studies have examined the influence of parental substance use upon the child, few have considered parental gender (23). These studies mostly report on a majority maternal sample due to mothers typically being the primary caregiver or their relationship with the child often being perceived to be central to the child development (24). Similarly, recent systematic reviews have either examined the impact of parental substance use on child outcomes without examining parental gender (25), or examined the impact of maternal substance use during pregnancy (26), wherein there may be very different mechanisms of transmission to that of post-birth parental substance use. Fathers have long been considered to be the 'forgotten contributors' (27), or research into their influence has been restricted to examining the effect of absent versus present fathers (28). The invisibility of fathers in research impedes our understanding of paternal effects upon children's substance use and other health problems, as well as our ability to develop appropriate preventative strategies and family interventions (29). More recently, research has begun to examine the role of fathers in child development (23), including how paternal substance use may influence child substance use and mental health outcomes. Studies that include gender have reported mixed results, with some highlighting the significance of maternal substance use (15, 30, 31), whilst other studies found only paternal and not maternal substance use to be positively associated with adolescent substance use (6, 14, 32). Given the importance of understanding the etiology of children's substance use and mental health disorders, it is imperative to investigate the differential effects of maternal and paternal substance use upon children.

We undertook a systematic review and meta-analyses of published studies. Our primary objective was to examine the association between maternal and paternal substance use and child substance use. Our secondary objective was to examine the association between maternal and paternal substance use and child externalising and internalising problems. To our knowledge, this is the first systematic review and meta-analysis examining these associations.

## **METHODS**

The review protocol was registered with PROSPERO (CRD42017070337). This protocol specified that substance misuse/abuse below the diagnostic threshold for dependency would be examined. However, due to inconsistency in how substance use levels were reported within papers, an amendment to the protocol was registered in February 2021 to include all substance use disorders including dependency. The following electronic databases were searched from inception to March 2022, using free text keywords and thesaurus headings: Medline (OVID), PsycInfo (OVID), CINAHL (EBSCO), SCOPUS, Applied Social Science Index and Abstract (ProQuest), International Bibliography of Social Science (ProQuest), ProQuest Criminal Justice (ProQuest), ProQuest Social Science Journals (ProQuest), ProQuest Sociology (ProQuest), Social Service Abstracts (ProQuest), Sociological Abstracts (ProQuest). No language or date restrictions were applied. Database searches were supplemented by searching for grey literature via SCOPUS and ProQuest and on key websites including Google and Google Scholar and hand-searching reference lists of relevant studies.

Two reviewers independently screened all titles and abstracts using pre-specified inclusion and exclusion criteria. Full papers for all potentially eligible studies were retrieved and evaluated for final inclusion. Relevant data were extracted independently by two reviewers, including: study design, sample characteristics, nature of parental substance use and child outcome. Discrepancies at each stage were resolved by discussion or by consulting a third reviewer if consensus could not be reached.

## **Eligibility**

Studies were included if they used a quantitative observational design, had a sample of children aged less than 18 years, and estimated the association between maternal and/or paternal substance use disorder (alcohol or drugs) and child substance use and/or mental health outcomes. We included studies which identified substance use disorders and mental health outcomes by a reliable, valid, formal assessment (validated screening tool, assessment by a health or child welfare practitioner) or diagnostic tool (DSM III, DSM IIIR, DSM IV, ICD8, ICD 9, ICD 10), or both. Studies were required to compare cohorts of children whose parents had a substance use disorder to cohorts of children whose parents did not have a substance use disorder. Where insufficient data were provided to permit meta-

analysis, we contacted the study authors to request these data. If we were unable to obtain this information, we excluded the study from the meta-analysis.

### **Quality appraisal**

We assessed study quality using the Newcastle-Ottawa Scale (33). The scale assesses quality in observational studies relating to three domains 1) selection of study groups; 2) comparability of groups and 3) ascertainment of exposure and outcomes. A star system is used to allow for semi-quantitative assessment of study quality. A total quality score of each individual study was calculated by adding all the stars (range = 0–9, with a higher score indicating higher overall quality).

### **Data synthesis**

Multiple meta-analyses were planned and carried out with a minimum requirement of three studies per meta-analysis. These were: by parental gender, outcome type (child substance use; externalising problems and internalising problems), and level of child risk (for substance use examining any alcohol/drug use and alcohol problem use). We defined alcohol problem use as weekly or more frequent and/or above the adult daily recommended consumption levels in line with the English Chief Medical Officer recommendations. Externalising problems included a range of maladaptive behaviours directed towards the child's environment (such as conduct disorders, antisocial behaviour, opposition disorder and non-specified externalising problems). Internalising problems included disturbances in emotion and mood (such as depression and anxiety). The odds ratio (OR) effect measure was used, for ease of computation. Data pertaining to parental alcohol use was prioritised over parental drug data and unadjusted OR estimates were prioritised over adjusted estimates. These decisions were taken to manage the impact of variation between studies in terms of the risk exposure and selection of covariates. Unadjusted OR estimates were derived from reported data, where possible. If no unadjusted estimate was available, the main adjusted estimate or estimate adjusted for the most covariates was used. Standardised mean difference data were converted to OR (34). Estimates stratified by gender and/or age were pooled using within-study random effects meta-analysis prior to inclusion in the main meta-analyses.

All meta-analyses used a random effects model with the inverse variance method for pooling OR estimates along with 95% confidence intervals. The restricted maximum likelihood method was used to estimate between-study variance (35). Forest plots and aggregate summary tables were obtained. Heterogeneity was explored using a range of statistics, including I-squared and tau-squared. Baujat plots were obtained but these were difficult to interpret due to the small number of studies (36).

Sensitivity analyses were carried out where data was available by substituting parental drug use for parental alcohol use, excluding adjusted estimates, excluding converted or within-study pooled estimates, and excluding estimates derived from cross-sectional studies. A leave-one-out influence analysis was carried out to find influential study estimates.

It was not possible to perform extensive meta-regression analyses due to the small number of studies in each analysis being below the recommended levels to produce robust estimates (37).

Multilevel network meta-regression analyses (38) were only used to assess the impact of small study effects, including publication bias. These analyses used Bayesian simulation methods, following established methodology. For all analyses, we included a burn-in of 30,000 iterations subsequently discarded after convergence was confirmed. We ran an additional 70,000 iterations to produce posterior estimates. We compared the goodness of fit between a 'small study bias adjustment' model compared with the same model without bias adjustment using total residual deviance (mean similar to the number of data points in the model indicates reasonable fit), between-study standard deviation (SD) (a smaller SD suggests bias adjustment is accounting for some of the heterogeneity across studies) and deviance information criterion (DIC, 3-5 points difference between models is generally agreed to be substantial).

Data management was carried out using Microsoft Excel (Microsoft Corp, Redmond, Washington, USA). Data analysis was carried out using R (R Foundation for Statistical Computing, Vienna, Austria) in RStudio (Rstudio, Boston, Massachusetts, USA) and WinBUGS (The BUGS Project, UK). The code is available via contact with the corresponding author.

## RESULTS

Our search identified 4365 potentially relevant references. Of those, 236 full papers were retrieved. Seventeen studies met the inclusion criteria and were included in our meta-analysis. One of these studies was included after the author provided additional, unpublished data (39).

[INSERT FIGURE 1]

The 17 included unique studies consisted of a total of 47,374 child participants. The age of the child participants within the studies ranged from 9 to 17 years at time of follow-up. All parents within the studies were biological relatives. Fifteen studies included data associated with parental alcohol misuse (7, 8, 39-51), four included data for illicit drug use (42, 45, 47, 52) and one included data on substance use (alcohol and/or drugs combined) (53). Fifteen studies examined child exposure to paternal substance use (7, 8, 40, 43-54) and 11 studies examined exposure to maternal substance use (7, 39, 41-45, 48, 49, 52, 53). Study outcomes varied with 12 studies reporting on child substance use (7, 8, 39-46, 48, 51), seven studies reporting on child externalising problems (39, 40, 47, 48, 50, 51, 53) and six studies

reporting on child internalising problems (39, 47, 48, 50, 51, 53). Six of the studies used a longitudinal design (8, 41, 42, 44, 45, 52), six were cross sectional studies (7, 40, 43, 46, 49, 50) and five were cohort studies (39, 47, 48, 51, 53). Six studies reported adjusted outcomes only (8, 41, 43, 46, 48, 53). Seven of the studies were conducted in USA (8, 39, 47, 48, 51-53), three in Australia (40, 41, 43), two in the UK (42, 45), and one each from Canada (50), Barbados (49), Finland (44), Norway (7) and Slovakia (46).

The quality of the studies varied. The most common risk of bias was the certainty of exposure to the risk factor. Whilst all studies described a reliable assessment of parental substance use, a minority utilised a validated tool or applied diagnostic criteria (39, 47, 50, 51, 53, 55). Whilst most studies were large cohort studies, with randomly selected and/or representative samples, some studies included selected samples (8, 40, 47, 52) or included small samples and did not provide a sample size calculation (40, 47, 48, 53). The majority of the studies provided a detailed description of the statistical analysis conducted however two studies failed to do so (8, 49).

## **INSERT TABLE 1 & TABLE 2**

### **Child substance use**

We investigated the odds of substance use in children exposed to maternal and/or paternal substance use. We found that children who were exposed to maternal substance use were more likely than children who were not exposed to use any alcohol (OR=2.26; 95% CI [1.08, 4.70]; 28,691 participants; five studies); use any drugs (OR=2.09; 95% CI [1.53, 2.86]; 12,349 participants; three studies) and to use alcohol problematically (OR=2.16; 95% CI [1.73, 2.71]; 7,339 participants; four studies). Children who were exposed to paternal substance use were more likely to use drugs (OR=2.86; 95% CI [1.25, 6.54]; 5,692 participants; three studies) and to use alcohol problematically (OR=1.70; 95% CI [1.36, 2.12]; 14,219 participants; six studies). Additionally, they were found to be more likely to use any alcohol than children who were not exposed to parental substance use however the 95% CI was wide and includes no association (OR=1.89; 95% CI [0.93, 3.83]; 28,429 participants; five studies).

### **Child externalising and internalising problems**

We investigated the odds of externalising and internalising problems in children exposed to maternal and paternal substance use. We found that children who were exposed to maternal substance use were more likely than children who were not exposed to experience externalising problems, although the interval was wide and included essentially no association (OR=1.81; 95% CI [1.01, 3.22]; 1,748 participants; three studies). Exposed children were also more likely to experience internalising problems (OR=1.60; 95% CI [1.25, 2.06]; 1,748 participants; three studies). Children who were exposed to paternal substance



use were more likely than children who were not exposed to parental substance use to experience externalising problems (OR=1.60; 95% CI [1.18, 2.17]; 2,508 participants; six studies) and more likely to experience internalising problems (OR=1.42; 95% CI [1.12, 1.81]; 2,248 participants; five studies).

### **INSERT TABLE 3**

#### **Heterogeneity**

According to the  $I^2$  statistic, substantial heterogeneity was observed across studies examining exposure to maternal substance use and child any alcohol use ( $I^2=86\%$ ;  $t^2=0.5879$ ;  $p<0.01$ ); child any drug use ( $I^2=66\%$ ;  $t^2=0.0583$ ;  $p=0.03$ ) and child externalising problems ( $I^2=73\%$ ;  $t^2=0.1551$ ;  $p<0.01$ ). The Baujat plot for the child any alcohol use analysis visually suggested that Oshi (2018) was heterogeneous and influential. This study was conducted in Barbados, with important cultural and healthcare differences. The other maternal Baujat plots were difficult to meaningfully interpret.

Substantial heterogeneity was also observed in studies examining exposure to paternal substance use and child any alcohol use ( $I^2=94\%$ ;  $t^2=0.5696$ ;  $p<0.01$ ). The Baujat plot for this analysis again indicated that Oshi (2018) was heterogeneous and influential. The Baujat plot for child problem alcohol use indicated that Jennison (2014) contributed substantially to heterogeneity ( $I^2=0.47$ ;  $t^2=0.04$ ;  $p=0.09$ ). The sampling approach used in this study may have led to (a cross-sectional sample that was supplemented with oversampling of African American, Hispanic and economically disadvantaged White youths). The plot for child externalising problems indicated that McCauley Ohannessian (2014) contributed substantially to heterogeneity ( $I^2=0.16$ ;  $t^2=0.04$ ;  $p=0.31$ ). It is unclear why the results of this study differed substantially from other included studies.

#### **Sensitivity analysis**

The analyses using parental drug use rather than alcohol use all had larger magnitude ORs for both fathers (child any alcohol: OR=2.28; 95% CI [1.28, 4.07]; child externalising problems: OR=2.15; 95% CI [1.17, 3.95]; child internalising problems: OR=1.63; 95% CI [0.74, 3.59]) and mothers (child any alcohol use: OR=2.41; 95% CI [1.16, 4.97]). The confidence interval for paternal substance use and child any alcohol use became significant (OR: 2.28, 95% CI [1.28 to 4.07]) while the interval for paternal substance use and child internalising problems became wide and included no association (OR: 1.63, 95% CI [0.74 to 3.59]).

The analyses excluding adjusted estimates had larger magnitude ORs for paternal substance use and child externalising problems (OR=2.89; 95% CI [1.44, 5.83]) and internalising problems (OR=2.38; 95% CI [0.81, 7]). The confidence interval for maternal substance use and child any alcohol use became wide and included no association (OR: 2.32, 95% CI [0.88 to 6.09]). It was not possible to carry out sensitivity analysis for four analyses due to insufficient numbers of studies.

There were no significant differences in OR magnitude or confidence intervals for all analyses excluding converted or within-study pooled estimates. It was not possible to carry out sensitivity analysis for six analyses due to insufficient numbers of studies.

There were no significant differences in OR magnitude for all analyses excluding cross sectional data. The confidence interval for paternal substance use and child externalising problems became insignificant (OR: 1.36, 95% CI [0.95 to 1.95]). It was not possible to carry out sensitivity analysis for two analyses due to insufficient numbers of studies.

The influence analysis found that there was at least one influential study estimate in each of the meta-analyses, including Oshi (2018), identified as influential in the Baujat plots. This study also appears to be an outlier in the analysis for paternal substance use and child any alcohol use. This association became significant when the study was omitted (OR=1.54; 95% CI [1.06; 2.25]).

Full details of sensitivity analyses are included in the supplementary file.

### **Small study effects**

There was no evidence of small study bias in any analyses. The credible intervals (95% CrIs) for study size covariates (variance of treatment effect was used as a proxy for study size) were very wide for all outcomes. Total residual deviance for all models was similar to the number of datapoints, indicating reasonable goodness of fit. There were negligible differences in goodness of fit between the bias adjustment model and simpler model without covariates for all outcomes (see table 4).

INSERT TABLE 4

## **DISCUSSION**

This systematic review and meta-analysis found evidence that both maternal and paternal substance use are associated with child substance use and mental health problems. All outcomes except paternal substance use and child any alcohol use were found to be significantly associated. Similarly, after excluding maternal studies which used adjusted ORs, maternal substance use was not associated with child any alcohol use. Epidemiological studies estimating lifetime prevalence of alcohol use in adolescence have reported a 40% prevalence rate in children aged 10-17 years (56). Rates of alcohol consumption have been found to increase considerably with age ranging from just 4% for those aged 10 to 90% for those aged 17 years(57). This high prevalence of alcohol use in adolescent populations may provide some explanation as to why the cohorts of children of substance using parents within our meta-analysis (which is mostly children within mid to late adolescence) may not be more likely to report any alcohol use than their non-exposed peers. Rather, our findings highlight the vulnerability within these children to progress beyond experimentation to risky substance use and dependency and to experience mental health problems.

Reducing the number of children using substances is of international priority (58-61). Further, there is growing concern about the prevalence of mental health problems in children and adolescents with a recent UK survey finding 1 in 6 children aged 5-16 years have a mental health problem (62). Whilst it is acknowledged that children of substance using parents may have genetic vulnerability (63, 64), there is a large literature showing that environmental risk factors may play an influential part in the intergenerational transmission of substance use and mental health problems (3, 65-67). The drugs strategy in England aspires to achieve a 'generational shift in the use of drugs in society, [where] fewer people take drugs or feel drawn toward taking drugs, and today's children and young people grow up in a safer and healthier environment', and identifies children whose parents are dependent upon substances as a vulnerable group at risk of substance use and mental health problems (60). Between 60% and 80% of parents with substance use problems do not receive treatment (68). Better engagement of parents in drug and alcohol treatment may therefore contribute to reducing the prevalence of child substance use. However, it is unlikely that providing treatment to substance using parents only will be enough to ameliorate vulnerability in affected children. Rather children whose parents use substances may require support in their own right (69, 70) to mitigate risk coming from genetic vulnerability (71), reduce intergenerational trauma (72) and the impact of the adversity (5). Supportive interventions for children who experience maternal and/or paternal substance use may also be needed to meet their needs as young carers (73).

Policies and practice approaches to address parental substance use typically refer to 'parents', failing to distinguish between 'mothers' and 'fathers' (74). This results in 'gender blindness' rather than gender neutrality (75), wherein the focus upon mothers as the primary caregiver and recipient of intervention is reinforced and the role of fathers is overlooked. This gender blindness is also reported within provision for vulnerable families highlighting a tendency of services to target mothers and a failure to engage fathers, due to viewing them as less important to child outcomes (76), 'hard to reach' (77) or as a risk to exclude rather than a potential family resource (78). Moreover, gender blindness contributes to the 'bad mothers' (79) and 'invisible fathers' (80) discourse. This stigmatises mothers making it harder for them to access and benefit from drug treatment (81), and neglects the importance of addressing fathers' substance use. A series of systematic reviews have demonstrated the effectiveness of integrated parenting interventions at reducing parental substance use (82-86). These interventions combine parenting skills training with drug and alcohol treatment. This evidence base has resulted in growing availability of parenting programmes for substance using mothers. Despite suggestion that integrated parenting interventions may be of benefit to fathers (82, 83, 87), these services are rarely available to fathers (82, 83, 87) or fail to engage them due to perceptions that the interventions are maternal services (88). Consequently, there is a need for substance use

interventions to be developed for fathers, with a focus upon the specific issues substance using fathers may experience (87, 89).

### Limitations

There are some limitations with our findings. Our analysis examines maternal and paternal substance use separately, thus making an important contribution to understanding the differential impact of parental substance use upon children. However, many children may be affected by the substance use of both parents or by caregivers who are not their biological parents. It is likely that having more than one parent/caregiver who use substances will increase the odds of child substance use or mental health problems (90). Few of the studies reported the residency status of the parents. Therefore, it is possible that children had varying degrees of exposure to parental alcohol and substance use. Further, parental substance use is likely to occur within a context of multiple family adversity (91), wherein other risk factors co-exist and accumulate with greater impact (5). It is possible that there may be other drivers of vulnerability in children. For example, parental substance use and child poverty often co-exist and poverty has been found to be associated with child substance use and mental health problems (92). Further research examining the clustering of parental substance use with other risk factors, including poverty is needed in order to understand more clearly the burden of parental substance use upon children.

The results of analyses using converted mean difference data (paternal substance use and child externalising and internalising problems) should be interpreted with caution, due to uncertainty in relation to reported proportions in exposed and unexposed groups (47) and the difficulties in interpreting ORs derived from continuous measures.

It should be noted that some of the meta-analyses showed substantial heterogeneity. These studies all relied upon self-report measures and often did not use validated tools for substance use. Whilst sample sizes ranged from 1,748- 28,685 participants per analysis, some of our pooled samples came from combining a small number of studies. It was not possible to fully explore the possible causes of heterogeneity due to the limited numbers of studies. The exposure hierarchy assumptions did not appear to have any significant impact on the findings. The findings of the sensitivity analyses excluding adjusted, converted ORs and cross-sectional studies should be interpreted with caution, given that it was not possible to carry out at least 50% of such analyses.

The quality of the evidence in the review varied. A minority of the studies included in this review used validated tools to assess risk exposure with some utilising imprecise measures. We intended to use a funnel plot to investigate publication bias; however, as there were fewer than 10 trials in each of our meta-analysis, the minimum number of trials required to enable a funnel plot was not met (93, 94). To mitigate this limitation, we have explored the

impact of small study effects using Bayesian methods (38). Whilst this found there was no evidence of publication bias, we are unable to rule this out.

## CONCLUSION

Both maternal and paternal substance use is associated with child substance use, dependency and mental health problems. To reduce vulnerability in children of substance using parents, fathers' substance use as well as mothers' should be targeted by drug policy and family support services. It may be necessary to develop substance use treatment specifically for fathers. Further research is needed to estimate the effectiveness of interventions for fathers.

## ACKNOWLEDGMENTS

Ruth McGovern has been funded to undertake this systematic review by the National Institute for Health Research (NIHR) personal fellowships (NIHR PDF-2014-07-045). Eileen Kaner is in receipt of support from NIHR Senior Scientist Award. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, or the Department of Health.

The authors thank Lachlan Fotheringham for assistance with coding.

## REFERENCES

1. European Monitoring Centre for Drugs and Drug Addiction (EMCDDA). Children's Voices: Experiences and perceptions of European children on drug and alcohol issues. Lisbon: EMCDDA; 2010.
2. Children's Commissioner. Childhood vulnerability in England. <https://www.childrenscommissioner.gov.uk/chldrnl/>; 2020.
3. Hettema JM, Prescott CA, Myers JM, Neale MC, Kendler KS. The structure of genetic and environmental risk factors for anxiety disorders in men and women. *Arch Gen Psychiatry*. 2005;62(2):182-9.
4. Kendler KS, Prescott CA, Myers J, Neale MC. The structure of genetic and environmental risk factors for common psychiatric and substance use disorders in men and women. *Arch Gen Psychiatry*. 2003;60(9):929-37.
5. Hughes K, Bellis M, Hardcastle K, Sethi D, Butchart A, Mikton C, et al. The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis. *The Lancet Public Health*. 2017;2(8):e356-e66.
6. Cranford JA, Zucker RA, Jester JM, Puttler LI, Fitzgerald HE. Parental Alcohol Involvement and Adolescent Alcohol Expectancies Predict Alcohol Involvement in Male Adolescents. *Psychology of Addictive Behaviors*. 2010;24(3):386-96.

7. Haugland SH, Holmen TL, Ravndal E, Bratberg GH. Parental alcohol misuse and hazardous drinking among offspring in a general teenage population: gender-specific findings from the Young-HUNT 3 study. *BMC Public Health*. 2013;13:1140.
8. Jennison KM. The impact of parental alcohol misuse and family environment on young people's alcohol use and behavioral problems in secondary schools. *Journal of Substance Use*. 2014;19(1-2):206-12.
9. Keeley HS, Mongwa T, Corcoran P. The association between parental and adolescent substance misuse: Findings from the Irish CASE study. *Irish Journal of Psychological Medicine*. 2015;32(Spec Iss1):107-16.
10. Haugland SH, Holmen TL, Krokstad S, Sund ER, Bratberg GH. Intergenerational Hazardous Alcohol Use and Area Factors: The HUNT Study, Norway. *Substance Use & Misuse*. 2015;50(14):1753-64.
11. Haugland SH, Strandheim A, Bratberg G. Is high-risk use of intoxicants more common among adolescents who have seen their parents intoxicated? *Tidsskrift for Den Norske Laegeforening*. 2012;132(4):410-3.
12. Kendler KS, Gardner CO, Edwards A, Hickman M, Heron J, Macleod J, et al. Dimensions of parental alcohol use/problems and offspring temperament, externalizing behaviors, and alcohol use/problems. *Alcoholism: Clinical & Experimental Research*. 2013;37(12):2118-27.
13. Lieb R, Merikangas KR, Hofler M, Pfister H, Isensee B, Wittchen HU. Parental alcohol use disorders and alcohol use and disorders in offspring: a community study. *Psychological Medicine*. 2002;32(1):63-78.
14. Shorey RC, Fite PJ, Elkins SR, Frissell KC, Tortolero SR, Stuart GL, et al. The association between problematic parental substance use and adolescent substance use in an ethnically diverse sample of 9th and 10th graders. *Journal of Primary Prevention*. 2013;34(6):381-93.
15. Torvik FA, Rognmo K, Ask H, Roysamb E, Tambs K. Parental alcohol use and adolescent school adjustment in the general population: results from the HUNT study. *BMC Public Health*. 2011;11:706.
16. Seay KD. Pathways From Parental Substance Use to Child Internalizing and Externalizing Behaviors in a Child Protective Services Sample. *Child Maltreatment*. 2020;25(4):446-56.
17. Degenhardt L, Stockings E, Patton G, Hall WD, Lynskey M. The increasing global health priority of substance use in young people. *Lancet Psychiatry*. 2016;3(3):251-64.
18. Wykes T, Bell A, Carr S, Coldham T, Gilbody S, Hotopf M, et al. Shared goals for mental health research: what, why and when for the 2020s. *Journal of mental health (Abingdon, England)*. 2021:1-9.
19. Newbury-Birch D, Walker J, Avery L, Beyer F, Brown N, Jackson K, et al. Impact of Alcohol Consumption on Young People: A systematic review of published reviews. 2009.
20. McCambridge J, McAlaney J, Rowe R. Adult consequences of late adolescent alcohol consumption: a systematic review of cohort studies. *PLoS Med*. 2011;8(2):e1000413-e.

21. Clayborne Z, Varin M, Colman I. Systematic Review and Meta-Analysis: Adolescent Depression and Long-Term Psychosocial Outcomes. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2019;58(1):72-9.
22. Yoon G, Westermeyer J, Kuskowski MA, Nesheim L. Impact of the number of parents with alcohol use disorder on alcohol use disorder in offspring: a population-based study. *Journal of Clinical Psychiatry*. 2013;74(8):676-81.
23. Barker B, Iles JE, Ramchandani PG. Fathers, fathering and child psychopathology. *Curr Opin Psychol*. 2017;15:87-92.
24. Weijers D, van Steensel FJA, Bögels SM. Associations between Psychopathology in Mothers, Fathers and Their Children: A Structural Modeling Approach. *J Child Fam Stud*. 2018;27(6):1992-2003.
25. Kuppens S, Moore SC, Gross V, Lowthian E, Siddaway AP. The Enduring Effects of Parental Alcohol, Tobacco, and Drug Use on Child Well-being: A Multilevel Meta-Analysis. *Dev Psychopathol*. 2020;32(2):765-78.
26. Ruisch IH, Dietrich A, Glennon JC, Buitelaar JK, Hoekstra PJ. Maternal substance use during pregnancy and offspring conduct problems: A meta-analysis. *Neuroscience & Biobehavioral Reviews*. 2018;84:325-36.
27. Lamb ME. Fathers: Forgotten Contributors to Child Development. *Human Development*. 1975;18(4):245-66.
28. Flouri E. Fathers' behaviors and children's psychopathology. *Clin Psychol Rev*. 2010;30(3):363-9.
29. Davison KK, Charles JN, Khandpur N, Nelson TJ. Fathers' Perceived Reasons for Their Underrepresentation in Child Health Research and Strategies to Increase Their Involvement. *Maternal and Child Health Journal*. 2017;21(2):267-74.
30. Kerr DCR, Capaldi DM, Pears KC, Owen LD. Intergenerational influences on early alcohol use: Independence from the problem behavior pathway. *Development and Psychopathology*. 2012;24(3):889-906.
31. Yule AM, Wilens TE, Martelon MK, Simon A, Biederman J. Does Exposure to Parental Substance Use Disorders Increase Substance Use Disorder Risk in Offspring? A 5-year Follow-Up Study. *The American Journal on Addictions*. 2013;22(5):460-5.
32. Ohannessian CM. Parental Problem Drinking and Adolescent Psychosocial Adjustment: The Mediating Role of Adolescent-Parent Communication. *Journal of Research on Adolescence*. 2012;22(3):498-511.
33. Wells G, Shea B, O'Connell D, Peterson J, Welch V, Losos M. The Newcastle–Ottawa Scale (NOS) for assessing the quality of non randomised studies in meta-analyses. [https://www.ohri.ca//programs/clinical\\_epidemiology/oxford.asp](https://www.ohri.ca//programs/clinical_epidemiology/oxford.asp): Ottawa Hospital Research Institute; 2015 [
34. Borenstein M, Hedges L, Higgins J, Rothstein H. Introduction to meta-analysis. . Chichester John Wiley & Sons; 2009.

35. Langan D, Higgins JPT, Jackson D, Bowden J, Veroniki AA, Kontopantelis E, et al. A comparison of heterogeneity variance estimators in simulated random-effects meta-analyses. *Res Synth Methods*. 2019;10(1):83-98.
36. Baujat B, Mahé C, Pignon J-P, Hill C. A graphical method for exploring heterogeneity in meta-analyses: application to a meta-analysis of 65 trials. *Statistics in Medicine*. 2002;21(18):2641-52.
37. Hedges LV, Tipton E, Johnson MC. Robust variance estimation in meta-regression with dependent effect size estimates. *Research Synthesis Methods*. 2010;1(1):39-65.
38. Dias S, Welton NJ, Marinho VCC, Salanti G, Higgins JPT, Ades AE. Estimation and adjustment of bias in randomized evidence by using mixed treatment comparison meta-analysis. *Journal of the Royal Statistical Society: Series A (Statistics in Society)*. 2010;173(3):613-29.
39. Malone SM, McGue M, Iacono W. Mothers' maximum drinks ever consumed in 24 hours predicts mental health problems in adolescent offspring. *Journal of Child Psychology & Psychiatry*. 2010;51(9):1067-75.
40. Cumes-Rayner D, Lucke J, Singh B, Adler B, Lewin T, Dunne M, et al. A high-risk community study of paternal alcohol consumption and adolescents' psychosocial characteristics. *Journal of Studies on Alcohol*. 1992;53(6):626-35.
41. Alati R, Baker P, Betts KS, Connor JP, Little K, Sanson A, et al. The role of parental alcohol use, parental discipline and antisocial behaviour on adolescent drinking trajectories. *Drug and Alcohol Dependence*. 2014;134:178-84.
42. Heron J, Barker ED, Joinson C, Lewis G, Hickman M, Munafo M, et al. Childhood conduct disorder trajectories, prior risk factors and cannabis use at age 16: birth cohort study. *Addiction*. 2013;108(12):2129-38.
43. Kelly AB, Toumbourou JW, O'Flaherty M, Patton GC, Homel R, Connor JP, et al. Family Relationship Quality and Early Alcohol Use: Evidence for Gender-Specific Risk Processes. *Journal of Studies on Alcohol and Drugs*. 2011;72(3):399-407.
44. Korhonen T, Huizink A, Dick D, Pulkkinen L, Rose R, Kaprio J. Role of individual, peer and family factors in the use of cannabis and other illicit drugs: a longitudinal analysis among Finnish adolescent twins. *Drug and Alcohol Dependence*. 2008;97(1-2):33-43.
45. Macleod J, Hickman M, Bowen E, Alati R, Tilling K, Smith GD. Parental drug use, early adversities, later childhood problems and children's use of tobacco and alcohol at age 10: Birth cohort study. *Addiction*. 2008;103(10):1731-43.
46. Rehorcikova V, Nemcovska E, Sklenarova Z, Kallay A, Kallayova D, Brazinova A, et al. Impact of family level factors on alcohol drinking in primary school children. *Central European Journal of Public Health*. 2013;21(4):202-6.
47. Fals-Stewart W, Kelley ML, Fincham FD, Golden J, Logsdon T. Emotional and behavioral problems of children living with drug-abusing fathers: Comparisons with children living with alcohol-abusing and non-substance-abusing fathers. *Journal of Family Psychology*. 2004;18(2):319-30.



48. Ohannessian CM, Hesselbrock VM, Kramer J, Bucholz KK, Schuckit MA, Kuperman S, et al. Parental substance use consequences and adolescent psychopathology. *J Stud Alcohol*. 2004;65(6):725-30.
49. Oshi SN, Abel WD, Oshi DC, Smith PW, Ricketts Roomes TF, Meka IA, et al. Parental Alcohol Drinking Habit as a Predictor of Alcohol Use among Secondary School Students in Barbados. *Asian Pac J Cancer Prev*. 2018;19(S1):57-62.
50. Carbonneau R, Tremblay RE, Vitaro F, Dobkin PL, Saucier J-F, Pihl RO. Paternal alcoholism, paternal absence and the development of problem behaviors in boys from age six to twelve years. *Journal of Studies on Alcohol*. 1998;59(4):387-98.
51. Malone SM, Iacono WG, McGue M. Drinks of the father: father's maximum number of drinks consumed predicts externalizing disorders, substance use, and substance use disorders in preadolescent and adolescent offspring. *Alcoholism: Clinical & Experimental Research*. 2002;26(12):1823-32.
52. Henry KL, Augustyn MB. Intergenerational continuity in cannabis use: The role of parent's early onset and lifetime disorder on child's early onset. *Journal of Adolescent Health*. 2016(Pagination):No Pagination Specified.
53. Merikangas KR, Dierker LC, Szatmari P. Psychopathology among offspring of parents with substance abuse and/or anxiety disorders: a high-risk study. *Journal of Child Psychology and Psychiatry and Allied Disciplines*. 1998;39(5):711-20.
54. Sibley MH, Pelham WE, Molina BS, Cox S, Kipp H, Gnagy EM, et al. The role of early childhood ADHD and subsequent CD in the initiation and escalation of adolescent cigarette, alcohol, and marijuana use. *Journal of Abnormal Psychology*. 2014;123(2):362-74.
55. Macleod J, Hickman M, Bowen E, Alati R, Tilling K, Smith GD. Parental drug use, early adversities, later childhood problems and children's use of tobacco and alcohol at age 10: birth cohort study.[Erratum appears in *Addiction*. 2008 Dec;103(12):2075]. *Addiction*. 2008;103(10):1731-43.
56. Donoghue K, Rose H, Boniface S, Deluca P, Coulton S, Alam MF, et al. Alcohol Consumption, Early-Onset Drinking, and Health-Related Consequences in Adolescents Presenting at Emergency Departments in England. *J Adolesc Health*. 2017;60(4):438-46.
57. ESPAD Group. ESPAD Report 2019: Results from the European School Survey Project on Alcohol and Other Drugs. Luxembourg: EMCDDA Joint Publications, Publications Office of the European Union; 2020.
58. Council of The European Union. EU Drugs Strategy 2021-2025. <https://www.emcdda.europa.eu/system/files/attachments/13642/Council%20of%20the%20EU%20%282020%29%20EU%20Drugs%20Strategy%202021-25.pdf2020>.
59. Department of Health. National Drug Strategy 2017-2026. Australia: Commonwealth of Australia; 2017.
60. HM Government. From harm to hope: A 10-year drugs plan to cut crime and save lives. <https://www.gov.uk/government/publications/from-harm-to-hope-a-10-year-drugs-plan-to-cut-crime-and-save-lives2021>.
61. Office of National Drug Control Policy. The Biden-Harris Administration's Statement of Drug Policy Priorities for Year One. USA: ONDCP; 2021.

62. NHS Digital. Mental Health of Children and Young People in England, 2020: Wave 1 follow up to the 2017 survey. 2020.
63. Goldman D, Oroszi G, Ducci F. The genetics of addictions: uncovering the genes. *Nat Rev Genet.* 2005;6(7):521-32.
64. Kendler KS, Prescott CA. Cannabis use, abuse, and dependence in a population-based sample of female twins. *Am J Psychiatry.* 1998;155(8):1016-22.
65. Kendler KS, Schmitt E, Aggen SH, Prescott CA. Genetic and environmental influences on alcohol, caffeine, cannabis, and nicotine use from early adolescence to middle adulthood. *Arch Gen Psychiatry.* 2008;65(6):674-82.
66. Boomsma DI, van Beijsterveldt CE, Hudziak JJ. Genetic and environmental influences on Anxious/Depression during childhood: a study from the Netherlands Twin Register. *Genes Brain Behav.* 2005;4(8):466-81.
67. Meyers JL, Dick DM. Genetic and Environmental Risk Factors for Adolescent-Onset Substance Use Disorders. *Child and Adolescent Psychiatric Clinics of North America.* 2010;19(3):465-77.
68. Public Health England. Parents with alcohol and drug problems: adult treatment and children and family services. 2021.
69. McGovern R, Smart D, Alderson H, Araújo-Soares V, Brown J, Buykx P, et al. Psychosocial Interventions to Improve Psychological, Social and Physical Wellbeing in Family Members Affected by an Adult Relative's Substance Use: A Systematic Search and Review of the Evidence. *International Journal of Environmental Research and Public Health.* 2021;18(4).
70. Muir C, Adams EA, Evans V, Geijer-Simpson E, Kaner E, Phillips SM, et al. A Systematic Review of Qualitative Studies Exploring Lived Experiences, Perceived Impact, and Coping Strategies of Children and Young People Whose Parents Use Substances. *Trauma, Violence, & Abuse.* 2022:15248380221134297.
71. Kaufman JS, Yang B, Douglas-Palumberi H, Grasso D, Lipschitz D, Houshyar S, et al. Brain-Derived Neurotrophic Factor-5-HTTLPR Gene Interactions and Environmental Modifiers of Depression in Children. *Biological Psychiatry.* 2006;59(8):673-80.
72. Isobel S, Goodyear M, Furness T, Foster KT. Preventing intergenerational trauma transmission: A critical interpretive synthesis. *Journal of Clinical Nursing.* 2019;28(7-8):1100-13.
73. Kallander EK, Weimand B, Ruud T, Becker S, Van Roy B, Hanssen-Bauer K. Outcomes for children who care for a parent with a severe illness or substance abuse. *Child & Youth Services.* 2018;39(4):228-49.
74. Brandon M, Philip G, Clifton J. 'Counting fathers in': Understanding men's experiences of the child protection system. Nuffield Foundation; 2017.
75. Daly M. Parenting support: Another gender-related policy illusion in Europe? *Women's Studies International Forum.* 2013;41:223-30.
76. Daniel B, Taylor J. *Engaging Fathers: Practical Issues for Health and Social Care.* London: Jessica Kingsley; 2001.

77. Davies L. Are young fathers “hard to reach”? Understanding the importance of relationship building and service sustainability. *Journal of Children's Services*. 2016;11(4):317-29.
78. Maxwell N, Scourfield J, Featherstone B, Holland S, Tolman R. Engaging fathers in child welfare services: a narrative review of recent research evidence. *Child & Family Social Work*. 2012;17(2):160-9.
79. Reid C, Greaves L, Poole N. Good, bad, thwarted or addicted? Discourses of substance-using mothers. *Critical Social Policy*. 2008;28(2):211-34.
80. Söderström K, Skårderud F. The good, the bad, and the invisible father: A phenomenological study of fatherhood in men with substance use disorder. *Fathering: A Journal of Theory, Research, and Practice about Men as Fathers*. 2013;11(1):31-51.
81. Guimarães RA, de Oliveira Landgraf de Castro V, do Valle Leone de Oliveira SM, Stabile AC, Motta-Castro ARC, dos Santos Carneiro MA, et al. Gender differences in patterns of drug use and sexual risky behaviour among crack cocaine users in Central Brazil. *BMC Psychiatry*. 2017;17(1):412.
82. McGovern R, Newham J, Addison M, Hickman M, Kaner E. The effectiveness of psychosocial interventions to reduce parental substance use. *Cochrane Database of Systematic Reviews*. 2021(3).
83. McGovern R, Newham J, Addison M, Hickman M, Kaner E. The effectiveness of psychosocial interventions at reducing the frequency of alcohol and drug use in parents: findings of a Cochrane Review and meta-analyses. *Addiction*. 2022;n/a(n/a).
84. Milligan K, Niccols A, Sword W, Thabane L, Henderson J, Smith A, et al. Maternal substance use and integrated treatment programs for women with substance abuse issues and their children: a meta-analysis. *Subst Abuse Treat Prev Policy*. 2010;5:21.
85. Niccols A, Milligan K, Sword W, Thabane L, Henderson J, Smith A. Integrated programs for mothers with substance abuse issues: A systematic review of studies reporting on parenting outcomes. *Harm reduction journal*. 2012;9:14-.
86. Moreland AD, McRae-Clark A. Parenting outcomes of parenting interventions in integrated substance-use treatment programs: A systematic review. *J Subst Abuse Treat*. 2018;89:52-9.
87. Stover CS, Carlson M, Patel S, Manalich R. Where's Dad? The Importance of Integrating Fatherhood and Parenting Programming into Substance Use Treatment for Men. *Child Abuse Rev*. 2018;27(4):280-300.
88. Benoit C, Magnus S. “Depends on the Father” Defining Problematic Paternal Substance Use During Pregnancy and Early Parenthood. *Canadian Journal of Sociology/Cahiers canadiens de sociologie*. 2017;42(4):379-402.
89. Recto P, Lesser J. The Parenting Experiences of Hispanic Adolescent Fathers: A Life Course Theory Perspective. *Western Journal of Nursing Research*. 2020;42(11):918-26.
90. McGovern R, Gilvarry E, Addison M, Alderson H, Geijer-Simpson E, Lingam R, et al. The Association Between Adverse Child Health, Psychological, Educational and Social

Outcomes, and Nondependent Parental Substance: A Rapid Evidence Assessment. *Trauma, Violence, & Abuse*. 2018;21(3):470-83.

91. The Children's Commissioner. Estimating the prevalence of the 'toxic trio', evidence from the Adult Psychiatric Morbidity Survey. London: The Children's Commissioner; 2018.

92. Adjei NK, Schlüter DK, Straatmann VS, Melis G, Fleming KM, McGovern R, et al. Impact of poverty and family adversity on adolescent health: a multi-trajectory analysis using the UK Millennium Cohort Study. *Lancet Reg Health Eur*. 2022;13:100279.

93. Sterne JA, Sutton AJ, Ioannidis JP, Terrin N, Jones DR, Lau J, et al. Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials. *Bmj*. 2011;343:d4002.

94. Sterne J, Egger M, Moher D. Chapter 10: Addressing reporting biases. In: Higgins J, Green S, editors. *Cochrane Handbook for Systematic Reviews of Interventions* 510, Vol 10. [www.cochrane-handbook.org](http://www.cochrane-handbook.org).: Cochrane; 2011.

Accepted Article

Figure 1: flow of studies




Table 1; Summary of included studies estimating impact of maternal substance use upon child outcomes

Study	Child Participants	Parent substance use/ child age at exposure	Child outcome measured/age outcome was assessed	Odds ratio; 95% CI	Adjusted	Converted
<i>Child any alcohol use</i>						
Kelly (2011) Cross sectional study Australian	6843 children (47% boys; 53% girls)	Alcohol abuse (child age 11-14 years)	Any alcohol use (11-14 years)	2.13 [1.44; 3.15]	Yes	Pooled by age and gender
MacLeod (2008) Longitudinal study UK	6895 (49% boys; 51% girls)	Daily alcohol use (exposure anytime in childhood)	Any alcohol use (10 years)	2.80 [0.80; 9.80]	No	No
Haugland (2013) Cross sectional study Norway	5032 children (50% boys; 50% girls)	Alcohol abuse (child age $\geq$ 13 years)	Any alcohol use ( $\geq$ 13 years)	0.99 [0.59; 1.66]	No	Pooled by gender
Oshi (2018) Cross sectional study Barbados	8538 (40% boys; 60% girls)	Daily alcohol use (exposure at point child assessed)	Any alcohol use (majority 15-16 years)	8.39 [4.33; 16.25]	No	No
Malone (2010) Cohort study USA	1383 children	Alcohol dependence (child age 14.8 years)	Any alcohol use (14.8 years)	1.41 [0.97; 2.05]	No	Pooled by age

<i>Child any drug use</i>						
Heron (2013) Longitudinal study UK	6837 children (47% boys; 53% girls)	Substance use (child age between 2 and 33 months)	Cannabis use (16.7 years)	1.70 [1.41; 2.05]	No	No
Malone (2010) Cohort study USA	1383 children	Alcohol dependence (child age 17 years)	Any drug use (17 years)	2.08 [1.29, 3.37]	No	Pooled by age
Korhonen (2008) Longitudinal study Finland	4129 children (48% boys, 52% girls)	Weekly alcohol intoxication (child age 11-12 years)	Ever used drugs (17.6 years)	2.73 [2.01; 3.72]	No	No
<i>Child problem alcohol use</i>						
Alati (2014) Longitudinal study Australia	751 children	Alcohol abuse (child age 13.5 years)	Alcohol intoxication (child age 13.5 years)	2.77 [1.86; 4.13]	Yes	No
Haugland (2013) Cross sectional study Norway	5032 children (50% boys; 50% girls)	Alcohol abuse (child age 13 years)	Alcohol intoxication (13 years)	1.01 [0.39; 2.65]	No	Pooled by gender
Malone (2010) Cohort study USA	1383 children	Alcohol dependence (child age 17 years)	Alcohol dependence (17 years)	2.11 [1.43; 3.10]	No	Pooled by age

McCauley Ohannessian (2004) Cohort study USA	173 children (51% boys; 49% girls) 94% White	Alcohol dependence (child age 15.17 years)	Alcohol dependence (15.17 years)	1.95 [1.27; 3.01]	Yes	No
<i>Child externalising problems</i>						
Malone (2010) Cohort study USA	1383 children 14 years of age 17 years of age	Alcohol dependence (child age 17 yaers)	Conduct disorder (17 years)	1.92 [1.29; 2.86]	No	Pooled by age
McCauley Ohannessian (2004) Cohort study USA	173 children (51% boys; 49% girls) 94% White	Alcohol dependence (child age 15.17 years)	Conduct disorder (15.17 years)	1.28 [0.96; 1.72]	Yes	No
Merikangas (1999) Longitudinal study UK	192 (49% boys; 51% girls) 12.25 years mean age	Substance abuse (child age 12.3 years)	Conduct disorder (12.3 years)	16.70 [1.90; 146.78]	Yes	No
<i>Child internalising problems</i>						
Malone (2010) Cohort study USA	1383 children 14 years of age 17 years of age	Alcohol dependence (child age 17 years)	Depression (17 years)	1.73 [1.14; 2.63]	No	Pooled by age



McCauley Ohannessian (2004) Cohort study USA	173 children (51% boys; 49% girls) 94% White	Alcohol dependence (child age 15.17 years)	Depression (15.17 years)	1.60 [1.17; 2.19]	Yes	No
Merikangas (1999) Longitudinal study UK	192 (49% boys; 51% girls) 12.25 years mean age	Substance abuse (child age 12.3 years)	Conduct disorder (12.3 years)	0.20 [0.02; 2.00]	Yes	No

Table 2; Summary of included studies examining the impact of paternal substance use upon child outcomes

Study	Child Participants	Parent substance use/ child age at exposure	Child outcome measured/age outcome was assessed	Odds ratio; 95% CI	Adjusted	Converted
<i>Any child alcohol use</i>						
Kelly (2011) Cross sectional study Australian	6843 children (47% boys; 53% girls)	Alcohol abuse (child age 11-14 years)	Any alcohol use (11-14 years)	2.21 [1.66; 2.96]	Yes	Pooled by age and gender
MacLeod (2008) Longitudinal study UK	6895 (49% boys; 51% girls)	Daily alcohol use (exposure anytime in childhood)	Any alcohol use (10 years)	0.60 [0.20; 1.80]	No	No

Haugland (2013) Cross sectional study Norway	5032 children (50% boys; 50% girls)	Alcohol abuse (child age $\geq 13$ years)	Any alcohol use ( $\geq 13$ years)	1.29 [0.69; 2.37]	No	Pooled by gender
Oshi (2018) Cross sectional study Barbados	8538 (40% boys; 60% girls)	Daily alcohol use (exposure at point child assessed)	Any alcohol use (majority 15-16 years)	6.03 [4.63; 7.86]	No	No
Malone (2002) Cohort study USA	1121 (49% boys; 51% girls)	Alcohol dependence (child age 14.8 years)	Any alcohol use (14.8 years)	1.53 [1.11; 2.11]	No	No
<i>Any child drug use</i>						
Henry (2017) Longitudinal study USA	442 children 68% black; 17% Hispanic	Cannabis use disorder (lifetime)	Cannabis use (<15 years of age)	4.26 [1.46; 12.43]	No	No
Korhonen (2008) Longitudinal study Finland	4129 children (48% boys, 52% girls)	Weekly alcohol intoxication (child age 11-12 years)	Ever used drugs (17.6 years)	2.39 [1.70; 3.36]	No	No
Malone (2002) Cohort study	1121 (49% boys; 51% girls)	Alcohol dependence (child age 14.8 years)	Any drug use (14.8 years)	1.67 [1.11; 2.51]	No	No

This article is protected by copyright. All rights reserved.

USA						
<i>Child problem alcohol use</i>						
Alati (2014) Longitudinal study Australia	751 children	Alcohol abuse Professional fathers and semi-skilled or housemaker mothers (child age 13.5 years)	Alcohol intoxication (13.5 years of age)	1.40 [1.04; 1.88]	Yes	No
Haugland (2013) Cross sectional study Norway	5032 children (50% boys; 50% girls)	Alcohol abuse (child age $\geq 13$ years of age)	Alcohol intoxication ( $\geq 13$ years of age)	1.35 [1.06; 1.73]	No	Pooled by gender
Rehorcikova (2013) Cross section study Slovakia	2494 (52% boys; 48% girls)	Daily alcohol use (child median age 13 years)	Weekly alcohol use consumption (median 13 years)	2.23 [1.19; 4.18]	Yes	No
Jennison (2014) Longitudinal USA	4648	Alcohol dependent (child age 16.3 years)	Risky alcohol use (16.3 years)	2.79 [1.72; 4.53]	Yes	No
McCauley Ohannessian (2004) Cohort study USA	173 children (51% boys; 49% girls) 94% White	Alcohol dependence (child age 15.2 years)	Alcohol dependence (15.2 years)	1.57 [1.02; 2.41]	Yes	No

This article is protected by copyright. All rights reserved.

Malone (2002) Cohort study USA	1121 (49% boys; 51% girls)	Alcohol dependence (child age 14.8 years)	Alcohol dependence (17 years)	1.93 [1.29; 2.89]	No	No
<i>Child externalising problems</i>						
Carbonneau (1998) Cross sectional study Canada	642 children	Alcohol dependency (child age 12 years)	Oppositional problems (12 years)	1.94 [1.30; 2.88]	No	Converted from means and standard deviations
Cumes-Rayner (1992) Cross sectional study Australia	260 boys	Alcohol abuse (child age 15 years)	Anti-social behaviour (17 years)	1.98 [0.92; 4.26]	No	Converted from means and standard deviations
Fals-Stewart (2004) Cohort study USA	120 children 68% White, 20% African American, 8% Hispanic; 5% other 8-12 years of age	Alcohol abuse, drug abuse & drug dependence (child age 9.1 years)	Externalising problems (9.1 years)	1.80 [0.77; 4.22]	No	Converted from means and standard deviations

Malone (2002) Cohort study USA	1121 (49% boys; 51% girls)	Alcohol dependence (child age 14.8 years)	Conduct disorder (17 years)	2.40 [0.92; 6.26]	No	Pooled by gender
McCauley Ohannessian (2004) Cohort study USA	173 children (51% boys; 49% girls) 15.2 years mean age 94% White	Alcohol dependence (child age 15.17 years)	Conduct disorder (15.17 years)	1.19 [0.87; 1.62]	Yes	No
Merikangas (1999) Cohort study USA	192 (49% boys; 51% girls) 12.25 years mean age	Substance abuse (child age 12.3 years)	Conduct disorder (12.3 years)	0.70 [0.10; 4.90]	Yes	No
<i>Child internalising problems</i>						
Carbonneau (1998) Cross sectional study Canada	642 children 12 years	Alcohol dependency (child age 12 years)	Anxiety problems (12 years)	1.18 [0.79; 1.75]	No	Converted from means and standard deviations
Fals-Stewart (2004) Cohort study USA	120 children 68% White, 20% African American, 8% Hispanic; 5% other	Alcohol abuse, drug abuse & drug dependence (child age 9.1 years)	Internalising problems (9.1 years)	2.92 [1.23; 6.94]	No	Converted from means and standard deviations

This article is protected by copyright. All rights reserved.

	8-12 years of age					
McCauley Ohannessian (2004) Cohort study USA	173 children (51% boys; 49% girls) 15.2 years mean age 94% White	Alcohol dependence (child age 15.17 years)	Depression (15.17 years)	1.36 [0.95; 1.92]	Yes	No
Malone (2002) Cohort study USA	1121 (49% boys; 51% girls)	Alcohol dependence (child age 14.8 years)	Depression (17 years)	1.76 [1.17; 2.65]	No	No
Merikangas (1999) Cohort study USA	192 (49% boys; 51% girls) 12.25 years mean age	Substance abuse (child age 12.3 years)	Anxiety disorder (12.3 years)	0.40 [0.10; 1.60]	Yes	No

Table 3; Results of meta-analyses

This article is protected by copyright. All rights reserved.

<b>Analysis</b>	<b>Result</b>	<b>Studies and Participants</b>
<b>Child substance use</b>		
1.1 Fathers substance use and child any alcohol use	OR: 1.89, 95% CI 0.93 to 3.83	5 studies; 28,429 participants
1.2 Fathers substance use and child drug use	OR: 2.86, 95% CI 1.25 to 6.54	3 studies; 5,692 participants
1.3 Fathers substance use and child problematic alcohol use	OR; 1.70, 95% CI 1.36 to 2.12	6 studies; 14,219 participants
2.1 Mothers substance use and child any alcohol use	OR: 2.26, 95% CI 1.08 to 4.70	5 studies; 28,691 participants
2.2 Mothers substance use and child drug use	OR: 2.09, 95% CI 1.53 to 2.86	3 studies; 12,349 participants
2.3 Mothers substance use and child problematic alcohol use	OR: 2.16, 95% CI 1.73 to 2.71	4 studies; 7,339 participants
<b>Child externalising problems</b>		
3.1 Fathers substance use and child externalising problems	OR: 1.60, 95% CI 1.18 to 2.17	6 studies; 2,508 participants
3.2 Mothers substance use and child externalising problems	OR: 1.81, 95% CI 1.01 to 3.22	3 studies; 1,748 participants
<b>Child internalising problems</b>		
4.1 Fathers substance use and child internalising problems	OR: 1.42, 95% CI 1.12 to 1.81	5 studies; 2248 participants
4.2 Mothers substance use and child internalising problems	OR: 1.60, 95% CI 1.25 to 2.06	3 studies; 1,748 participants

Table 4; Comparing the goodness of fit for models with a study size (variance) covariate with models without covariates



Outcomes	Covariate for small-study effect (95% CrI)	OR (95% CrI): parental exposure of substance use	Total residual deviance	Between-study SD (95% CrI)	DIC
Child alcohol use-covariate model	Beta=1.29 (-7.17 to 4.63)	Father: 1.80 (0.63 to 4.76) Mother: 2.27 (0.85 to 6.42)	Mean=10.23	0.92 (0.50 to 1.98)	9.53
Child alcohol use-no covariates	-	Father: 1.90 (0.73 to 4.57) Mother: 2.25 (0.90 to 5.81)	Mean=10.11	0.86 (0.49 to 1.76)	9.18
Child alcohol problematic use-covariate model	Beta=-0.89 (-6.72 to 4.15)	Father: 1.67(1.23 to 2.34) Mother: 2.01 (1.20 to 3.13)	Mean=11.16	0.22 (0.02 to 0.64)	7.32
Child alcohol problematic use-without covariate	-	Father: 1.68 (1.32 to 2.27) Mother: 2.10 (1.43 to 2.94)	Mean=11.1	0.19 (0.01 to 0.55)	6.12
Child drug use- covariate model	Beta=0.16 (-5.32 to 5.75)	Father: 2.23 (0.91 to 6.17) Mother: 1.90 (0.85 to 3.78)	Mean=8.03	0.39 (0.06 to 1.60)	6.34
Child drug use- without covariate	-	Father: 2.25 (1.25 to 4.71) Mother: 1.93 (1.07 to 3.16)	Mean=7.74	0.32 (0.05 to 1.14)	5.50

This article is protected by copyright. All rights reserved.

<b>Child externalising problems-covariate model</b>	Beta=0.94 (-0.56 to 2.39)	Fathers: 1.95 (1.13 to 3.22) Mothers: 2.14 (1.14 to 4.53)	Mean=10.2	0.26 (0.02 to 1.02)	5.96
<b>Child externalising problems-no covariates</b>	-	Fathers: 1.62 (1.03 to 2.61) Mothers: 1.70 (1.02 to 4.01)	Mean=10.46	0.30 (0.03 to 1.03)	5.45
<b>Child internalising problems-covariate model</b>	Beta=-1.51 (-3.21 to 0.13)	Fathers: 1.04 (0.61 to 1.80) Mothers: 1.14 (0.58 to 2.16)	Mean=8.91	0.23 (0.01 to 1.06)	5.59
<b>Child internalising problems-no covariates</b>	-	Fathers: 1.42 (0.83 to 2.27) Mothers: 1.54 (0.63 to 2.61)	Mean=11.28	0.23 (0.01 to 1.27)	4.93

OR=odds ratio CrI=credible interval SD=standard deviation DIC=deviance information criterion

This article is protected by copyright. All rights reserved.